

Type 2 Diabetes - Chronic Kidney Disease

CKD is eGFR < 60ml/min or kidney damage for ≥ 3 months (e.g. urine sediment, abnormal imaging, or albuminuria (UACR < 30mg/g = nl, 30-300 = micro, >300 = macro))

Stages of Chronic Kidney Disease (CKD)

	1	2	3	4	5
eGFR	> 60	> 60	30-59	15-29	< 15 ml/min/1.73m ²

Markers of progression: decreasing eGFR, increasing albuminuria, poor BP control

Workup of CKD and to r/o non-diabetes causes

CMP, UA, UACR, Uric Acid, Phos, CBC, ANA, RF, C3, C4, HepB sAg, HepC Ab, dilated retinal exam, and renal U/S; if pat >40 yrs & UACR is pos then check SPEP and UPEP

Referrals

Nephrologist: When eGFR < 30 or sooner if unsure of etiology or problems
 Nutrition: Refer to RD for consult (protein, Na+, K+, PO4, fluids, saturated fat)

Managing Complications of CKD – Stages 3-5

Acidosis		
If CO2 < 22mmol/L	Start sodium bicarbonate 325-650mg (1-2 tabs) TID-QID	Goal: CO2 ≥ 22mmol/L
Anemia		
<p>Check Hb at least yearly: Anemia = Hb <13.5 g/dL adult men, <12 g/dL adult women; r/o B12/folate deficiency, GI blood loss, other causes</p> <p>Baseline Labs: Ferritin, transferrin % sat, iron studies (Fe, % Sat, TIBC), CBC+diff</p> <p>Start oral iron therapy if ferritin/iron studies low</p> <p>Ferrous Sulfate (FeSO4) 325mg daily to TID</p> <p>Consider docusate 100mg BID to reduce constipation</p> <p>Monitor ferritin to avoid iron overload</p> <p>Consider IV iron or blood transfusion if needed</p> <p>Safety of erythropoiesis stimulating agents (ESA) unclear; reserve for patients on dialysis, pending renal transplant, or Hb < 9 with symptoms unresponsive to treatment above</p>		
Blood Pressure		
Most effective CKD intervention: BP goal <130/80; continue ACEI/ARB (watch K+)		
Cardiovascular Disease (CVD)		
<p>CVD: CKD increases CVD risk – patients on aspirin (if no contraindications)</p> <p>Achieve lipid targets, encourage tobacco cessation</p>		
Diabetes		
<p>Blood sugar control—as renal fxn declines pts' BGs often improve—titrate meds down as needed; Caution setting an A1c target <7% if advanced CKD or CVD</p> <p>D/C metformin when Creatinine >1.5 men or >1.4 women</p> <p>Peripheral Neuropathy: Foot ulcers common, check feet each visit, refer to shoe clinic</p> <p>Retinopathy: Ophth/retinal visits regularly</p> <p>Autonomic Neuropathy: Frequent BP fluctuations, including orthostatic symptoms.</p>		

Type 2 Diabetes - Chronic Kidney Disease

Edema/Fluid Overload

Establish patient's dry wt; Titrate furosemide 20-240mg BID (diuresis lasts 6 hours-give AM & mid-day)

Metabolic Bone Disease

Evidence Based: Phosphorus (PO₄): if >4.6 mg/dL, start binder (calcium); Refer to RD for dietary PO₄ restriction

Calcium (Ca): If <8.4, start/increase calcium supplementation; target: 8.4-9.5 mg/dL

If >10.2, correct causes (often 2° meds), need to hold Ca and/or Vit D/calcitriol

Consensus Opinion: If iPTH elevated, measure 25(OH) Vitamin D; If 25(OH)D >=30mg/mL, start calcitriol

If 25(OH) Vitamin D <30mg/mL, start ergocalciferol (Vitamin D2)

Follow Ca, PO₄, iPTH, and 25(OH)D (Vitamin D): if Ca or PO₄ above target or if iPTH below target, hold calcitriol and/or calcium

CKD Stage	eGFR	iPTH goal	PO ₄ Goal	Ca Goal	Ca goal Ca x PO ₄
3	30-59	35-70	2.7-4.6	8.4-9.5	< 55
4	15-29	70-110	2.7-4.6	8.4-9.5	< 55
5	< 15	150-300	3.5-5.4	8.4-9.5	< 55

Medication*	iPTH effect	PO ₄ effect	Ca effect	Comments
Phosphate Binders				
<i>CaCO₃ (Oyst-Cal or TUMS) 500-2000mg with meals</i>	–	↓	↑	Use if Ca < 8.4; No more than 7g/d
<i>Ca Acetate 1334-2868mg with meals</i>	–	↓↓	↑	Use if Ca < 8.4 & PO ₄ > 5
<i>Sevelamer (Renagel) 800-1600mg TID</i>	–	↓↓		Decrease PO ₄ , no effect on Ca; cost
<i>Lanthanum 1500-3750mg/day w/ meals</i>	–	↓↓	↓	Decrease PO ₄ and Ca; cost
<i>Aluminum 600-1200mg TID between meals & HS</i>	–	↓↓	–	ONLY if PO ₄ > 7 and Ca x PO ₄ > 55; not more than 30 days (toxicity)
Vitamin D and Analogs				
<i>Vit D2 (Ergocalciferol) 1.25-5mg daily</i>	↓	–	↑↑	Use if Vit D < 30 mg/mL
<i>Calcitriol 0.25-1mcg daily or 0.5-3mcg TIW</i>	↓	–	↑↑	Use only if Ca & PO ₄ in normal range
<i>Doxercalciferol 1-3mcg daily or 10-20mcg TIW</i>	↓	–	↑	Hold if Ca x PO ₄ > 55
Other				
<i>Cinacalcet 30-180mg daily</i>	↓	↓	↓↓	Do not use if Ca < 8.4

*Always include dietary phosphorous restriction

Drugs in italics are not on the IHS National Core Formulary

Lab Monitoring

Parameter	GFR > 60	GFR 30-59	GFR 15-29	GFR<15 not on Dialysis
Creatinine and eGFR	Annual	Each visit	Each visit	Each visit
UACR	Annual	Q3-6mos*	Each visit*	Each visit*
Hb	Annual	Q3mos	Q3mos	Q3mos
Fe, Transferrin Sat, Ferritin		Q3mos	Q3mos	Q3mos
Ca, PO ₄ , and iPTH		At least annually	Q3mos	Q3mos

Monitor more often if values are worsening or on medications that affect these labs

***Frequency of checking depends on rate of rise of urine albumin**

Ref: KDOQI/NKF and UK Renal Assoc 4th Ed. Clinical Practice Guidelines for Complications of CKD
 ADA Clinical Practice Recommendations 2010. J Am Soc Nephrol 2010; 21:2-6.