

Standards of Care and Clinical Practice
Recommendations:
Type 2 Diabetes

Bibliography

Updated December 2011

Indian Health Service
Division of Diabetes Treatment and Prevention
5300 Homestead Road NE
Albuquerque, New Mexico 87110
<http://www.ihs.gov/MedicalPrograms/Diabetes/>



TABLE OF CONTENTS

Introduction	4
Alcohol and Other Substance Use	5
Antiplatelet Therapy	5
Bariatric Surgery.....	6
Blood Pressure Management	6
Cancer Screening	7
Cardiovascular Care	7
Contraception	8
Depression	8
Diagnosis of Type 2 Diabetes and Prediabetes.....	9
Diabetes Self-Management Education	10
Eye Care	10
Foot Care.....	11
Glycemic Control.....	12
Immunizations	13
Kidney Care	13
Lipid Management	15
Medical Nutrition Therapy.....	16
Neuropathy	17
Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis	18
Obstructive Sleep Apnea.....	18
Older Adults	18
Oral Care	19
Peripheral Arterial Disease	19
Physical Activity	19
Preconception, Pregnancy, and Postpartum	20

Sexual Health	21
Tobacco Use	21
Tuberculosis Screening	22
Weight Management, Overweight, and Obesity.....	22
Youth and Type 2 Diabetes	23

Bibliography

This bibliography contains journal articles and reports that provide more in-depth information on the topics addressed in the *Standards of Care and Clinical Practice Recommendations: Type 2 Diabetes*. Additional references can be found in the *Indian Health Diabetes Best Practices* documents.

Introduction

Acton KJ, Burrows NR, Moore K, Querec L, Geiss LS, Engelgau MM. Trends in diabetes prevalence among American Indian and Alaska Native children, adolescents, and young adults. *Am J Public Health*. 2002 Sep;92(9):1485-90.

American Diabetes Association. Standards of medical care in diabetes,2011. 2011 Jan;34 Suppl 1:S11-61.

Dabelea D, Pettitt DJ. Intrauterine diabetic environment confers risks for type 2 diabetes mellitus and obesity in the offspring, in addition to genetic susceptibility. *J Pediatr Endocrinol Metab*. 2001;14:1085-91.

Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the Adverse Childhood Experiences (ACE) Study. *Am J Prev Med*.1998 May;14(4):245-58.

Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med*. 2008 Jul 3;359:61-73.

Hillier TA, Pedula KL, Schmidt MM, Mullen JA, Charles MA, Pettitt DJ. Childhood obesity and metabolic imprinting: the ongoing effects of maternal hyperglycemia. *Diabetes Care*. 2007;30:2287-92.

Krieger, N. *Epidemiology and the people's health: theory and context*. New York: Oxford University Press; 2011.

Mayfield JA, Rith-Najarian SJ, Acton KJ, Schraer CD, Stahn RM, Johnson MH, et al. Assessment of diabetes care by medical record review: the Indian Health Service model. *Diabetes Care*. 1994 Aug; 17(8):918-23.

National Committee for Quality Assurance. *Comprehensive diabetes care*. In: *The state of health care quality reform: the quality agenda and resource use*. Washington (DC): National Committee for Quality Assurance; 2010. p. 47-51.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Alcohol and Other Substance Use

Bantle JP, Wylie-Rosett J, Albright AL, Apovian CM, Clark NG, Franz MJ, et al. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. *Diabetes Care*. 2008;31 Suppl 1:S61-78. (See p. S67 for alcohol recommendations)

Boyd DR, Milman K, Stuart P, Dekker A, Flaherty J. Alcohol screening and brief intervention (ASBI) program implementation and operations manual. Rockville (MD): Indian Health Service, Office of Clinical and Preventive Services, Emergency Services; 2008 Mar. 36 p.

Guard A, Rosenblum L. Alcohol screening and brief intervention: a guide for public health practitioners. Produced by the American Public Health Association under contract to the National Highway Traffic Safety Administration, U.S. Department of Transportation. Washington (DC): American Public Health Association and Education Development Center, Inc.; 2008. 28 p.

Indian Health Service National Tribal Advisory Committee on Behavioral Health (NTAC) and the Behavioral Health Work Group. American Indian/Alaska Native behavioral health briefing book. Rockville (MD): U.S. Department of Health and Human Services, Indian Health Service, Division of Behavioral Health, Office of Clinical and Preventive Services; 2011 Aug.

Substance Abuse and Mental Health Services Administration, Office of Applied Studies. The national survey on drug use and health (NSDUH) report: substance use among American Indian or Alaska Native adults. Rockville (MD); 2010 Jun 24.

Antiplatelet Therapy

Belch J, MacCuish A, Campbell I, Cobbe S, Taylor R, Prescott R, et al. The prevention of progression of arterial disease and diabetes (POPADAD) trial: factorial randomised placebo controlled trial of aspirin and antioxidants in patients with diabetes and asymptomatic peripheral arterial disease. *BMJ*. 2008;337:a1840.

Pignone M, Alberts MJ, Colwell JA, Cushman M, Inzucchi SE, Mukherjee D, et al.; American Diabetes Association; American Heart Association; American College of Cardiology Foundation. Aspirin for primary prevention of cardiovascular events in people with diabetes: a position statement of the American Diabetes Association, a scientific statement of the American Heart Association, and an expert consensus document of the American College of Cardiology Foundation. *Diabetes Care*. 2010;33:1395-1402.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Bariatric Surgery

Buchwald H, Estok R, Fahrback K, Banel D, Jensen MD, Pories WJ, et al. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *Am J Med.* 2009;122:248-56.

Dixon JB, Zimmet P, Alberti KGMM, Rubino F; IDF Taskforce on Epidemiology and Prevention. Bariatric surgery: an IDF statement for obese type 2 diabetes. *Diabet Med.* 2011;28(6):628-42.

Poirier P, Cornier M, Mazzone T, Stiles S, Cummings S, Klein S, et al. Bariatric surgery and cardiovascular risk factors: a scientific statement from the American Heart Association. *Circulation.* 2011. Epub 2011 Mar 14.

Blood Pressure Management

Action to Control Cardiovascular Risk in Diabetes (ACCORD) Study Group. Effects of intensive blood-pressure control in type 2 diabetes. *N Eng J Med.* 2010 Apr 29;362:1575-85.

Holman RR, Paul SK, Bethel MA, Neil AW, Matthews DR. Long-term follow-up after tight control of blood pressure in type 2 diabetes. *N Eng J Med.* 2008;359(15):1565-76.

Indian Health Service diabetes care and outcomes audit, unpublished data. Albuquerque (NM): Department of Health and Human Services (US), Indian Health Service, Office of Clinical and Preventive Services, Division of Diabetes Treatment and Prevention; 2011.

National High Blood Pressure Education Program. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. Bethesda (MD): Department of Health and Human Services (US), National Institutes of Health, National Heart, Lung, and Blood Institute; 2004 Aug. 88 p. (NIH Publication No. 04-5230).

Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al.; the DASH-Sodium Collaborative Research Group. Effects on blood pressure of reduced dietary sodium and the dietary approaches to stop hypertension (DASH) diet. *N Engl J Med.* 2001 Jan 4;344(1):3-10.

United Kingdom Prospective Diabetes Study (UKPDS) Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS. *BMJ.* 1998;317:703-13.

Zhang Y, Galloway JM, Welty TK, Wiebers DO, Whisnant JP, Devereaux RB, et al. Incidence and risk factors for stroke in American Indians: the Strong Heart Study. *Circulation.* 2008;118:1577-84.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Cancer Screening

Giovannucci E, Harlan DM, Archer DM, Bergenstal RM, Gapstur SM, Habel LA, et al. Diabetes and cancer: a consensus report. *Diabetes Care*. 2010;33:1674-85.

Smith RA, Cokkinides V, Brooks D, Saslow D, Shah M, Brawley OW. Cancer screening in the United States, 2011: a review of current American Cancer Society guidelines and issues in cancer screening. *CA Cancer J Clin*. 2011;61(1):8-30.

Wright TC, Massad LS, Dunton CJ, Spitzer M, Wilkinson EJ, Solomon D; 2006 American Society for Colposcopy and Cervical Pathology-sponsored consensus conference. American Society for Colposcopy and Cervical Pathology (ASCCP) cervical consensus guidelines and algorithms. *Am J Obstet Gyn*. 2007;197(4):346-55.

Cardiovascular Care

Action to Control Cardiovascular Risk in Diabetes (ACCORD) Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med*. 2008;358(24):2545-59.

ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008;358(24):2560-72.

Buse JB, Ginsberg HN, Bakris GL, Clark NG, Costa F, Eckel R, et al. AHA/ADA scientific statement: primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Circulation*. 2007;115:114-26.

Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med*. 2009 Jan 8;360:129-39.

Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, 2010. *Circulation*. 2010 Nov 12;122:e584-e636.

Lee ET, Howard BV, Wang W, Welty TK, Galloway JM, Best LG, et al. Prediction of coronary heart disease in a population with high prevalence of diabetes and albuminuria: the Strong Heart Study. *Circulation*. 2006;113(25):2897-2905.

Skyler JS, Bergenstal R, Bonow RO, Buse J, Deedwania P, Gale EAM, et al. Intensive glycemic control and the prevention of cardiovascular events: implications of the ACCORD, ADVANCE, and VA Diabetes Trials: a position statement of the American Diabetes Association and a scientific statement of the American College of Cardiology Foundation and the American Heart Association. *J Am Coll Cardiol*. 2009; 53:298-304.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Wilson PWF, Meigs JB, Sullivan L, Fox CS, Nathan DM, D'Agostino RB. Prediction of incipient diabetes mellitus in middle-aged adults: the Framingham Offspring Study. *Arch Intern Med.* 2007;167:1068-74.

Contraception

Tepper NK, Curtis KM, Jamieson DJ, Marchbanks PA. Update to CDC's U.S. medical eligibility criteria for contraceptive use, 2010: revised recommendations for the use of contraceptive methods during the postpartum period. *Morb Mortal Wkly Rep.* 2011 Jul 8;60(26):878-83. Corrected and republished from: U.S. medical eligibility criteria for contraceptive use, 2010: adapted from the World Health Organization medical eligibility criteria for contraceptive use, 4th edition. *Morb Mortal Wkly Rep.* 2010 Jun 18;59(No. RR04):1-6.

Depression

Beck, AT, Steer, RA, Brown, GK. *Manual for the Beck Depression Inventory–II.* San Antonio (TX): Psychological Corporation; 1996.

Ell K, Katon W, Xie B, Lee P-J, Kapetanovic S, Guterman J, et al. Collaborative care management of major depression among low-income, predominantly Hispanic subjects with diabetes. *Diabetes Care.* 2010 Apr;33(4):706-13.

Fisher L, Glasgow RE, Mullan JT, Skaff MM, Polonsky WH. Development of a brief diabetes distress screening instrument. *Ann Fam Med.* 2008 May/Jun;6(3):246-52.

Fisher L, Skaff MM, Mullan JT, Arean P, Mohr D, Masharani U, et al. Clinical depression versus distress among patients with type 2 diabetes: not just a question of semantics. *Diabetes Care.* 2007 Mar;30(3):542-8.

Gonzalez JS, Safren SA, Cagliero E, Wexler DJ, Delahanty L, Wittenberg E, et al. Depression, self-care, and medication adherence in type 2 diabetes: relationships across the full range of symptom severity. *Diabetes Care.* 2007 Sep;30(9):2222–7.

Jiang L, Beals J, Whitesell NR, Roubideaux Y, Manson SM; the AISUPERPFP Team. Stress burden and diabetes in two American Indian reservation communities. *Diabetes Care.* 2008 Mar;31(3):427-9.

Katon WJ, Lin EHB, Von Korff M, Ciechanowski P, Ludman EJ, Young B, et al. Collaborative care for patients with depression and chronic illnesses. *N Engl J Med.* 2010 Dec 30;363(27):2611-20.

Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care.* 2003 Nov;41(11):1284-92.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Li C, Ford ES, Strine TW, Mokdad AH. Prevalence of depression among U.S. adults with diabetes: findings from the 2006 Behavioral Risk Factor Surveillance System. *Diabetes Care*. 2008 Jan;31(1): 105-7.

Lin EHB, Rutter CM, Katon W, Heckbert SR, Ciechanowski P, Oliver MM, et al. Depression and advanced complications of diabetes: a prospective cohort study. *Diabetes Care*. 2010 Feb;33(2):264-9.

Manson SM, Beals J, Klein SA, Croy CD; AISUPERPFP Team. Social epidemiology of trauma among 2 American Indian reservation populations. *Am J Public Health*. 2005;95:851-9.

Pan A, Lucas M, Sun Q, van Dam RM, Franco OH, Manson JE, et al. Bidirectional association between depression and type 2 diabetes mellitus in women. *Arch Intern Med*. 2010;170:1884-91.

Sahota PKC, Knowler WC, Looker HC. Depression, diabetes, and glycemic control in an American Indian community. *J Clin Psychiatry*. 2008;69:800-9.

Singh PK, Looker HC, Hanson RL, Kraffoff J, Bennett PH, Knowler WC. Depression, diabetes, and glycemic control in Pima Indians. *Diabetes Care*. 2004;27(2):618-9.

Diagnosis of Type 2 Diabetes and Prediabetes

American Diabetes Association. Diagnosis and classification of diabetes mellitus: position statement. *Diabetes Care*. 2011 Jan;34 Suppl 1:S62-9.

American Diabetes Association. Standards of medical care in diabetes—2011. *Diabetes Care*. 2011 Jan;34 Suppl 1:S11-61.

Endocrine Society Statement on the use of A1C for diabetes diagnosis and risk estimation. Chevy Chase (MD); January 2010.

Goldstein DE, Little RR, Lorenz RA, Malone JI, Nathan D, Peterson CM, et al. Tests of glycemia in diabetes. *Diabetes Care*. 2004;27(7):1761-73.

Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D, Heine RJ; A1C-Derived Average Glucose (ADAG) Study Group. Translating the A1C assay into estimated average glucose values. *Diabetes Care*. 2008;31(8):1473-8.

Pogach L, Conlin PR, Hobbs C, Vigersky RA, Aron D. VA-DoD update of diabetes guidelines: what clinicians need to know about absolute risk of benefits and harms and A1C laboratory accuracy. *Fed Pract*. 2011;28(4):39-44.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

US Department of Veterans Affairs Office of Quality and Performance; US Army Medical Command Quality Management Division. VA/DoD clinical practice guideline for the management of diabetes mellitus. Version 4.0. Washington (DC): Veterans Health Administration and Department of Defense; 2010. 146 p.

Diabetes Self-Management Education

American Association of Diabetes Educators. AADE position statement: individualization of diabetes self-management education. *Diabetes Educ*. 2007 Jan-Feb;33(1):45-9.

American Association of Diabetes Educators. AADE7 self-care behaviors. *Diabetes Educ*. 2008;34:445-9.

American Association of Diabetes Educators. The art and science of diabetes self-management education desk reference. Mensing C, editor. 2nd ed. Chicago (IL): American Association of Diabetes Educators; 2010.

Funnell MM, Brown TL, Childs BP, Haas LB, Hoseney GM, Jensen B, et al. National standards for diabetes self-management education. *Diabetes Care*. 2011; 34 Suppl 1:S89-96.

Funnell MM, Tang TS, Anderson RM. From DSME to DSMS: developing empowerment-based diabetes self-management support. *Diabetes Spectrum*. 2007;20(4):221-6.

Eye Care

American Academy of Ophthalmology Retina Panel. Preferred practice pattern: diabetic retinopathy. San Francisco (CA): American Academy of Ophthalmology; 2008.

American Optometric Association. Optometric clinical practice guideline care of the patient with diabetes mellitus: reference guide for clinicians. St. Louis (MO): American Optometric Association;2009. 74 p.

Carroll M, Cullen T, Ferguson S, Hogge N, Horton M, Kokesh J. Innovation in Indian healthcare: using health information technology to achieve health equity for American Indian and Alaska Native populations. *Perspect Health Inf Manag*. (2011 Winter):1-9.

Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Research Group. Prolonged effect of intensive therapy on the risk of retinopathy complications in patients with type 1 diabetes mellitus 10 years after the Diabetes Control and Complications Trial. *Arch Ophthalmol*. 2008;126:1707-15.

Fong DS, Aiello LP, Ferris FL 3rd, Klein R. Diabetic retinopathy: technical review. *Diabetes Care*. 2004;27(10):2540-53.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Gómez-Ulla F, Fernandez MI, Gonzalez F, Rey P, Rodriguez M, Rodriguez-Cid MJ, et al. Digital retinal images and teleophthalmology for detecting and grading diabetic retinopathy. *Diabetes Care*. 2002;25(8):1384-9.

Wilson C, Horton M, Cavallerano J, Aiello LM. Addition of primary care-based retinal imaging technology to an existing eye care professional referral program increased the rate of surveillance and treatment of diabetic retinopathy. *Diabetes Care*. 2005; 28:318-22.

Foot Care

Boulton AM, Armstrong DG, Albert SF, Frykberg RG, Hellman R, Kirkman MS, et al. Comprehensive foot examination and risk assessment: a report of the Task Force of the Foot Care Interest Group of the American Diabetes Association, with endorsement by the American Association of Clinical Endocrinologists. *Diabetes Care*. 2008;31:1679-85.

Lavery LA, Peters EJ, Williams JR, Murdoch DP, Hudson A, Lavery DC. Reevaluating the way we classify the diabetic foot: restructuring the diabetic foot risk classification system of the International Working Group on the Diabetic Foot. *Diabetes Care*. 2008 Jan;31(1):154-6.

Mayfield JA, Reiber GE, Sanders LJ, Pogach LM. Preventive foot care in diabetes: American Diabetes Association position statement. *Diabetes Care*. 2004 Jan; 27 Suppl 1:S63-4.

Reiber GE, Raugi GJ. Preventing foot ulcers and amputations in people with diabetes: future promise based on lessons learned. *Lancet*. 2005 Nov 12;366(9498):1676-7.

Rith-Najarian S, Branchaud C, Beaulieu O, Gohdes D, Simonson G, Mazze R. Reducing lower-extremity amputations due to diabetes: application of the staged diabetes management approach in a primary care setting. *J Fam Pract*. 1998;47(2):127-32.

Rogers LC, Frykberg RG, Armstrong DG, Boulton AJM, Edmonds M, Ha Van G, et al. The Charcot foot in diabetes: consensus report of the American Diabetes Association and the American Podiatric Medical Association. *Diabetes Care*. 2011 Sep;34:2123-9.

Schraer CD, Weaver D, Naylor JL, Provost E, Mayer AM. Reduction of amputation rates among Alaska Natives with diabetes following the development of a high risk foot program. *Int J Circumpolar Health*. 2004;63 Suppl 2:114-9.

Sumpio BE, Armstrong DG, Lavery LA, Andros G. The role of interdisciplinary team approach in the management of the diabetic foot: a joint statement from the Society for Vascular Surgery and the American Podiatric Medical Association. *J Am Podiatr Med Assoc*. 2010 Jul 1;100(4):309-11.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Glycemic Control

Action to Control Cardiovascular Risk in Diabetes (ACCORD) Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med*. 2008;358(24):2545-59.

ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008;358(24):2560-72.

American Diabetes Association. Position Statement Executive Summary: Guidelines and Recommendations for Laboratory Analysis in the Diagnosis and Management of Diabetes Mellitus. *Diabetes Care*. 2011 Jun;34(6):1419-23.

Boussageon R, Bejan-Angoulvant T, Saadatian-Elahi M, Lafont S, Kassaï B, Erpeldinger S, et al. Effect of intensive glucose lowering treatment on all-cause mortality, cardiovascular death, and microvascular events in type 2 diabetes: meta-analysis of randomised controlled trials. *BMJ*. 2011;343:d5301.

Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med*. 2009 Jan 8;360(2):129-39.

Hemmingsen B, Lund SS, Gludd C, Vaag A, Almdal T, Hemmingsen C. Intensive glycaemic control for patients with type 2 diabetes: systematic review with meta-analysis and trial sequential analysis of randomised clinical trials. *BMJ*. 2011;343:d6898.

Holman RR, Paul SK, Bethel MA, Matthews DR, Neil AW. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Eng J Med*. 2008 Oct 9;359(15):1577-89.

Kuritzky L. Managing type 2 diabetes in the primary care setting: beyond gluco-centricity. *Am J Med Sci*. 2010;340:133-43.

Nathan DM, Buse JB, Davidson MB, Ferrannini E, Holman RR, Sherwin R, et al. Medical management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*. 2009;32(1):193-203.

O'Connor PJ, Bodkin NL, Fradkin J, Glasgow RE, Greenfield S, Gregg E, et al. Diabetes performance measures: current status and future directions. American Diabetes Association Consensus Report. *Diabetes Care*. 2011 Jul;34(7):1651-9.

Patel A, MacMahon S, Chalmers J, Neal B, Billot L, Woodward M, et al.; ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008 Jun 12;358:2560-72.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Preiss D, Ray KK. Intensive glucose lowering treatment in type 2 diabetes: editorial. *BMJ*. 2011;343:d4243.

Rodbard HW, Jellinger PS, Davidson JA, Einhorn D, Garber AJ, Grunberger G, et al. Statement by an American Association of Clinical Endocrinologists/American College of Endocrinology consensus panel on type 2 diabetes mellitus: an algorithm for glycemic control. *Endocr Pract*. 2009;15(6):540-59. Erratum in *Endocr Pract*. 2009 Nov-Dec;15(7):768-70.

Skyler JS, Bergenstal R, Bonow RO, Buse J, Deedwania P, Gale EA, et al. Intensive glycemic control and the prevention of cardiovascular events: implications of the ACCORD, ADVANCE, and VA diabetes trials: a position statement of the American Diabetes Association and a scientific statement of the American College of Cardiology Foundation and the American Heart Association. *Diabetes Care*. 2009 Jan;32(1):187-92. Epub 2008 Dec 17.

Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, et al.; United Kingdom Prospective Diabetes Study Group. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*. 2000 Aug 12;321(7258):405-12.

Immunizations

Centers for Disease Control and Prevention. Recommended adult immunization schedule—United States, 2011. *Morb Mort Wkly Rep*. 2011 Feb 4;60(40):1-4.

Centers for Disease Control and Prevention. Recommended immunization schedules for persons aged 0-18 years—United States, 2011. *Morb Mort Wkly Rep*. 2011 Feb 11. Erratum in: *Morb Mort Wkly Rep*. 2011 Mar 18;60(10):315.

Kidney Care

American Diabetes Association. Position statement executive summary: guidelines and recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus. *Diabetes Care*. 2011 Jun;34(6):1419-23.

Chronic Kidney Disease Prognosis Consortium. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet*. 2010 Jun 12;375(9731):2073-81.

Di Angelantonio E, Chowdhury R, Sarwar N, Aspelund T, Danesh J, Gudnason V. Chronic kidney disease and risk of major cardiovascular disease and non-vascular mortality: prospective population based cohort study. *BMJ*. 2010;341:c4986.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Holman RR, Paul SK, Bethel MA, Neil AW, Matthews DR. Long-term follow-up after tight control of blood pressure in type 2 diabetes. *N Eng J Med*. 2008;359(15):1565-76.

Kim NH, Pavkov ME, Knowler WC, Hanson RL, Weil EJ, Curtis JM, et al. Predictive value of albuminuria in American Indian youth with or without type 2 diabetes. *Pediatrics*. 2010;125:e844-51.

Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW, et al.; National Kidney Foundation. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med*. 2003 Jul 15;139(2):137-47. Erratum in: *Ann of Intern Med*. 2003 Oct 7;139(7):605.

McFarlane SI, Chen S, Whaley-Connell AT, Sowers JR, Vassalotti JA, Salifu MO, et al. Prevalence and associations of anemia of CKD: Kidney Early Evaluation Program (KEEP) and National Health and Nutrition Examination Survey (NHANES) 1999-2004. *Am J Kidney Dis*. 2008;51 4 Suppl 2:S46-55.

Miller WG. Estimating glomerular filtration rate. *Clin Chem Lab Med*. 2009;47:1017-9.

National Kidney Disease Education Program (US). Urine albumin-to-creatinine ratio (UACR) in evaluating patients with diabetes for kidney disease. Bethesda (MD): National Institutes of Health; 2010. 2 p. (NIH Publication No. 10-6286).

National Collaborating Centre for Chronic Conditions. Chronic kidney disease: national clinical guideline for early identification and management in adults in primary and secondary care. London: Royal College of Physicians; 2008 Sep. 204 p.

Narva, AS. Reducing the burden of chronic kidney disease among American Indians. *Adv Chronic Kidney Dis*. 2008;15:168-73.

Nelson RG, Pavkov ME, Hanson RL, Knowler WC. Changing course of diabetic nephropathy in the Pima Indians. *Diabetes Res Clin Pract*. 2008;82 Suppl 1: S10-4.

Pavkov ME, Knowler WC, Hanson RL, Bennett PH, Nelson RG. Predictive power of sequential measures of albuminuria for progression to ESRD or death in Pima Indians with type 2 diabetes. *Am J Kidney Dis*. 2008;51(5):759-66.

Pavkov ME, Knowler WC, Hanson RL, Nelson RG. Diabetic nephropathy in American Indians, with a special emphasis on the Pima Indians. *Curr Diab Rep*. 2008;8:486-93.

Pavkov ME, Mason CC, Bennett PH, Curtis JM, Knowler WC, Nelson RG. Change in the distribution of albuminuria according to estimated glomerular filtration rate in Pima Indians with type 2 diabetes. *Diabetes Care*. 2009;32(10):1845-50.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Schwartz GJ, Muñoz A, Schneider MF, Mak RH, Kaskel F, Warady BA, et al. New equations to estimate GFR in children with CKD. *J Am Soc Nephrol*. 2009;20(3):629-37.

Schwartz GJ, Work DF. Measurement and estimation of GFR in children and adolescents. *J Am Soc Nephrol*. 2009 Nov;4(11):1832-43.

Van Wyck DB, Eckardt K, Adamson JW, Bailie GR, Berns JS, Fishbane S, et al.; Anemia in Chronic Kidney Disease Work Group. National Kidney Foundation (NKF)-Kidney Disease Outcomes Quality Initiative (KDOQI) clinical practice guidelines and clinical practice recommendations for anemia in chronic kidney disease. New York: National Kidney Foundation; 2006. Available at http://www.kidney.org/professionals/KDOQI/guidelines_anemia

Vassalotti JA, Stevens LA, Levey AS. Testing for chronic kidney disease: a position statement from the National Kidney Foundation. *Am J Kidney Dis*. 2007;50(2):169-80.

Zelmanovitz T, Gerchman F, Balthazar APS, Thomazelli FCS, Matos JD, Canani LH. Diabetic nephropathy: review. *Diabetol Metab Syndr* [Internet]. 2009 Sep;doi:10.1186./1758-5996-1-10. Available at <http://www.dmsjournal.com/content/1/1/10>

Lipid Management

ACCORD Study Group. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med*. 2010 Apr 29;362:1563-74.

Brunzell JD, Davidson M, Furberg CD, Goldberg RB, Howard BV, Stein JH. Lipoprotein management in patients with cardiometabolic risk: consensus statement from the American Diabetes Association and the American College of Cardiology Foundation. *Diabetes Care*. 2008 Apr;31(4): 811-22.

Heart Protection Study Collaborative Group. MRC/BHF heart protection study of cholesterol lowering with simvastatin in 5963 people with diabetes: a randomized placebo-controlled trial. *Lancet*. 2003;361:2005-16.

Indian Health Service diabetes care and outcomes audit, unpublished data. Albuquerque (NM): Department of Health and Human Services (US), Indian Health Service, Office of Clinical and Preventive Services, Division of Diabetes Treatment and Prevention; 2011.

Indian Health Service, Division of Diabetes Treatment and Prevention. Guidelines for care of adults with prediabetes and/or the metabolic syndrome in the clinical setting. Albuquerque (NM): Department of Health and Human Services (US), Indian Health Service, Office of Clinical and Preventive Services, Division of Diabetes Treatment and Prevention; 2008 Sep. 31 p.

Indian Health *Standards of Care and Clinical Practice Recommendations:*
Type 2 Diabetes – Bibliography

Updated December 2011

Liu J, Sempos C, Donahue RP, Dorn J, Trevisan M, Grundy SM. Joint distribution of non-HDL and LDL cholesterol and coronary heart disease risk prediction among individuals with and without diabetes. *Diabetes Care*. 2005;28(8):1916-21.

Lu W, Resnick HE, Jablonski KA, Jones KL, Jain AK, Howard WJ, et al. Non-HDL cholesterol as a predictor of cardiovascular disease in type 2 diabetes: the Strong Heart Study. *Diabetes Care*. 2003;26(1):16-23.

National Cholesterol Education Program. Third report of the expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (ATP III final report). Bethesda (MD): National Heart, Lung, and Blood Institute (US); 2002 Sep. 280 p. (NIH Publication No. 02-5215).

Wing RR, Lang W, Wadden TA, Safford M, Knowler WC, Bertoni AG, et al.; Look AHEAD Research Group. Benefits of weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care*. 2011;34(7):1481-6.

Medical Nutrition Therapy

Bantle JP, Wylie-Rosett J, Albright AL, Apovian CM, Clark NG, Franz MJ, et al. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. *Diabetes Care*. 2008;31 Suppl 1:S61-78.

Coleman-Jensen A, Nord M, Andrews M, Carlson S. Household food security in the United States in 2010: Economic Research Report 125, U.S. Department of Agriculture, Economic Research Service; 2011 Sep. 29 p.

Coppell KJ, Kataoka M, Williams SM, Chisholm AW, Vorgers SM, Mann JI. Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment—Lifestyle Over and Above Drugs in Diabetes (LOADD) study: randomised controlled trial. *BMJ*. 2010;341;c3337.

Franz MJ, Powers A, Leontos C, Holzmeister LA, Kulkarni K, Monk A, et al. The evidence for medical nutrition therapy for type 1 and type 2 diabetes in adults. *J Am Diet Assoc*. 2010 Dec;110(12):1852-89.

O'Connell M, Buchwald DS, Duncan GE. Food access and cost in American Indian communities in Washington State. *J Am Diet Assoc*. 2011 Sept;111(9):1375-9.

Paxton AE, Strycker LA, Toobert DJ, Ammerman A, Glasgow RE. Starting the conversation: performance of a brief dietary assessment and intervention tool for health professionals. *Am J Prev Med*. 2011 Jan;40(1):67-71.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Salas-Salvadó J, Bulló M, Babio N, Martínez-González MA, Ibarrola-Jurado N, Basora J, et al. Reduction in the incidence of type 2 diabetes with the Mediterranean diet: results of the PREDIMED-Reus Nutrition Intervention Randomized Trial. *Diabetes Care*. 2011 Jan;34(1):14-9.

Seligman JL, Schillinger D. Hunger and socioeconomic disparities in chronic disease. *N Engl J Med*. 2010 Jul;363:6-9.

Stang J. Improving health among American Indians through environmentally-focused nutrition interventions. *J Am Diet Assoc*. 2009 Sep;109(9):1528-31.

U.S. Department of Agriculture; U.S. Department of Health and Human Services. *Dietary guidelines for Americans, 2010*. 7th ed. Washington (DC): U.S. Government Printing Office; December 2010. 95 p.

Van Horn L, McCoin M, Kris-Etherton PM, Burke F, Carson JA, Champagne CM, et al. The evidence for dietary prevention and treatment of cardiovascular disease. *J Am Diet Assoc*. 2008 Feb;108(2):287-331.

Wilson C, Brown T, Acton K, Gilliland S. Effects of clinical nutrition education and educator discipline on glycemic control outcomes in the Indian Health Service. *Diabetes Care*. 2003;26(9):2500-4.

Neuropathy

Argoff CE, Misha-Miroslav B, Belgrade MJ, Bennett GJ, Clark MR, Cole E, et al. Consensus guidelines: assessment, diagnosis, and treatment of diabetic peripheral neuropathic pain. *Mayo Clin Proc*. 2006 Apr;81 Suppl 4 :S12-25.

Boulton AJ, Vinik AI, Arezzo JC, Bril V, Feldman EL, Freeman R, et al. Diabetic neuropathies: a statement by the American Diabetes Association. *Diabetes Care*. 2005 Apr;28(4):956-62.

Bril V, England J, Franklin GM, Backonja M, Cohen J, Toro DD. Evidence-based guideline: treatment of painful diabetic neuropathy. Report of the American Academy of Neurology, the American Association of Neuromuscular and Electrodiagnostic Medicine, and the American Academy of Physical Medicine and Rehabilitation. *Neurology*. 2011 May 17;76(20):1758-65.

Freeman R. Not all neuropathy in diabetes is of diabetic etiology: differential diagnosis of diabetic neuropathy. *Curr Diab Rep*. 2009;9:423-31.

Vinik AI, Casellini C, Nakave A, Patel C. Chapter 31, Diabetic neuropathies. In: Goldfine ID, Rushakoff RJ, editors. *Diabetes and carbohydrate metabolism*, [Internet] *Diabetes Manager*. Endotext.org; 2009. Available at <http://www.endotext.org/diabetes/diabetes31/diabetesframe31.htm>

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Wile DJ, Toth C. Association of metformin, elevated homocysteine, and methylmalonic acid levels and clinically worsened diabetic peripheral neuropathy. *Diabetes Care*. 2010;33:156-61.

Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis

Medina J, Fernández-Salazar LI, García-Buey L, Moreno-Otero R. Approach to the pathogenesis and treatment of nonalcoholic steatohepatitis. *Diabetes Care*. 2004;27(8):2057-66.

Tolman KG, Fonseca V, Dalpiaz A, Tan MH. Spectrum of liver disease in type 2 diabetes and management of patients with diabetes and liver disease: review. *Diabetes Care*. 2007 Mar;30(3):734-43.

Vuppalanchi R, Chalasani N. Non-alcoholic fatty liver disease and non-alcoholic steatohepatitis: selected practical issues in their evaluation and management. *Hepatology*. 2009 Jan;49(1):306-17.

Obstructive Sleep Apnea

Chung F, Yegneswaran B, Liao P, Chung SA, Vairavanathan S, Islam S, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology*. 2008 May;108(5):822-30.

International Diabetes Federation. The IDF consensus statement on sleep apnoea and type 2 diabetes. Brussels: International Diabetes Federation; 2008. 24 p. Available at http://www.idf.org/webdata/docs/APNOEA_final.pdf

Shaw JE, Punjabi NM, Wilding JP, Alberti KG, Zimmet PZ. Sleep-disordered breathing and type 2 diabetes: a report from the International Diabetes Federation Taskforce on Epidemiology and Prevention. *Diabetes Res Clin Pract*. 2008;81:2-12.

Older Adults

Bethel MA, Sloan FA, Belsky D, Feinglos MN. Longitudinal incidence and prevalence of adverse outcomes of diabetes mellitus in elderly patients. *Arch Intern Med*. 2007 May 14;167(9):921-7.

Brown AF, Mangione CM, Saliba DM, Sarjustan CA; California Healthcare Foundation/American Geriatrics Society Panel on Improving Care for Elders with Diabetes. Guidelines for improving the care of the older person with diabetes mellitus. *J Am Geriatr Soc*. 2003 May;51 Suppl 5:S265-80.

Sloan FA, Bethel MA, Ruiz D Jr, Shea AM, Feinglos MN. The growing burden of diabetes mellitus in the US elderly population. *Arch Intern Med*. 2008 Jan 28;168(2):192-9.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Oral Care

Lamster IB, Lalla E, Borgnakke WS, Taylor GW. The relationship between oral health and diabetes mellitus. *J Am Dent Assoc.* 2008 Oct 1;139 Suppl 5:S19-24.

Mealey BL, Oates TW. Diabetes mellitus and periodontal diseases. *J Periodontol.* 2006;77:1289-1303.

Taylor GW, Borgnakke WS. Periodontal disease: associations with diabetes, glycemic control and complications. *Oral Dis.* 2008;14(3):204-5.

Teeuw WJ, Gerdes VE, Loos BG. Effect of periodontal treatment on glycemic control of diabetic patients: a systematic review and meta-analysis. *Diabetes Care.* 2010;33(2):421-7.

Peripheral Arterial Disease

Lamar Welch VL, Casper M, Greenlund K, Zheng ZJ, Giles W, Rith-Najarian S. Prevalence of lower extremity arterial disease defined by the ankle-brachial index among American Indians: the Inter-Tribal Heart Project. *Ethn Dis.* 2002;12(1):S1-63-7.

Mehler PS, Coll JR, Estacio R, Esler A, Schrier RW, Hiatt WR. Intensive blood pressure control reduces the risk of cardiovascular events in patients with peripheral arterial disease and type 2 diabetes. *Circulation.* 2003;107(5):753-6.

Resnick HE, Lindsay RS, McDermott MM, Devereux RB, Jones KL, Fabsitz RR, et al. Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: the Strong Heart Study. *Circulation.* 2004;109(6):733-9.

Rooke TW, Hirsch AT, Misra S, Sidawy AN, Beckman JA, Findeiss LK, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline); a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2011;58:2020-45. Epub 2011 Sep 29.

Physical Activity

American College of Sports Medicine. *ACSM's guidelines for exercise testing and prescription*, 8th ed. Baltimore: Lippincott Williams & Wilkins; 2009 Apr. 400 p.

Colberg SR, Sigal RJ, Fernhall B, Regensteiner JG, Blissmer BJ, Rubin RR, et al. Exercise and type 2 diabetes. The American College of Sports Medicine and the American Diabetes Association: joint position statement. *Diabetes Care.* 2010 Dec;33(12):2692-6.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Donnelly J, Blair SN, Jackicic J, Manore MM, Rankin J, Smith BK, et al. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc.* 2009 Feb;41(2):459-71.

Hansen D, Dendale P, Jonkers A, Beelan M, Manders RJF, Corluy L, et al. Continuous low- to moderate-intensity exercise training is as effective as moderate- to high-intensity exercise training at lowering blood HbA(1c) in obese type 2 diabetes patients. *Diabetologia.* 2009 Sep;52(9):1789-97.

Topolski TD, LoGerfo J, Patrick DL, Williams B, Walwick J, Patrick MB. The Rapid Assessment of Physical Activity (RAPA) among older adults. *Prev Chronic Dis.* [Internet]. 2006 Oct; [cited 2011 Nov 8]. Epub 2006 Sep 15. Available at http://www.cdc.gov/pcd/issues/2006/oct/06_0001.htm.

Preconception, Pregnancy, and Postpartum

American College of Obstetricians and Gynecologists Committee on Obstetric Practice. Committee opinion number 504: screening and diagnosis of gestational diabetes mellitus. *Obstet Gynecol.* 2011 Sep;118(3):751-3.

American Diabetes Association. Diagnosis and classification of diabetes mellitus: position statement. *Diabetes Care.* 2011 Jan;34 Suppl 1:S62-9. (See p. S68 for GDM)

Correa A, Gilboa SM, Besser LM, Botto LD, Moore CA, Hobbs CA, et al. Diabetes mellitus and birth defects. *Am J Obstet Gynecol.* 2008;199(3):237.e1-9.

Metzger BE, Gabbe SG, Persson B, Buchanan TA, Catalano PA, Damm P, et al.; International Association of Diabetes and Pregnancy Study Groups Consensus Panel. International Association of Diabetes and Pregnancy Study Groups recommendations in the diagnosis and classification of hyperglycemia in pregnancy. *Diabetes Care.* 2010 Mar;33(3):676-82.

Kitzmiller JL, Block JM, Brown FM, Catalano PM, Conway DL, Coustan, DR, et al. Managing preexisting diabetes for pregnancy: summary of evidence and consensus recommendations for care. *Diabetes Care.* 2008;31:1060-79.

Murphy N. ANMC OB/GYN Service diabetes mellitus in pregnancy screening and management guidelines. Anchorage: Alaska Native Medical Center; 2011. [cited 2011 Nov 8]. Available at <http://www.ihs.gov/MedicalPrograms/MCH/F/documents/DMpregn4611.pdf>

Wahabi HA, Alzeidan RA, Bawazeer GA, Alansari LA, Esmail SA. Preconception care for diabetic women for improving maternal and fetal outcomes: a systematic review and meta-analysis. *BMC Pregnancy Childbirth.* 2010 Oct 14;10:63. Available at <http://www.biomedcentral.com/content/pdf/1471-2393-10-63.pdf>

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Sexual Health

American College of Obstetricians and Gynecologists (ACOG). Female sexual dysfunction. Washington (DC): American College of Obstetricians and Gynecologists (ACOG). ACOG Practice Bulletin No. 119; 2011 April. 12 p.

Dhindsa F, Miller MG, McWhirter CL, Mager DE, Ghanim H, Chaudhuri A, et al. Testosterone concentrations in diabetic and nondiabetic obese men. *Diabetes Care*. 2010;33(6), 1186-92.

Lindau ST, Schumm LP, Laumann EO, Levinson W, O’Muirheartaigh CA, Waite LJ. A study of sexuality and health among older adults in the United States. *N Engl J Med*. 2007 Aug 23;357(8): 762-74.

Wang C, Jackson G, Jones TH, Matsumoto AM, Nehra A, Perelman MA, et al. Low testosterone associated with obesity and the metabolic syndrome contributes to sexual dysfunction and cardiovascular disease risk in men with type 2 diabetes. *Diabetes Care*. 2011 Jul;34:1669-73.

Tobacco Use

Caraballo RS, Yee SL, Pechacek TF, Henson R, Gfroerer JC. Tobacco use among racial and ethnic population subgroups of adolescents in the United States. *Prev Chron Dis*. 2006 Apr;3(2):A39. Epub 2006 Mar 15.

Centers for Disease Control and Prevention. Cigarette smoking among adults and trends in smoking—United States, 2008. *Morb Mortal Wkly Rep*. 2009 Nov 13;58(44);1227-32.

Fiore MC, Jaén CR, Baker TB, Bailey WC, Benowitz NL, Curry SJ, et al. Treating tobacco use and dependence: 2008 update: clinical practice guideline. Rockville (MD): Department of Health and Human Services (US), Public Health Service; 2008 May.

Ranney L, Melvin C, Lux L, McClain E, Lohr KN. Systematic review: smoking cessation intervention strategies for adults and adults in special populations. *Ann Intern Med*. 2006;145:845-56.

Sidransky D , Norman LA , McCarthy A , Taylor PL, editors. How tobacco smoke causes disease: the biology and behavioral basis for smoking-attributable disease: a report of the Surgeon General. Atlanta (GA): Department of Health and Human Services (US), Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010. 704 p.

Yeh HC, Duncan BB, Schmidt MI, Wang NY, Brancati FL. Smoking, smoking cessation, and risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med*. 2010 Jan 5;152(1):10-7.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Tuberculosis Screening

LoBue P, Sizemore C, Castro KG. Plan to combat extensively drug-resistant tuberculosis: recommendations of the Federal Tuberculosis Task Force. *Morb Mortal Wkly Rep.* 2009 Feb 13;58(RR03):1-43. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5803a1.htm>.

Mazurek GH, Jereb J, Vernon A, LeBue P, Goldberg S, Castro K. Updated guidelines for using interferon gamma release assays to detect mycobacterium tuberculosis infection—United States, 2010. *Morb Mortal Wkly Rep.* 2010 Jun 25;59(RR05):1-25. Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5905a1.htm?s_cid=rr5905a1_e.

Weight Management, Overweight, and Obesity

Franz MJ, VanWormer JJ, Crain AL, Boucher JL, Histon T, Caplan W, et al. Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc.* 2007;107:1755-67.

Klein S, Sheard NF, Pi-Sunyer X, Daly A, Wylie-Rosett J, Kulkarni K, et al. Weight management through lifestyle modification for the prevention and management of type 2 diabetes: rationale and strategies: a statement of the American Diabetes Association, the North American Association for the Study of Obesity, and the American Society for Clinical Nutrition. *Diabetes Care.* 2004;27(8):2067-73.

NHLBI Obesity Education Initiative Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: the evidence report. Bethesda (MD): Department of Health and Human Services (US), Public Health Service, National Institutes of Health; 1998 Sep. (NIH Publication No. 98-4083).

Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009 Feb 26;360:859-73.

Seagle HM, Strain GW, Makris A, Reeves RS; American Dietetic Association. Position of the American Dietetic Association: weight management. *J Am Diet Assoc.* 2009 Feb;109(2):330-46.

Wilson C, Gilliland S, Moore K, Acton K. The epidemic of extreme obesity among American Indian and Alaska Native adults with diabetes. *Prev Chronic Dis.* 2007 Jan;4(1):A06. Epub 2006 Dec 15.

Wing RR, Bahnson JL, Bray GA, Clark JM, Coday M, Egan C, et al.; Look AHEAD Research Group. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med.* 2010;170(17):1566-75.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011

Wing RR, Lang W, Wadden TA, Safford M, Knowler WC, Bertoni AG, et al.; Look AHEAD Research Group. Benefits of weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care*. 2011;34(7):1481-6.

Youth and Type 2 Diabetes

Acton KJ, Burrows NR, Moore K, Querec L, Geiss LS, Engelgau MM. Trends in diabetes prevalence among American Indian and Alaska Native children, adolescents, and young adults. *Am J Public Health*. 2002 Sep;92(9):1485-90.

Dabelea D, DeGroat J, Sorrelman C, Glass M, Percy CA, Avery C, et al. Diabetes in Navajo youth—prevalence, incidence and clinical characteristics: the SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2009;32(2) Suppl:S141-7.

Dabelea D, Mayer-Davis EJ, Lamichhane AP. Association of intrauterine exposure to maternal diabetes and obesity with type 2 diabetes in youth: the SEARCH Case-Control Study. *Diabetes Care*. 2008;31(7):1422-6.

Dabelea D, Pihoker C, Talton JW, D'Agostino RB Jr, Fujimoto W, Klingensmith GJ, et al.; SEARCH for Diabetes in Youth Study. Etiological approach to characterization of diabetes type: the SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2011 Jul;34(7):1628-33.

Flint A, Arslanian S. Treatment of type 2 diabetes in youth. *Diabetes Care*. 2011 May;34 Suppl 2: S177-83.

Indian Health Service diabetes care and outcomes audit, unpublished data. Albuquerque (NM): Department of Health and Human Services (US), Indian Health Service, Office of Clinical and Preventive Services, Division of Diabetes Treatment and Prevention; 1990-2009.

Kreba NF, Himes JH, Jacobson D, Nicklas T, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. *Pediatrics*. 2007 Dec 1;120 Suppl 4:S193-228.

Moore K. Youth-onset type 2 diabetes among American Indians and Alaska Natives. *J Public Health Manag Pract*. 2010;16(5):388-93.

National High Blood Pressure Education Program. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Bethesda (MD): National Heart, Lung, and Blood Institute (US); 2005 May. (NIH Publication No. 05-5267).

Peterson K, Silverstein J, Kaufman F, Warren-Boulton E. Management of type 2 diabetes in youth: an update. *Am Fam Physician*. 2007 Sep 1;76(5):658-64.

Rosenbloom AL, Silverstein JH, Amemiya S, Zeitler P, Klingensmith G. Type 2 diabetes in children and adolescents. *Pediatric Diabetes*. 2009;10 Suppl 12: 17-32.

*Indian Health Standards of Care and Clinical Practice Recommendations:
Type 2 Diabetes – Bibliography*

Updated December 2011