

Toolkit 4: CRS Breastfeeding Guidelines/Logic

1.1.1 Breastfeeding Rates

No changes from Version 11.1

Note: This measure is used in conjunction with the Childhood Weight Control GPRA measure to support the reduction of the incidence of childhood obesity.

Owner/Contact

Tina Tah, RN, BSN, MBA

National Reporting

OTHER NATIONAL (included in new Other National Measures Report; not reported to OMB and Congress)

Denominators

1. Active Clinical patients who are 30–394 days old.
2. **PART:** Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of two months (45–89 days).
3. Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of six months (165–209 days).
4. Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of nine months (255–299 days).
5. Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of 1 year (350–394 days)

Numerators

1. Patients who were screened for infant feeding choice at least once.
2. Patients who were screened for infant feeding choice at the age of two months (45–89 days).
3. Patients were screened for infant feeding choice at the age of six months (165–209 days).
4. Patients who were screened for infant feeding choice at the age of nine months (255–299 days).
5. Patients who were screened for infant feeding choice at the age of 1 year (350–394 days).
6. **PART:** Patients who, at the age of two months (45–89 days), were either exclusively or mostly breastfed.
7. Patients who, at the age of six months (165–209 days), were either exclusively or mostly breastfed.
8. Patients who, at the age of nine months (255–299 days), were either exclusively or mostly breastfed.
9. Patients who, at the age of 1 year (350–394 days), were either exclusively or mostly breastfed.

Definitions

Infant Feeding Choice

The documented feeding choice from the file V Infant Feeding Choice that is closest to the exact age that is being assessed will be used. For example, if a patient was assessed at 45 days old as 1/2 breastfed and 1/2 formula and assessed again at 65 days old as mostly breastfed, the mostly breastfed value will be used since it is closer to the exact age of 2 months (i.e. 60 days). Another example is a patient who was assessed at 67 days as mostly breastfed and again at 80 days as mostly formula. In this case, the 67 days value of mostly breastfed will be used. The other exact ages are 180 days for 6 months, 270 days for 9 months, and 365 days for 1 year.

In order to be included in the age-specific screening numerators, the patient must have been screened at the specific age range. For example, if a patient was screened at 6 months and was exclusively breastfeeding but was not screened at 2 months, then the patient will only be counted in the 6 months numerator.

2011 Performance Description

During FY 2011, achieve the target rate of 28.6% for the proportion of 2-month olds who are mostly or exclusively breastfeeding.

Patient List

List of patients 30–394 days old, with infant feeding choice value, if any.