

Measure	FY	Target	Result
<p>1: Diabetes: Poor Glycemic Control: Proportion of patients with diagnosed diabetes that have poor glycemic control (A1c > 9.5). IHS-All (Outcome)</p> <p><i>NOTE: This measure is directed at decreasing the percentage of diabetic patients with poor glycemic control; therefore, a lower number is better.</i></p> <p>Prior to 2004, results were reported from the Diabetes (DM) Audit.</p>	2012	18.6%	N/A
	2011	19.4%	19.1% (Target Exceeded)
	2010	16%	18% (Target Not Met)
	2009	18%	18% (Target Met)
	2008	16%	17% (Target Not Met)
	2007	15%	16% (Target Not Met)
	2006	15%	16% (Target Not Met)
	2005	17%	15% (Target Exceeded)
	2004	Set Baseline	17% (Baseline)

Measure	FY	Target	Result
<p>2: Diabetes: Ideal Glycemic Control: Proportion of patients with diagnosed diabetes with ideal glycemic control (A1c < 7.0). IHS-All (Outcome)</p> <p>Prior to 2002, results were reported from the DM Audit.</p>	2012	32.7%	N/A
	2011	30.2%	31.9% (Target Exceeded)
	2010	33%	32% (Target Not Met but Improved)
	2009	30%	31% (Target Exceeded)
	2008	31%	32% (Target Exceeded)
	2007	32%	31% (Target Not Met)
	2006	32%	31% (Target Not Met but Improved)
	2005	27%	30% (Target Exceeded)
	2004	29%	27% (Target Not Met)
	2003	25%	28% (Target Exceeded)
	2002		25% (result reported from CRS)

Measure	FY	Target	Result
3: Diabetes: Blood Pressure Control: Proportion of patients with diagnosed diabetes that have achieved blood pressure control (<130/80). IHS-All (<i>Outcome</i>)	2012	38.7%	N/A
	2011	35.9%	37.8% (Target Exceeded)
	2010	40%	38% (Target Not Met but Improved)
	2009	36%	37% (Target Exceeded)
	2008	39%	38% (Target Not Met)
	2007	37%	39% (Target Exceeded)
	2006	37%	37% (Target Met)
	2005	35%	37% (Target Exceeded)
	2004	38%	35% (Target Not Met)
	2003	36%	37% (Target Exceeded)
	2002		36% (result reported from CRS)

Year	Percent
2002	36
2003	37
2004	35
2005	37
2006	37
2007	39
2008	38
2009	37
2010	38
2011	37.8
2012	38.7

Prior to 2002, results were reported from the DM Audit.

Measure	FY	Target	Result
4: Diabetes: Dyslipidemia Assessment: Proportion of patients with diagnosed diabetes assessed for dyslipidemia (LDL cholesterol). IHS-All (<i>Outcome</i>)	2012	70.3%	N/A
	2011	63.3%	68.7% (Target Exceeded)
	2010	69%	67% (Target Not Met but Improved)
	2009	60%	65% (Target Exceeded)
	2008	61%	63% (Target Exceeded)
	2007	60%	61% (Target Exceeded)
	2006	53%	60% (Target Exceeded)
	2005	53%	53% (Target Met)
	2004	49%	53% (Target Exceeded)
	2003	44%	48% (Target Exceeded)
	2002		44% (result reported from CRS)

Year	Percent
2002	44
2003	48
2004	53
2005	53
2006	60
2007	61
2008	63
2009	65
2010	67
2011	68.7
2012	70.3

Prior to 2002, results were reported from the DM Audit.

Measure	FY	Target	Result
5: Diabetes: Nephropathy Assessment: Proportion of patients with diagnosed diabetes assessed for nephropathy. IHS-All (Outcome)	2012	57.8%	N/A
	2011	51.9%	56.5% (Target Exceeded)
	2010	54%	55% (Target Exceeded)
	2009	47%	50% (Target Exceeded)
	2008	40%	50% (Target Exceeded)
	2007	Set Baseline (Based on new standards of care)	40% (Baseline)


Year	Percent
2007	40
2008	50
2009	50
2010	55
2011	56.5
2012	57.8

Measure	FY	Target	Result
6: Diabetic Retinopathy: Proportion of patients with diagnosed diabetes who receive an annual retinal examination. IHS - All (Outcome)	2012	54.8%	N/A
	2011	50.1%	53.5% (Target Exceeded)
	2010	55%	53% (Target Not Met but Improved)
	2009	47%	51% (Target Exceeded)
	2008	49%	50% (Target Exceeded)
	2007	49%	49% (Target Met)
	2006	Set Baseline	49% (Baseline)


Year	Percent
2006	49
2007	49
2008	50
2009	51
2010	53
2011	53.5
2012	54.8

Measure	FY	Target	Result
12: Topical Fluorides: Number of American Indian and Alaska Native patients receiving at least one topical fluoride application. (Outcome)	2012	161,461 patients	N/A
	2011	135,604 patients	161,461 patients (Target Exceeded)
	2010	136,978 patients	145,181 patients (Target Exceeded)
	2009	114,716 patients	136,794 patients (Target Exceeded)
	2008	107,934 patients	120,754 patients (Target Exceeded)
	2007	95,439 patients	107,934 patients (Target Exceeded)
	2006	85,318 patients	95,439 patients (Target Exceeded)
2005	Set Baseline	85,318 patients (Baseline)	

Year	Number of Patients
2005	85,318
2006	95,439
2007	107,934
2008	120,754
2009	136,794
2010	145,181
2011	161,461
2012	161,461

Measure	FY	Target	Result
13: Dental Access: Percent of patients who receive dental services. (Outcome) 	2012	26.9%	N/A
	2011	23.0%	26.9% (Target Exceeded)
	2010	27%	25% (Target Not Met)
	2009	24%	25% (Target Exceeded)
	2008	25%	25% (Target Met)
	2007	24%	25% (Target Exceeded)
	2006	24%	23% (Target Not Met)
	2005	24%	24% (Target Met)
	2004	25%	24% (Target Not Met)
	2003	27%	25% (Target Not Met)
	2002		27% (result reported from CRS)

Prior to 2002, results were reported from the IHS National Patient Information Reporting System (NPIRS).

Measure	FY	Target	Result
14: Dental Sealants: Number of sealants placed per year in AI/AN patients. (Outcome) 	2012	276,893 sealants	N/A
	2011	257,261 sealants	276,893 sealants (Target Exceeded)
	2010	257,920 sealants	275,459 sealants (Target Exceeded)
	2009	229,147 sealants	257,067 sealants (Target Exceeded)
	2008	245,449 sealants	241,207 sealants (Target Not Met)
	2007	246,645 sealants	245,449 sealants (Target Not Met)
	2006	249,882 sealants	246,645 sealants (Target Not Met)
	2005	230,295 sealants	249,882 sealants (Target Exceeded)
	2004	230,295 sealants	230,295 sealants

Prior to 2005, results were reported from NPIRS.

Measure	FY	Target	Result
25: Adult Immunizations: Influenza: Influenza vaccination rates among adult patients aged 65 years and older. IHS-All (<i>Outcome</i>)	2012	63.4%	N/A
	2011	58.5%	62.0% (Target Exceeded)
	2010	60%	62% (Target Exceeded)
	2009	62%	59% (Target Not Met)
	2008	59%	62% (Target Exceeded)
	2007	59%	59% (Target Met)
	2006	59%	58% (Target Not Met)
	2005	On hold because of Influenza vaccine Shortage in 2005 (54% original target)	On hold (59% result provided for context)
	2004	51%	54% (Target Exceeded)
	2003	51%	51% (Target Met)
	2002		51% (result reported from CRS)

Percent of Influenza 65+ by Year

Year	Percent
2002	51
2003	51
2004	54
2005	59
2006	58
2007	59
2008	62
2009	59
2010	62
2011	62.0
2012	63.4

Prior to 2002, results reported from NPIRS.

Measure	FY	Target	Result
26: Adult Immunizations: Pneumovax: Pneumococcal vaccination rates among adult patients aged 65 years and older. IHS-All (<i>Outcome</i>)	2012	87.5%	N/A
	2011	79.3%	85.5% (Target Exceeded)
	2010	83%	84% (Target Exceeded)
	2009	82%	82% (Target Met)
	2008	79%	82% (Target Exceeded)
	2007	76%	79% (Target Exceeded)
	2006	72%	74% (Target Exceeded)
	2005	69%	69% (Target Met)
	2004	65%	69% (Target Exceeded)
	2003	64%	65% (Target Exceeded)
	2002	Set Baseline	64% (Baseline)

Percent of Pneumovax 65+ by Year

Year	Percent
2002	64
2003	65
2004	69
2005	69
2006	74
2007	79
2008	82
2009	82
2010	84
2011	85.5
2012	87.5

Prior to 2002, results reported from NPIRS.

Measure	FY	Target	Result
24: Combined (4:3:1:3:3:1:4) Childhood immunization rates: AI/AN children patients aged 19-35 months. In 2010 CDC added the Varicella vaccine to the basic series [4:3:1:3:3:1] and in 2011, Pneumococcal conjugate was added. IHS - All (<i>Outcome</i>)	2012	77.8% [4:3:1:3:3:1:4]	N/A
	2011	74.6% [4:3:1:3:3:1:4]	75.9% (Target Exceeded)
	2010	80% [4:3:1:3:3:1]	79% (Target Not Met)
	2009	78% [4:3:1:3:3]	79% (Target Exceeded)
	2008	78% [4:3:1:3:3]	78% (Target Met)
	2007	78% [4:3:1:3:3]	78% (Target Met)
	2006	Set Baseline for reporting from Clinical Reporting System (CRS) [4:3:1:3:3 Series] 4DTap, 3IPV, 1MMR, 3Hib, 3HepB)	78% (Baseline)

Percent of Childhood Immunizations by Year

Year	Percent
2006	78
2007	78
2008	78
2009	79
2010	79
2011	75.9
2012	77.8

Measure	FY	Target	Result
7: Pap Smear Rates: Proportion of eligible women who have had a Pap screen within the previous three years. IHS - All (<i>Outcome</i>)	2012	59.5%	N/A
	2011	55.7%	58.1% (Target Exceeded)
	2010	60%	59% (Target Not Met)
	2009	59%	59% (Target Met)
	2008	59%	59% (Target Met)
	2007	60%	59% (Target Not Met)
	2006	60%	59% (Target Not Met)
	2005	58%	60% (Target Exceeded)
	2004	61%	58% (Target Not Met)
	2003	62%	61% (Target Not Met)
	2002		62% (result reported from CRS)


Percent Pap Smears by Year


Year	Percent
2002	62
2003	61
2004	58
2005	60
2006	59
2007	59
2008	59
2009	59
2010	59
2011	58.1
2012	59.5

Prior to 2002, results reported from NPIRS.


Measure	FY	Target	Result
8: Mammogram Rates: Proportion of eligible women who have had mammography screening within the previous two years. IHS - All (Outcome) <i>Prior to 2002, results reported from NPIRS.</i>	2012	51.7%	N/A
	2011	46.9%	49.8% (Target Exceeded)
	2010	47%	48% (Target Exceeded)
	2009	45%	45% (Target Met)
	2008	43%	45% (Target Exceeded)
	2007	41%	43% (Target Exceeded)
	2006	41%	41% (Target Met)
	2005	40%	41% (Target Exceeded)
	2004	40%	40% (Target Met)
	2003	42%	40% (Target Not Met)
	2002		42% (result reported from CRS)

Measure	FY	Target	Result
9: Colorectal Cancer Screening Rates: Proportion of eligible patients who have had appropriate colorectal cancer screening. IHS - All (Outcome) 	2012	43.2%	N/A
	2011	36.7%	41.7% (Target Exceeded)
	2010	36%	37% (Target Exceeded)
	2009	29%	33% (Target Exceeded)
	2008	26%	29% (Target Exceeded)
	2007	22%	26% (Target Exceeded)
	2006 (New measure)	Set Baseline	22% (Baseline)


Measure	FY	Target	Result
32: Tobacco Cessation Intervention: Proportion of tobacco-using patients that receive tobacco cessation intervention. IHS-All (<i>Outcome</i>) 	2012	30.0%	N/A
	2011	23.7%	29.4% (Target Exceeded)
	2010	27%	25% (Target Not Met but Improved)
	2009	21%	24% (Target Exceeded)
	2008	16%	21% (Target Exceeded)
	2007	12%	16% (Target Exceeded)
	2006 (New Measure)	Set Baseline	12% (Baseline)

Measure	FY	Target	Result
11: Alcohol Screening (FAS Prevention): Alcohol-use screening (to prevent Fetal Alcohol Syndrome) among appropriate female patients. IHS-All (<i>Outcome</i>) 	2012	58.7%	N/A
	2011	51.7%	57.8% (Target Exceeded)
	2010	55%	55% (Target Met)
	2009	47%	52% (Target Exceeded)
	2008	41%	47% (Target Exceeded)
	2007	28%	41% (Target Exceeded)
	2006	12%	28% (Target Exceeded)
	2005	7%	11% (Target Exceeded)
	2004 (New measure)	Set Baseline	7% (Baseline)

Measure	FY	Target	Result
16: Domestic (Intimate Partner) Violence Screening: Proportion of women who are screened for domestic violence at health care facilities. IHS-All (<i>Outcome</i>)	2012	55.3%	N/A
	2011	52.8%	55.3% (Target Exceeded)
	2010	53%	53% (Target Met)
	2009	42%	48% (Target Exceeded)
	2008	36%	42% (Target Exceeded)
	2007	28%	36% (Target Exceeded)
	2006	14%	28% (Target Exceeded)
	2005	4%	13% (Target Exceeded)
	2004	15%	4% (Target Not Met)
	2004 (New measure)		



Measure	FY	Target	Result
18: Behavioral Health: Proportion of adults ages 18 and over who are screened for depression. IHS-All (<i>Outcome</i>)	2012	56.5%	N/A
	2011	51.9%	56.5% (Target Exceeded)
	2010	53%	52% (Target Not Met but Improved)
	2009	35%	44% (Target Exceeded)
	2008	24%	35% (Target Exceeded)
	2007	15%	24% (Target Exceeded)
	2006	Set Baseline	15% (Baseline)
	2006 (New measure)		



Measure	FY	Target	Result
30: CVD Comprehensive Assessment: Proportion of Ischemic Heart Disease (IHD) patients who have a comprehensive assessment for all five CVD-related risk factors. <i>(Outcome)</i>	2012	40.6%	N/A
	2011	33.0%	39.8% (Target Exceeded)
	2010	33%	35% (Target Exceeded)
	2009	30%	32% (Target Exceeded)
	2008	30%	30% (Target Met)
	2007	Set Baseline based on logic changes to the measure	30% (Baseline)

Year	Percent
2007	30
2008	30
2009	32
2010	35
2011	39.8
2012	40.6

Measure	FY	Target	Result
33: HIV Screening: Proportion of pregnant women screened for HIV. <i>(Outcome)</i>	2012	81.8%	N/A
	2011	73.6%	80.0% (Target Exceeded)
	2010	77%	78% (Target Exceeded)
	2009	75%	76% (Target Exceeded)
	2008	74%	75% (Target Exceeded)
	2007	65%	74% (Target Exceeded)
	2006	55%	65% (Target Exceeded)
	2005 (New measure)	Set Baseline	54% (Baseline)

Year	Percent
2005	54
2006	65
2007	74
2008	75
2009	76
2010	78
2011	80.0
2012	81.8