

# ED Discharge Template Screen Shots

**Template: EMER DEPT DC INSTRUCTIONS**

---

Phoenix Indian Medical Center                      Patient: DEMO,PATIENT JOHN TEEN  
4212 N. 16th Street                                      DOB: JAN 1,2000  
Phoenix, AZ 85016                                        HR#: 99-99-76  
Visit: 02/27/17 09:26

---

PHOENIX INDIAN MEDICAL CENTER EMERGENCY DEPARTMENT

PATIENT DISCHARGE INSTRUCTIONS

---

GENERAL INSTRUCTIONS:

(1) PATIENT RESPONSIBILITY: It is YOUR RESPONSIBILITY to obtain follow up care as instructed below. It is your responsibility to call the clinic you are referred to for follow up care and make an appointment. When you call to make an appointment advise the clinic that your were referred by the Emergency Department.

(2) MEDICAL CONDITION WORSENS: If you become concerned about your condition and are unable to contact your physician, or if your condition is a medical emergency go to the Emergency Department.

(3) XRAYs: If X-Rays were taken today, the interpretation in the Emergency Department is PRELIMINARY. If the final radiologist interpretation reveals a different significant finding, you will be contacted.

---

During todays visit at the Phoenix Indian Medical Center Emergency Department, you were seen by the following medical medical provider.

PIMC ER MEDICAL PROVIDER:

DIAGNOSIS: Your diagnosis for this visit was the following:

No Diagnoses Found

CONDITION SPECIFIC CARE INSTRUCTIONS

PIMC Specific

MALE

Epididymitis

ABDOMINAL PAIN male

Urethritis Male

Figure 1: EMER Dept DC Instructions

## ED Discharge Template Screen Shots

Female

- Pregnancy
- MASTITIS
- STD, Suspected
- Bacterial Vaginosis
- Ovarian Cyst
- Irregular vaginal bleeding
- Threatened Miscarriage
- Abdominal pain female
- BLADDER INFECTION, Female

Muscle skeletal

- Shoulder Fracture (Shoulder Immobilizer)
- DISLOCATION: SHOULDER (Reduced)
- Nursemaid's Elbow
- SCIATICA
- Bursitis, Elbow (Olecranon)
- Muscle spasm
- SPRAIN: KNEE
- FRACTURE: HAND (closed)
- Fracture: Toe (Closed)
- Fracture:Foot
- Fracture: Forearm (Radius/Ulna)
- SPRAIN: ELBOW
- Costochondritis
- Rib: Contusion Vs Minor Fracture
- CONTUSION: LOWER EXTREMITY
- CONTUSION: HAND
- CONTUSION, SOFT TISSUE

Figure 2: Female and Muscle skeletal

## ED Discharge Template Screen Shots

Neck Sprain or Strain

Carpal tunnel syndrome

Bursitis

BACK PAIN [acute or chronic]

SPRAIN: ANKLE

FRACTURE ANKLE

Muscle Strain in the Abdomen

Shoulder Blade or Collarbone Fracture

Contusion, soft tissue

Back pain

ANKLE SPRAIN

SPRAIN: A-C JOINT

Strep throat

Your illness has the signs of a strep throat infection. Strep throat is a contagious illness. It is spread by coughing, kissing or by touching others after touching your mouth or nose. Symptoms include throat pain worse with swallowing, aching all over, headache and fever. You will be treated with an antibiotic, which should make you start to feel better within 1-2 days.

HOME CARE:

1. Rest at home and drink plenty of fluids to avoid dehydration.
2. No school or work for the first two days on antibiotics. You will not be contagious after this time, and if you are feeling better, you can return to school or work.
3. Take your antibiotics for a full 10 days, even if you feel better after the first few days of treatment. This is very important to prevent complications from the strep infection (such as heart or kidney disease).
4. Children: Use acetaminophen (Tylenol) for fever, fussiness or discomfort. In infants over six months of age, you may use ibuprofen (Children's Motrin) instead of Tylenol. NOTE: If your child has chronic liver or kidney disease or ever had a stomach ulcer or GI bleeding, talk with your doctor before using these medicines. (Aspirin should never be used in anyone under 18 years of age who is ill with a fever. It may cause severe liver damage.)  
Adults: You may use acetaminophen (Tylenol) or ibuprofen (Motrin, Advil) to

Figure 3: Discharge Instructions – Strep Throat

## ED Discharge Template Screen Shots

Adults: You may use acetaminophen (Tylenol) or ibuprofen (Motrin, Advil) to control pain or fever, unless another medicine was prescribed for this. NOTE: If you have chronic liver or kidney disease or ever had a stomach ulcer or GI bleeding, talk with your doctor before using these medicines.

5. Throat lozenges or sprays (Chloraseptic and others) will reduce pain. Gargling with warm salt water will also reduce throat pain. Dissolve 1/2 teaspoon of salt in 1 glass of warm water. This is especially useful just before meals.

FOLLOW UP with your doctor or express care within the next 2 days.

GET PROMPT MEDICAL ATTENTION if any of the following occur:

- Fever over 100.5°F (38.0°C) oral, or over 101.5°F (38.6°C) rectal for more than three days
- New or worsening ear pain, sinus pain or headache
- Painful lumps in the back of your neck
- Unable to swallow liquids or open your mouth wide due to throat pain
- Trouble breathing or noisy breathing
- Muffled voice
- New rash

LARYNGITIS

BRONCHITIS

Asthma

CROUP, Viral (Child)

Pneumonia (Child)

PNEUMONIA (Adult)

VIRAL SYNDROME [Adult]

VIRAL SYNDROME [Child]

INFLUENZA (Child)

INFLUENZA (Adult)

Otitis Externa

EUSTACHIAN TUBE OBSTRUCTION

ACUTE OTITIS MEDIA WITH INFECTION [Child]

DOG BITE

Diarrhea

Figure 4: Discharge Instructions – Strep Throat continued

## ED Discharge Template Screen Shots

Diarrhea

Atopic dermatitis

DERMATITIS (Non-specific)

Dehydration

Benign Positional Vertigo

COPD Flare

Croup

Corneal Abrasion

Bacterial Conjunctivitis

Chronic Pain

Heart Failure

Chest Pain, Uncertain Cause

Bursitis

Burn 2ed Degree

Bronchitis

High blood pressure

Bell's Palsy

Asthma

Nasal Allergy

Allergic Reaction

Abdominal pain male

ABSCESS

ALLERGIC REACTION, GENERALIZED

CHEST PAIN

KRAMES DISCHARGE INSTRUCTIONS:

KRAMES ON DEMAND: A printed copy of diagnosis specific discharge instructions generated using Krames on Demand Exit-Writer were provided to the patient/care giver on discharge from the Emergency Department.

[Click to open Krames Exit-Writer Discharge Instructions](#)

Figure 5: Krames Discharge Instructions

# ED Discharge Template Screen Shots

Importing Krames Exit-Writer discharge instructions into EHR

1. Click on the link to open the Krames Patient Education application
2. For Discharge Instructions click on Exit-Writer on the left side of the page
3. Select the desired instruction sheet
4. Enter the Patient's name and personalized discharge instructions
5. Click "GO" to print a copy for the instructions for the patient or if giving the patient several sheets save to the "Education Cart" to print as a group when your EHR note is complete.
6. Highlight and copy the content of the instruction sheet you want to include in EHR
7. Close this template
8. Paste the discharge information you want to document in EHR into your note.
9. Repeat steps 2-3 for each instruction sheet selected.

\_\_\_\_\_  
The following is the text of the Instructional Material provided to the patient from Krames Exit Writer.  
The patient copy may also include graphics and diagrams that cannot be saved in this note

\_\_\_\_\_

PASTE KRAMES CONTENT HERE AFTER THE TEMPLATEW IS CLOSED

\_\_\_\_\_

FOLLOW UP CARE

Please schedule a follow up appointment with your Provider / Clinic as noted

MEDICATION/PRESCRIPTIONS

MODE OF DEPARTURE:

Patient left the Emergency Department by

Patient left the Emergency Department with

ADDITIONAL INSTRUCTIONS:

WORK RELEASE

-----

To Whom It May Concern:


DEMO, PATIENT JOHN TEEN was seen at the Phoenix Indian Medical Center Emergency Department on MAR 03, 2017. PATIENT was seen by a emergency room medical provider who reccommends the following:

o Patient may return to work/school with the following limitations/modifications:

Figure 6: Import Krames Exit-Writer discharge instructions into EHR

## ED Discharge Template Screen Shots

---



o As PIMC employee, we recommend the patient be off work 1 days. This requires your supervisor approval.

o The medical provider recommends patient be off work/school for days.

YOU MUST TALK TO YOUR EMPLOYER OR SCHOOL  
BEFORE YOU TAKE TIME OFF FROM WORK OR SCHOOL.

o Patient may return to work/school today without any limitations.

Figure 7: Work Release