



RESOURCE AND PATIENT MANAGEMENT SYSTEM

Clinical Reporting System (BGP)

CRS Clinical Performance Measure Logic Manual for FY 2011 Clinical Measures

Version 11.1
June 2011

Office of Information Technology (OIT)
Division of Information Resource Management
Albuquerque, New Mexico

Preface

The Government Performance and Results Act (GPRA) requires federal agencies to report annually on how the agency measured up against the performance targets set in its annual Plan. The Indian Health Service (IHS) GPRA report includes measures for clinical prevention and treatment, quality of care, infrastructure, and administrative efficiency functions.

The IHS Clinical Reporting System (CRS) is a Resource and Patient Management System (RPMS) software application designed for national reporting as well as Area Office and local monitoring of clinical GPRA and developmental measures. CRS was first released for Fiscal Year (FY) 2002 performance measures (as GPRA+) and is based on a design by the Aberdeen Area (GPRA2000).

This manual contains the FY 2011 clinical performance measure definitions and logic for the CRS 2011 Version 11.1 software. CRS is the reporting tool used by the IHS Office of Planning and Evaluation to collect and report clinical performance results annually to the Department of Health and Human Services and to Congress.

Each year, an updated version of CRS software is released to reflect changes in the logic descriptions of the different denominators and numerators. Additional performance measures may also be added. Local facilities can run reports as often as they want to and can also use CRS to transmit data to their Area. The Area Office can use CRS to produce an aggregated Area report for either annual GPRA or Area Director Performance reports.

CRS produces reports on demand from local RPMS databases for both GPRA and developmental clinical measures that are based on RPMS data, thus eliminating the need for manual chart audits for evaluating and reporting clinical measures.

To produce reports with comparable data across every facility, the GPRA measures definitions was “translated” into programming code with the assistance of clinical subject matter experts. CRS uses predefined taxonomies to find data items in the RPMS Patient Care Component (PCC) to determine if a patient meets the performance measure criteria. Taxonomies contain groups of codes (e.g., diagnoses or procedures) or site-specific terms. Each performance measure has one or more defined denominators and numerators.

Administrative and clinical users can produce reports for selected measures at any time to:

- Identify potential data issues in their RPMS, i.e., missing or incorrect data
- Monitor their site’s performance against past national performance and upcoming agency goals

- Identify specific areas where the facility is not meeting the measure in order to initiate business process or other changes
- Quickly measure impact of process changes on performance measures
- Identify areas meeting or exceeding measures to provide lessons learned

Users of the RPMS CRS include:

- Area Office and site quality improvement staff
- Compliance Officers
- GPRA coordinators
- Clinical staff, such as physicians, nurses, nurse practitioners, and other providers
- Area Office directors
- Any staff involved with quality assurance initiatives
- Staff who run various CRS reports

Table of Contents

1.0 About This Manual.....	1
2.0 Performance Measure Logic.....	1
2.1 Performance Measure Logic Basics	1
2.1.1 CRS Denominator Definitions.....	1
2.1.2 Performance Measure Logic Example.....	7
2.1.3 Age Ranges	9
2.1.4 Standard Health Care Codes.....	9
2.2 Diabetes Related Measure Topics	10
2.2.1 Diabetes Prevalence	10
2.2.2 Diabetes Comprehensive Care.....	13
2.2.3 Diabetes: Glycemic Control	17
2.2.4 Diabetes: Blood Pressure Control	22
2.2.5 Diabetes: LDL Assessment	25
2.2.6 Diabetes: Nephropathy Assessment.....	29
2.2.7 Diabetic Retinopathy	32
2.2.8 ACEI/ARB Use in Diabetic Patients.....	36
2.2.9 Diabetes: Access to Dental Services.....	40
2.3 Dental Measure Topics	42
2.3.1 Access to Dental Services.....	42
2.3.2 Dental Sealants.....	45
2.3.3 Topical Fluoride.....	47
2.4 Immunization Measure Topics	50
2.4.1 Influenza	50
2.4.2 Adult Immunizations	54
2.4.3 Childhood Immunizations	59
2.4.4 Adolescent Immunizations.....	70
2.5 Childhood Diseases Group	77
2.5.1 Appropriate Treatment for Children with Upper Respiratory Infection	77
2.5.2 Appropriate Testing for Children with Pharyngitis	80
2.6 Cancer Related Measure Topics.....	82
2.6.1 Cancer Screening: Pap Smear Rates.....	82
2.6.2 Cancer Screening: Mammogram Rates.....	86
2.6.3 Colorectal Cancer Screening.....	89
2.6.4 Comprehensive Cancer Screening.....	93
2.6.5 Tobacco Use and Exposure Assessment	98
2.6.6 Tobacco Cessation.....	105
2.7 Behavioral Health Related Performance Measure Topics.....	110
2.7.1 Alcohol Screening (FAS Prevention)	110
2.7.2 Alcohol Screening and Brief Intervention (ASBI) in the ER.....	114
2.7.3 Intimate Partner (Domestic) Violence Screening	118
2.7.4 Depression Screening	121
2.7.5 Antidepressant Medication Management.....	126
2.8 Cardiovascular Disease Related Measure Topics	131

2.8.1	Obesity Assessment.....	131
2.8.2	Childhood Weight Control.....	136
2.8.3	Nutrition and Exercise Education for At Risk Patients.....	139
2.8.4	Physical Activity Assessment	143
2.8.5	Comprehensive Health Screening	146
2.8.6	Cardiovascular Disease and Cholesterol Screening	152
2.8.7	Cardiovascular Disease and Blood Pressure Control	155
2.8.8	Controlling High Blood Pressure.....	158
2.8.9	Comprehensive CVD-Related Assessment	161
2.8.10	Appropriate Medication Therapy after a Heart Attack	167
2.8.11	Persistence of Appropriate Medication Therapy after a Heart Attack.....	178
2.8.12	Appropriate Medication Therapy in High Risk Patients	190
2.8.13	Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge	201
2.8.14	Cholesterol Management for Patients with Cardiovascular Conditions	204
2.8.15	Heart Failure and Evaluation of LVS Function	207
2.9	STD-Related Measure Topics.....	210
2.9.1	HIV Screening	210
2.9.2	HIV Quality of Care	213
2.9.3	Chlamydia Screening	215
2.9.4	Sexually Transmitted Infection Screening.....	218
2.10	Other Clinical Measures Topics	224
2.10.1	Osteoporosis Management	224
2.10.2	Osteoporosis Screening in Women	227
2.10.3	Rheumatoid Arthritis Medication Monitoring	229
2.10.4	Osteoarthritis Medication Monitoring	234
2.10.5	Asthma.....	237
2.10.6	Asthma Quality of Care	239
2.10.7	Asthma and Inhaled Steroid Use.....	243
2.10.8	Medication Therapy for Persons with Asthma.....	246
2.10.9	Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation.....	249
2.10.10	Chronic Kidney Disease Assessment.....	251
2.10.11	Prediabetes/Metabolic Syndrome.....	253
2.10.12	Proportion of Days Covered by Medication Therapy.....	260
2.10.13	Medications Education	265
2.10.14	Public Health Nursing.....	267
2.10.15	Breastfeeding Rates.....	270
2.10.16	Use of High-Risk Medications in the Elderly	274
2.10.17	Functional Status Assessment in Elders.....	279
2.10.18	Fall Risk Assessment in Elders	281
2.10.19	Palliative Care	283
2.10.20	Annual Wellness Visit.....	286
2.10.21	Goal Setting	287
Contact Information		290

1.0 About This Manual

This manual provides information on the performance measure logic used by the Clinical Reporting System (CRS) Version 11.1 Selected Measures (Local) Report (Fiscal Year [FY] 2011 Clinical Performance Measures). For information on system setup, available reports and steps for running the reports, and performing Area Office functions, refer to the CRS Version 11.1 User Manual.

2.0 Performance Measure Logic

This section provides the following information for each performance measure topic:

- For Government Performance and Results Act (GPRA) measures, the measure description is provided as stated in the Indian Health Service (IHS) Annual Performance Report to Congress
- Definitions of all denominators and numerators for each performance measure topic
- Detailed description of the logic for the denominator and numerator, including specific codes, fields, taxonomies, and/or values searched
- Key changes to logic from the previous year, if any
- Description of which patients and information are contained on the patient list
- Performance measure source and past IHS performance, if any, and IHS or Healthy People (HP) 2020 targets for the performance measure
- Report examples
- Patient list examples

Note: All report examples and patient list examples used in this section were produced from “scrubbed” demonstration databases and do not represent individual patient data.

2.1 Performance Measure Logic Basics

2.1.1 CRS Denominator Definitions

Each performance measure topic has one or more define denominators and numerators. The denominator is the total population that is being reviewed for a specific measure. For the National GPRA & Program Assessment Rating Tool (PART) Report, only one denominator for each topic is reported. These denominators are pre-defined, based on the Active Clinical Population definition. For the Selected Measures reports for local use (CRS Version 11.1 User Manual, Section 5.12), multiple denominators may be reported to provide a complete picture of clinical performance. There are also additional options available to further refine denominator definitions.

2.1.1.1 Denominator Definitions for National GPRA Reporting

The Active Clinical population is the denominator definition used as the basis for *most* GPRA measures. This denominator was developed in FY 2003 specifically for clinical performance measures because it is more representative of the active clinical population.

Note: There are facilities that do not offer direct care. Patients in these facilities receive only Contract Health Services (CHS) and therefore do not meet the requirements of the Active Clinical population. A new site parameter, Contract Health Site Only, was added for these facilities in FY2006.

Prior to FY 2003, the GPRA User Population denominator definition was used for national reporting, similar to the agency's IHS User Population definition.

The *Active Clinical* population for the National GPRA & PART Report is defined by the following criteria:

- Patients with the name of "DEMO,PATIENT" or who are included in the Resource and Patient Management System (RPMS) Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two visits to medical clinics in the past three years. At least one visit must be to one of the following core medical clinics:

01	General	24	Well Child
06	Diabetic	28	Family Practice
10	GYN	57	EPSDT
12	Immunization	70	Women's Health
13	Internal Medicine	80	Urgent Care
20	Pediatrics	89	Evening

The second visit can be either to one of the core medical clinics in the previous list or to one of the following additional medical clinics:

02	Cardiac	37	Neurology
03	Chest And TB	38	Rheumatology
05	Dermatology	49	Nephrology
07	ENT	50	Chronic Disease
08	Family Planning	69	Endocrinology
16	Obstetrics	75	Urology
19	Orthopedic	81	Men's Health Screening
23	Surgical	85	Teen Clinic
25	Other	88	Sports Medicine
26	High Risk	B8	Gastroenterology - Hepatology
27	General Preventive	B9	Oncology - Hematology
31	Hypertension	C3	Colposcopy
32	Postpartum		

- Patient must be alive on the last day of the report period.
- Patient must be American Indian/Alaska Native (AI/AN) (defined as Beneficiary 01). This data item is entered and updated during the patient registration process.
- Patient must reside in a community included in the site's "official" GPRA community taxonomy, defined as all communities of residence in the CHS catchment area specified in the community taxonomy specified by the user.

The *Active Clinical CHS Population* for National GPRA & PART Reports is defined as follows:

- Patients with the name of "DEMO,PATIENT" or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the Patient Care Component [PCC] Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two CHS visits in the three years prior to the end of the report period.
- Patient must be alive on the last day of the report period.
- Patient must be AI/AN (defined as Beneficiary 01). This item is entered and updated during the patient registration process.
- User must reside in a community included in the site's official GPRA community taxonomy, defined as all communities of residence in the CHS catchment area specified in the community taxonomy that is specified by the user.

The *Active Clinical Behavioral Health Population* for National GPRA & PART Reports is defined as follows:

- Patients with the name of “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the Patient Care Component [PCC] Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two behavioral health visits in the three years prior to the end of the report period.
- Patient must be alive on the last day of the report period.
- Patient must be AI/AN (defined as Beneficiary 01). This item is entered and updated during the patient registration process.
- User must reside in a community included in the site’s official GPRA community taxonomy, defined as all communities of residence in the Urban Outreach & Referral catchment area specified in the community taxonomy that is specified by the user.

The *GPRA User Population* for the National GPRA & PART Report is defined by the following criteria:

- Patients with the name of “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have been seen at least once in the three years prior to the end of the time period, regardless of the clinic type.
- Patient must be alive on the last day of the report period.
- Patient must be AI/AN (defined as Beneficiary 01). This data item is entered and updated during the patient registration process.
- Patient must reside in a community included in the site’s “official” GPRA community taxonomy, defined as all communities of residence in the CHS catchment area specified in the community taxonomy specified by the user.

Note: The GPRA User Population definition is similar, but not identical, to the definition used by IHS headquarters (HQ) for annual user population statistics. GPRA “visits” are not required to be workload reportable as defined by IHS HQ.

2.1.1.2 Denominator Definitions for Selected Measures Reports

In addition to the National GPRA & PART Report, CRS provides Selected Measures reports intended for local facility use for specific public health and/or performance improvement initiatives (CRS Version 11.1 User Manual, Section 5.12). Multiple denominators and numerators will be reported for each measure (e.g., *both* Active Clinical and GPRA User Population). Users have additional options to define the denominators as explained below.

The *Active Clinical Population* for Selected Measures (Local) Reports is defined as follows:

- Patients with name “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two visits to medical clinics in the past three years. At least one visit must be to one of the following core medical clinics:

01	General	24	Well Child
06	Diabetic	28	Family Practice
10	GYN	57	EPSDT
12	Immunization	70	Women's Health
13	Internal Medicine	80	Urgent Care
20	Pediatrics	89	Evening

The second visit can be either to one of the core medical clinics in the previous list or to one of the following additional medical clinics:

02	Cardiac	37	Neurology
03	Chest And TB	38	Rheumatology
05	Dermatology	49	Nephrology
07	ENT	50	Chronic Disease
08	Family Planning	69	Endocrinology
16	Obstetrics	75	Urology
19	Orthopedic	81	Men's Health Screening
23	Surgical	85	Teen Clinic
25	Other	88	Sports Medicine
26	High Risk	B8	Gastroenterology - Hepatology
27	General Preventive	B9	Oncology - Hematology
31	Hypertension	C3	Colposcopy
32	Postpartum		

- Patient must be alive on the last day of the Report period.
- User defines population type: AI/AN patients only, non AI/AN, or both. This data item is entered and updated during the patient registration process.
- User defines general population: single community; group of multiple communities (community taxonomy); user-defined list of patient (patient panel); or all patients regardless of community of residence.

The *Active Clinical CHS Population* for Selected Measures (Local) Reports is defined as follows:

- Patients with the name of “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two CHS visits in the three years prior to the end of the Report Period.
- Patient must be alive on the last day of the report period.
- User defines population type: AI/AN patients only, non AI/AN, or both.
- User defines general population: single community; group of multiple communities (community taxonomy); user-defined list of patient (patient panel); or all patients regardless of community of residence.

The *Active Clinical Behavioral Health Population* for Selected Measures (Local) Reports is defined as follows:

- Patients with the name of “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be automatically excluded from the denominator.
- Patient must have two behavioral health visits in the three years prior to the end of the Report Period.
- Patient must be alive on the last day of the report period.
- User defines population type: AI/AN patients only, non AI/AN, or both.
- User defines general population: single community; group of multiple communities (community taxonomy); user-defined list of patient (patient panel); or all patients regardless of community of residence.

The *User Population* for Selected Measures (Local) reports is defined as follows:

- Patients with the name of “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (DPST option located in the PCC Management Reports, Other section) will be excluded from the denominator automatically.
- Patient must have been seen at least once in the three years prior to the end of the time period, regardless of the clinic type.
- Patient must be alive on the last day of the report period.
- User defines population type: AI/AN patients only, non AI/AN, or both.
- User defines general population: single community, group of multiple communities (community taxonomy); user-defined list of patient (patient panel); or all patients regardless of community of residence.

2.1.2 Performance Measure Logic Example

Cancer Screening: Pap Smear Rates: During FY 2011, achieve the tentative target rate of 55.7% for the proportion of female patients ages 21 through 64 without a documented history of hysterectomy who have had a Pap screen within the previous three years.

For CRS, the GPRA measure definition is defined as:

- Denominator (total number of patients evaluated): Active Clinical female patients ages 21 through 64, excluding those with documented history of hysterectomy. (The clinical owner of the measure has determined based on current medical guidelines that “eligible” women are defined as ages 21–64.)
- Numerator (those from the denominator who meet the criteria for the measure): patients with documented Pap smear in past three years or refusal in past year.

For the programmer, the Pap Smear measure is described in terms of the following logic:

1. Begin with the Active Clinical population definition.
 - Exclude any patients with the name of “DEMO,PATIENT.”
 - Exclude any patient records that are included in the RPMS Demo/Test Patient Search Template.
 - Exclude any patients with a date of death in the Patient Registration file.
 - Exclude any patients who do *not* have value 01 (AI/AN) in the Beneficiary field in Patient Registration file.
 - Exclude any patients whose Community of Residence is not included in the site’s defined GPRA Community Taxonomy for this report.

- For the remaining patients, search visit files for the three years prior to the selected report end date; exclude any patients whose visits do not meet the “two medical clinics” definition; *or*, for facilities with the CHS-Only site parameter set to Yes, exclude any patients who do not have two CHS visits in the past three years.
2. From these patients, identify the subset that are female and that are at least age 21 on the first day of the current report period and less than age 65 on the last day of the report period.
 3. Exclude patients with documented hysterectomy by searching the V Procedure file for procedure codes 68.4–68.8 or V CPT for CPT codes 51925, 56308 (old code), 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200–58294, 58548, 58550–58554, 58570–58573, 58951, 58953–58954, 58956, 59135 or V POV 618.5, V88.01, V88.03 or Women's Health procedure called Hysterectomy any time before the end of the report period.
 4. For these patients (the denominator), check for a Pap smear in the past three years in the following order:
 - a. Check V Lab for a lab test called Pap Smear and for any site-populated pap smear lab test documented in the BGP PAP SMEAR TAX taxonomy; *or*
 - b. Check V Lab for any LOINC code listed in the predefined BGP PAP LOINC CODES taxonomy (see the *CRS Technical Manual* for specific codes); *or*
 - c. Check the Purpose of Visit file (V POV) for: a diagnosis of: V67.01 Follow-up Vaginal Pap Smear, V76.2-Screen Mal Neop-Cervix, V72.32 Encounter for Pap Cervical Smear to Confirm Findings of Recent Normal Smear Following Initial Abnormal Smear, V72.3 Gynecological Examination, Pap Cervical Smear as Part of General Gynecological Exam, Pelvic Exam (annual) (periodic) (old code, to be counted for visits prior to 10/1/04 only), V76.47 Vaginal Pap Smear for Post-Hysterectomy Patients, 795.0*, 795.10–16, 795.19; *or*
 - d. Check V Procedures for a procedure of 91.46; *or*
 - e. Check V CPT for the following CPT codes: 88141–88167, 88174–88175, G0123, G0124, G0141, G0143–G0145, G0147, G0148, P3000, P3001, Q0091; *or*
 - f. Check the Women's Health Tracking package for documentation of a procedure called Pap Smear and where the result does *not* have "ERROR/DISREGARD".

If a visit with any of the specified codes is found, the patient is considered to have met the measure, and the program checks the next patient.

2.1.3 Age Ranges

Unless otherwise noted, for the purposes of CRS reports, the age of a patient is calculated at the beginning of the report period. For example, for a report period of July 1, 2010 through June 30, 2011, Jane Doe is defined as age 74 if her birth date is June 10, 1936, even though she becomes age 75 during the report period.

2.1.4 Standard Health Care Codes

2.1.4.1 Current Procedural Terminology Codes

One of several code sets used by the healthcare industry to standardize data, and allow for comparison and analysis. Current Procedural Terminology (CPT) was developed and is updated annually by the American Medical Association, and is widely used in producing bills for services rendered to patients. CPTs include codes for diagnostic and therapeutic procedures, and specify information that differentiates the codes based on cost. CPT codes are the most widely accepted nomenclature in the United States for reporting physician procedures and services for federal and private insurance third-party reimbursement. CRS searches for CPT and other codes as specified in the logic definition to determine if a patient meets a denominator or numerator definition.

2.1.4.2 International Classification of Disease Codes

One of several code sets used by the healthcare industry to standardize data. The International Classification of Disease (ICD) is an international diagnostic coding scheme. In addition to diseases, ICD also includes several families of terms for medical-specialty diagnoses, health status, disablements, procedure, and reasons for contact with healthcare providers. IHS currently uses ICD, Ninth Revision (ICD-9) for coding. CRS searches for ICD and other codes as specified in the logic definition, to determine if a patient meets a denominator or numerator definition.

2.1.4.3 Logical Observation Identifiers Names and Codes

Logical Observation Identifiers Names and Codes (LOINC®). A standard coding system originally initiated for laboratory values, the system is being extended to include nonlaboratory observations (electrocardiograms, vital signs, etc.). Standard code sets are used to define individual tests and mitigate variations in local terminologies for laboratory and other healthcare procedures, for example, Glucose or Glucose Test. IHS began integrating LOINC values into RPMS in several pilot sites in 2002.

Refer to the CRS Version 11.1 Technical Guide for a list of specific LOINC codes included in each LOINC taxonomy.

2.2 Diabetes Related Measure Topics

2.2.1 Diabetes Prevalence

Denominators

All *User Population patients*. Broken down by gender and age groups (<15, 15–19, 20–24, 25–34, 35–44, 45–54, 55–64, >64 yrs).

Numerators

Anyone diagnosed with Diabetes *at any time* before the end of the Report Period.

Anyone diagnosed with Diabetes *during* the Report Period.

Logic Description

Age is calculated at the beginning of the Report Period.

Diabetes definition: At least one diagnosis of 250.00–250.93 recorded in V POV file.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of diabetic patients with most recent diagnosis.

Measure Source

HP 2010 5–2, 5–3

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	12.0%
IHS FY 2009 Performance	12.0%
IHS FY 2008 Performance	12.0%
IHS FY 2007 Performance	11.0%
IHS FY 2006 Performance	11.0%
IHS FY 2005 Performance	11.0%
IHS FY 2004 Performance	10.0%

DU

May 25, 2011

Page 1

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes Prevalence								
Denominator(s):								
- All User Population patients. Broken down by gender and by age groups: < 15, 15-19, 20-24, 25-34, 35-44, 45-54, 55-64, > 64.								
Numerator(s):								
- Anyone diagnosed with Diabetes at any time before the end of the Report Period.								
- Anyone diagnosed with Diabetes during the Report Period.								
Logic:								
Age is calculated at the beginning of the Report Period. Diabetes diagnosis is defined as at least one diagnosis 250.00-250.93 recorded in the V POV file.								
Performance Measure Description:								
Continue tracking (i.e., data collection and analyses) Area age-specific diabetes prevalence rates to identify trends in the age-specific prevalence of diabetes (as a surrogate marker for diabetes incidence) for the AI/AN population.								
Past Performance and/or Target:								
IHS Performance: FY 2010 - 12%, FY 2009 - 12%, FY 2008 - 12%, FY 2007 - 11%, FY 2006 - 11%, FY 2005 - 11%, FY 2004 - 10%								
Source:								
HP 2010 5-2, 5-3								
	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
# User Pop	2,896		2,456			2,346		
# w/ any DM DX	265	9.2	243	9.9	-0.7	197	8.4	+0.8
# w/ DM DX w/in past year	173	6.0	146	5.9	+0.0	100	4.3	+1.7
# Male User Pop	1,368		1,152			1,109		
# w/ any DM DX	114	8.3	104	9.0	-0.7	72	6.5	+1.8
# w/DM DX w/in past year	82	6.0	78	6.8	-0.8	48	4.3	+1.7
# Female User Pop	1,528		1,304			1,237		
# w/ any DM DX	151	9.9	139	10.7	-0.8	125	10.1	-0.2
# w/ DM DX w/in past year	91	6.0	68	5.2	+0.7	52	4.2	+1.8

Figure 2-1: Sample Summary Report, Diabetes Prevalence Topic

DU	May 25, 2011	Page 2
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		

Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes Prevalence (con't)								
TOTAL USER POPULATION								
Age Distribution								
	<15	15-19	20-24	25-34	35-44	45-54	55-64	>64 yrs
CURRENT REPORT PERIOD								
Total # User Pop	730	239	261	403	389	390	262	222
# w/ DM DX ever	1	3	7	37	57	67	50	43
% w/ DM DX ever	0.1	1.3	2.7	9.2	14.7	17.2	19.1	19.4
# w/DM DX in past yr	0	2	3	14	42	49	34	29
% w/DM DX in past yr	0.0	0.8	1.1	3.5	10.8	12.6	13.0	13.1
PREVIOUS YEAR PERIOD								
Total # User Pop	711	227	243	352	316	277	181	149
# w/ DM DX ever	3	4	9	31	54	57	45	40
% w/ DM DX ever	0.4	1.8	3.7	8.8	17.1	20.6	24.9	26.8
# w/DM DX in past yr	1	3	3	9	33	37	31	29
% w/DM DX in past yr	0.1	1.3	1.2	2.6	10.4	13.4	17.1	19.5
CHANGE FROM PREV YR %								
w/ DM DX ever	-0.3	-0.5	-1.0	+0.4	-2.4	-3.4	-5.8	-7.5
w/DM DX in past yr	-0.1	-0.5	-0.1	+0.9	+0.4	-0.8	-4.1	-6.4
BASELINE REPORT PERIOD								
Total # User Pop	787	208	217	329	293	228	141	143
# w/ DM DX ever	2	4	12	20	38	46	31	44
% w/ DM DX ever	0.3	1.9	5.5	6.1	13.0	20.2	22.0	30.8
# w/DM DX in past yr	2	1	3	7	18	21	20	28
% w/DM DX in past yr	0.3	0.5	1.4	2.1	6.1	9.2	14.2	19.6
CHANGE FROM BASE YR %								
w/ DM DX ever	-0.1	-0.7	-2.8	+3.1	+1.7	-3.0	-2.9	-11.4
w/DM DX in past yr	-0.3	+0.4	-0.2	+1.3	+4.7	+3.4	-1.2	-6.5

Figure 2-2: Sample Age Breakdown Page, Diabetes Prevalence Topic

DU	May 25, 2011					Page 1	
*** IHS 2011 Clinical Performance Measure Patient List ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Entire Patient List							

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic							
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Diabetes Prevalence: List of diabetic patients with most recent diagnosis							

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	UP	01/28/11	250.00
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP	03/21/11	250.00
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP	02/12/11	250.80

PATIENT4 ,WINONA	000004	COMMUNITY	#1	F	53	UP	04/21/11	250.80
PATIENT5 ,NADINE	000005	COMMUNITY	#1	F	61	UP	02/01/11	250.00
PATIENT6 ,RUTH	000006	COMMUNITY	#1	F	64	UP	08/09/11	250.00

Figure 2-3: Sample Patient List, Diabetes Prevalence, Patients with Diabetes Diagnosis

2.2.2 Diabetes Comprehensive Care

Denominators

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two diabetes mellitus- (DM-) related visits ever.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two diabetes mellitus- (DM-) related visits ever, without a documented history of bilateral foot amputation or two separate unilateral foot amputations.

Numerators

Patients with hemoglobin A1c documented during the Report Period, regardless of result

Patients with blood pressure (BP) documented during the Report Period

Patients with controlled BP, defined as < 130/80, i.e., the mean systolic value is less than 130 and the mean diastolic value is less than 80

Patients with low-density lipoprotein (LDL) completed during the Report Period, regardless of result

Patients with nephropathy assessment, defined as an estimated GFR with result and a quantitative urinary protein assessment during the Report Period *or* with the evidence of diagnosis and/or treatment of end-stage renal disease (ESRD) at any time before the end of the Report Period

Patients receiving a qualified retinal evaluation during the Report Period

Note: This numerator does *not* include refusals.

Patients with diabetic foot exam during the Report Period

Note: This numerator does *not* include refusals.

Patients with comprehensive diabetes care (Documented A1c *and* Blood Pressure *and* LDL *and* Nephropathy Assessment *and* Retinal Exam *and* Diabetic Foot Exam).

Logic Description

Diabetes definition: First Purpose of Visit (POV) 250.00–250.93 recorded in the POV file prior to the Report Period.

A1c definition: Searches for most recent A1c test with a result during the Report Period. If none found, CRS searches for the most recent A1c test without a result. A1c defined as: CPT 83036, 83037, 3044F–3046F, 3047F (old code); LOINC taxonomy; or site-populated taxonomy DM AUDIT HGB A1C TAX.

BP documented definition: Having a minimum of two BPs documented on non-Emergency Room (ER) visits during the Report Period.

CRS uses mean of last three BPs documented on non-ER visits during the Report Period. If three BPs are not available, uses mean of last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) BPs and dividing by three (or two).

If CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F–3080F documented on a non-ER visit during the Report Period.

Controlled BP definition: CRS uses a mean, as described above. If the mean systolic and diastolic values do not *both* meet the criteria for controlled, then the value is considered not controlled.

BP documented and Controlled BP: If CRS is not able to calculate a mean BP from BP measurements, it will search for the most recent of any of the following CPT codes documented on non-ER visits during the Report Period: BP Documented: 0001F or 2000F; *or* Systolic 3074F, 3075F, or 3077F *with* Diastolic: 3078F, 3079F, or 3080F. The systolic and diastolic values do not have to be recorded on the same day. If there are multiple values on the same day, the CPT indicating the lowest value will be used. The following combination represents BP < 130/80 and will be included in the Controlled BP numerator: CPT 3074F *and* 3078F. All other combinations *will not* be included in the Controlled BP numerator.

LDL definition: Finds last test done during the Report Period; defined as CPT 80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F; LOINC taxonomy; site-populated taxonomy DM AUDIT LDL CHOLESTEROL TAX.

Nephropathy assessment definition: (1) Estimated GFR with result during the report period, defined as any of the following: (A) Site-populated taxonomy BGP GPRA ESTIMATED GFR TAX or (B) LOINC taxonomy, *and* (2) Quantitative Urinary

Protein Assessment during the Report Period, defined as any of the following: (A) CPT 82042, 82043, or 84156; (B) LOINC taxonomy; or (C) site-populated taxonomy BGP QUANT URINE PROTEIN.

Note: Be sure and check with your laboratory supervisor that the names you add to your taxonomy reflect quantitative test values.

(3) ESRD diagnosis/treatment defined as any of the following ever: (A) V CPT 36145, 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831–36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951–90970 or old codes 90918–90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, G0257, G0308–G0327 (old codes), G0392 (old code), G0393 (old code), or S9339; (B) V POV 585.5, 585.6, V42.0, V45.1 (old code), V45.11, V45.12, or V56.*; (C) V Procedure 38.95, 39.27, 39.42, 39.43, 39.53, 39.93–39.95, 54.98, or 55.6*.

Qualified retinal evaluation definition: (1) diabetic retinal exam or (2) other eye exam. The following methods are qualifying for this measure: (1) dilated retinal evaluation by an optometrist or ophthalmologist, or (2) seven standard fields stereoscopic photos (ETDRS) evaluated by an optometrist or ophthalmologist, or (3) any photographic method formally validated to seven standard fields (ETDRS).

- *Diabetic Retinal Exam:* Any of the following during the Report Period: (1) Exam Code 03 Diabetic Eye Exam (dilated retinal examination or validated photographic equivalent), (2) CPT 2022F Dilated retinal eye exam; 2024F Seven standard field stereoscopic photos with interpretation by an ophthalmologist or optometrist; 2026F Eye imaging validated to match the diagnosis from seven standard field stereoscopic photos; S0620 Routine ophthalmological examination including refraction; new patient; S0621 Routine ophthalmological examination including refraction; established patient; S3000 Diabetic indicator; retinal eye exam, dilated, bilateral.
- *Other Eye Exam:* (1) Non-DNKA (did not keep appointment) visits to ophthalmology, optometry or validated teleophthalmology retinal evaluation clinics or (2) non-DNKA visits to an optometrist or ophthalmologist. Searches for the following codes in the following order: Clinic Codes A2, 17, 18, 64; Provider Code 24, 79, 08; CPT 67028, 67038 (old code), 67039, 67040, 92002, 92004, 92012, 92014; POV V72.0; Procedure 95.02.

Diabetic foot exam definition: (1) Exam Code 28 Diabetic Foot Exam, Complete; (2) non-DNKA visit with a podiatrist (Provider Codes 33, 84 or 25), (3) non-DNKA visit to Podiatry Clinic (Clinic Code 65), or (4) CPT 2028F.

Bilateral foot amputation definition: CPT 27290.50-27295.50, 27590.50-27592.50, 27598.50, 27880.50-27882.50 (.50 modifier indicates bilateral)

Unilateral foot amputation definition: Must have 2 separate occurrences for either CPT or Procedure codes on 2 different dates of service: (1) CPT 27290-27295, 27590-27592, 27598, 27880-27882, or (2) ICD Procedure codes 84.10, 84.13-84.19.

Key Logic Changes from CRS Version 11.0

1. Excluded patients with bilateral foot amputations or two separate unilateral foot amputations from the denominator for diabetic foot exam.
2. Added CPT code 36147 to ESRD definition.
3. Added CPT codes 0001F and 2000F to BP Documented definition.
4. For Other Eye Exam, noted that CPT code 67038 is an old code.
5. Added codes 62238-1 and 50110-4 to LOINC taxonomy for Estimated GFR.
6. Added codes 55454-3, 59261-8, and 62388-4 to LOINC taxonomy for A1c.
7. Added code 43396-1 to LOINC taxonomy for LDL.
8. Added codes 56553-1, 57369-1, 58448-2, 58992-9, and 59159-4 to LOINC taxonomy for Quantitative Urine Protein.

Patient List Description

List of diabetic patients with documented tests, if any.

Measure Source

Foot Exam: HP 2020 D-9

Measure Past Performance and Long-Term Targets

Target	Percent
HP 2020 goal for foot exam	74.8%

DU	May 25, 2011						Page 9
*** IHS 2011 Selected Measures with Community Specified Report ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Previous Year Period: Jan 01, 2010 to Dec 31, 2010							
Baseline Period: Jan 01, 2000 to Dec 31, 2000							

Diabetes Comprehensive Care							
	REPORT	%	PREV YR	%	CHG from	BASE	%
	PERIOD		PERIOD		PREV YR %	PERIOD	BASE %
Active Diabetic Pts	144		97			87	
# w/Comp Diabetes							

Care	10	6.9	1	1.0	+5.9	0	0.0	+6.9
# w/A1c done								
w/ or w/o result	92	63.9	70	72.2	-8.3	52	59.8	+4.1
# w/ BPs documented	120	83.3	78	80.4	+2.9	74	85.1	-1.7
# w/Controlled BP								
<130/80	28	19.4	20	20.6	-1.2	13	14.9	+4.5
# w/ LDL done	78	54.2	46	47.4	+6.7	23	26.4	+27.7
# w/ est GFR & quant								
UP assmt or								
w/ESRD	53	36.8	11	11.3	+25.5	6	6.9	+29.9
# w/Retinal Evaluation								
-No Refusals	54	37.5	39	40.2	-2.7	44	50.6	-13.1
Active Diabetic Pts w/o								
Hx of Bilateral								
Amputation	137		97			87		
# w/Diabetic Foot Exam								
-No Refusals	20	14.6	18	18.6	-4.0	16	18.4	-3.8

Figure 2-4: Sample Summary Report, Diabetes Comprehensive Care Topic

CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT

DU

May 25, 2011

Page 11

IHS 2011 Clinical Performance Measure Patient List

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic

PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetes Comprehensive Care: List of Diabetic patients with documented tests, if any

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	AD	A1c: 02/28/11
6.6;BPs: 133/82 UNC;LDL: 10/28/11 119;EYE: 01/07/11 Cl: 18						
PATIENT2,TARA	000002	COMMUNITY #1	F	51	AD	BP: <130/80: BPs:
118/61; ESRD: 03/03/03 90951; FOOT AMPUTATION						
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	AD	A1c: 04/09/11
6.5;BPs: 138/66 UNC;GFR: 09/09/11 & QUANT UP: QUANT URINE PROTEIN-03/31/11;EYE: 07/30/11 Cl: 18;FOOT EXAM: 01/07/11 Cl: 65						

Figure 2-5: Sample Patient List, Diabetes Comprehensive Care

2.2.3 Diabetes: Glycemic Control

GPRA Measure Description, Poor Glycemic Control

During FY 2011, achieve the target rate of 19.4% for the proportion of patients with diagnosed diabetes who have poor glycemic control (defined as A1c > 9.5).

GPRA Measure Description, Ideal Glycemic Control

During FY 2011, achieve the target rate of 30.2% for the proportion of patients with diagnosed diabetes who have ideal glycemic control (defined as A1c < 7).

Denominators

All User Population patients diagnosed with diabetes prior to the report period.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, and at least two visits during the Report Period, and two DM-related visits ever. (GPRA Denominator)

Active Adult Diabetic patients, defined by meeting the following criteria: (1) who are age 19 or older at the beginning of the Report Period, (2) whose first ever DM diagnosis occurred prior to the Report Period; (3) who had at least two DM related visits ever; (4) at least one encounter with DM POV in a primary clinic with a primary provider during the Report Period; and (5) never have had a creatinine value greater than five.

Numerators

Number of patients with a Hemoglobin A1c documented during the Report Period, regardless of result.

Poor Control: Total of Poor and Very Poor Control: Patients with A1c greater than (>) 9.5. (GPRA Numerator)

Very Poor Control: Patients with A1c equal to or greater than (=>) 12.

Poor Control: Patients with A1c greater than (>) 9.5 and less than (<) 12.

Fair Control: Patients with A1c equal to or greater than (=>) 8 and less than or equal to (<=) 9.5.

Good Control: Patients with A1c equal to or greater than (=>) 7 and less than (<) 8.

Ideal Control: Patients with A1c less than (<) 7. (GPRA Numerator)

Without Result: Patients with A1c documented but no value.

Logic Description

Diabetes definition: First Purpose of Visit 250.00–250.93 recorded in the V POV file prior to the Report Period.

Hemoglobin A1c definition: Searches for most recent A1c test with a result during the Report Period. If more than one A1c test is found on the same day and/or the same visit and one test has a result and the other does not, the test with the result will be used. If an A1c test with a result is not found, CRS searches for the most recent A1c test without a result. Without result is defined as A1c documented but with no value.

CRS uses the following definitions:

	CPT Codes	LOINC Codes	Taxonomy
Creatinine (for Active Adult Diabetic denominator)	CPT codes are not included since they do not store the result, which is used in this topic	Yes	DM AUDIT CREATININE TAX
Hemoglobin A1c	83036, 83037, 3044F-3046F, 3047F (old code) Note: CPT 3044F represents A1c <7 and will be included in the Ideal Control numerator.	Yes	DM AUDIT HGB A1C TAX

In the CPT Codes column, specific LOINC codes used CRS are located in the *CRS Technical Manual*.

Key Logic Changes from CRS Version 11.0

1. Added codes 55454-3, 59261-8, and 62388-4 to LOINC taxonomy for A1c.

Patient List Description

List of diabetic patients with most recent A1c value, if any.

Measure Source

HEDIS; HP 2020 D-11, D-5

Measure Past Performance and Long-Term Targets

Hemoglobin A1c Documented

Performance	Percent
IHS FY 2010 Performance	82.0%
IHS FY 2009 Performance	80.0%
IHS FY 2008 Performance	79.0%
IHS FY 2007 Performance	79.0%
IHS FY 2006 Performance	79.0%
IHS FY 2005 Performance	78.0%
IHS FY 2004 Performance	77.0%
IHS FY 2003 Performance	75.0%

Performance	Percent
IHS FY 2002 Performance	73.0%
HP 2020 Goal	71.1%

Poor Glycemic Control

Performance	Percent
IHS FY 2010 Performance	18.0%
IHS FY 2009 Performance	18.0%
IHS FY 2008 Performance	17.0%
IHS FY 2007 Performance	16.0%
IHS FY 2006 Performance	16.0%
IHS FY 2005 Performance	15.0%
IHS FY 2004 Performance	17.0%
IHS FY 2003 Performance	17.0%
IHS FY 2002 Performance	18.0%

Ideal Glycemic Control

Performance	Percent
IHS FY 2010 Performance	32.0%
IHS FY 2009 Performance	31.0%
IHS FY 2008 Performance	32.0%
IHS FY 2007 Performance	31.0%
IHS FY 2006 Performance	31.0%
IHS FY 2005 Performance	30.0%
IHS FY 2004 Performance	27.0%
IHS FY 2003 Performance	28.0%
IHS FY 2002 Performance	25.0%
HP 2020 Goal	58.9%

DU	May 25, 2011						Page 11	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes: Glycemic Control								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
User Pop w/ DM DX								
prior to report								
end date	229		202			180		
# w/A1c done w/								
or w/o result	95	41.5	72	35.6	+5.8	53	29.4	+12.0
# w/A1c =>12	3	1.3	1	0.5	+0.8	3	1.7	-0.4

# w/A1c >9.5 and <12	15	6.6	3	1.5	+5.1	8	4.4	+2.1
# w/A1c =>8 and =<9.5	12	5.2	19	9.4	-4.2	10	5.6	-0.3
# w/A1c=>7 and <8	13	5.7	16	7.9	-2.2	7	3.9	+1.8
# w/A1c <7	38	16.6	33	16.3	+0.3	23	12.8	+3.8
# w/A1c w/o Result	14	6.1	0	0.0	+6.1	2	1.1	+5.0
Active Diabetic Pts (GPRA)	144		97			87		
# w/A1c done w/ or w/o result	92	63.9	70	72.2	-8.3	52	59.8	+4.1
# w/A1c > 9.5 (GPRA)	18	12.5	4	4.1	+8.4	11	12.6	-0.1
# w/A1c =>12	3	2.1	1	1.0	+1.1	3	3.4	-1.4
# w/A1c >9.5 and < 12	15	10.4	3	3.1	+7.3	8	9.2	+1.2
# w/A1c =>8 and =<9.5	12	8.3	19	19.6	-11.3	10	11.5	-3.2
# w/A1c=>7 and <8	13	9.0	16	16.5	-7.5	7	8.0	+1.0
# w/A1c <7 (GPRA)	35	24.3	31	32.0	-7.7	22	25.3	-1.0
# w/A1c w/o Result	14	9.7	0	0.0	+9.7	2	2.3	+7.4
Active Adult Diabetic Patients	102		75			63		
# w/A1c done w/ or w/o result	71	69.6	61	81.3	-11.7	46	73.0	-3.4
# w/A1c =>12	3	2.9	1	1.3	+1.6	3	4.8	-1.8
# w/A1c >9.5 and <12	13	12.7	2	2.7	+10.1	7	11.1	+1.6
# w/A1c =>8 and =<9.5	11	10.8	18	24.0	-13.2	8	12.7	-1.9
# w/A1c=>7 and <8	10	9.8	12	16.0	-6.2	6	9.5	+0.3
# w/A1c <7	28	27.5	28	37.3	-9.9	22	34.9	-7.5
# w/A1c w/o Result	6	5.9	0	0.0	+5.9	0	0.0	+5.9

Figure 2-6: Sample Report, Diabetes: Glycemic Control Topic

***** CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT *****
DU May 25, 2011 Page 16
*** IHS 2011 Clinical Performance Measure Patient List ***
DEMO INDIAN HOSPITAL
Report Period: Jan 01, 2011 to Dec 31, 2011
Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetes: Glycemic Control: List of diabetic patients with most recent Alc value, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DEBORA	000001	COMMUNITY #1	F	45	UP,AD,AAD	03/28/11 6.6
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP,AD,AAD	02/20/11 12.4
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP,AD,AAD	04/09/11 6.5
PATIENT4,WINONA	000004	COMMUNITY #1	F	53	UP	
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	UP,AD,AAD	02/01/11 6.5
PATIENT6,RUTH	000006	COMMUNITY #1	F	64	UP	

Figure 2-7: Sample Patient List, Diabetes: Glycemic Control

2.2.4 Diabetes: Blood Pressure Control

GPRA Measure Description

During FY 2011, achieve the target rate of 35.9% for the proportion of patients with diagnosed diabetes who have achieved blood pressure control (defined as <130/80).

Denominators

All User Population patients diagnosed with diabetes prior to the Report Period.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, and at least two visits during the Report Period, and two DM-related visits ever. (GPRA Denominator)

Active Adult Diabetic patients, defined by meeting the following criteria: (1) who are age 19 or older at the beginning of the Report Period, (2) whose first ever DM diagnosis occurred prior to the Report Period; (3) who had at least two DM-related visits ever, (4) at least one encounter with DM POV in a primary clinic with a primary provider during the Report Period; and (5) never have had a creatinine value greater than five.

Numerators

Patients with BP documented during the report period

Patients with controlled BP, defined as < 130/80, i.e., the mean systolic value is less than 130 and the mean diastolic value is less than 80 (GPRA Numerator)

Patients with BP that is not controlled

Logic Description

Diabetes definition: First DM POV 250.00–250.93 recorded in the V POV file prior to the Report Period.

BP documented definition: CRS uses mean of last three BPs documented on non-ER visits during the Report Period. If three BPs are not available, uses mean of the last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) blood pressures and dividing by three (or two). If the systolic and diastolic values do not *both* meet the criteria for controlled, then the value is considered not controlled.

If CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F-3080F documented on a non-ER visit during the Report Period.

Controlled BP definition: CRS uses a mean, as described above where BP is <130/80. If the mean systolic and diastolic values do not *both* meet the criteria for controlled, then the value is considered not controlled.

BP documented and Controlled BP: If CRS is not able to calculate a mean BP from blood pressure measurements, it will search for the most recent of any of the following CPT codes documented on non-ER visits during the Report Period: BP Documented: 0001F or 2000F; *or* Systolic: 3074F, 3075F or 3077F *with* Diastolic: 3078F, 3079F, or 3080F. The systolic and diastolic values do *not* have to be recorded on the same day. If there are multiple values on the same day, the CPT indicating the lowest value will be used. The following combination represents BP <130/80 and will be included in the Controlled BP numerator: CPT 3074F *and* 3078F. All other combinations will *not* be included in the Controlled BP numerator.

CRS uses the following definition:

	CPT Codes	LOINC Codes	Taxonomy
Creatinine (for Active Adult Diabetic denominator)	CPT codes are not included since they do not store the result, which is used in this topic	Yes	DM AUDIT CREATININE TAX

In the LOINC Codes column, specific LOINC codes by CRS are location in the *CRS Technical Manual*.

Key Logic Changes from CRS Version 11.0

1. Added CPT codes 0001F and 2000F to BP Documented definition.

Patient List Description

List of diabetic patients with BP value, if any.

Measure Source

HP 2020 D-7

Measure Past Performance and Long-Term Targets**Controlled BP**

Performance	Percent
IHS FY 2010 Performance	38.0%
IHS FY 2009 Performance	37.0%
IHS FY 2008 Performance	38.0%
IHS FY 2007 Performance	39.0%
IHS FY 2006 Performance	37.0%
IHS FY 2005 Performance	37.0%
IHS FY 2004 Performance	35.0%
IHS FY 2003 Performance	37.0%
IHS FY 2002 Performance	36.1%
<i>HP 2020 Goal</i>	<i>57.0%</i>

BP Assessed

Performance	Percent
IHS FY 2008 Performance	89.0%
IHS FY 2005 Performance	89.0%

DU	May 25, 2011						Page 13	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes: Blood Pressure Control								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
User Pop w/ DM DX prior to report period	229		202			180		
# w/ BPs Documented	132	57.6	88	43.6	+14.1	84	46.7	+11.0
# w/controlled BP < 130/80	32	14.0	24	11.9	+2.1	18	10.0	+4.0
# w/Not controlled BP	100	43.7	64	31.7	+12.0	66	36.7	+7.0
Active Diabetic Pts (GPRA)	144		97			87		
# w/ BPs Documented	120	83.3	78	80.4	+2.9	74	85.1	-1.7
# w/Controlled BP								

< 130/80 (GPRA)	28	19.4	20	20.6	-1.2	13	14.9	+4.5
# w/Not controlled BP	92	63.9	58	59.8	+4.1	61	70.1	-6.2
Active Adult Diabetic Patients	102		75			63		
# w/ BPs Documented	82	80.4	61	81.3	-0.9	56	88.9	-8.5
# w/Controlled BP < 130/80	20	19.6	14	18.7	+0.9	8	12.7	+6.9
# w/Not controlled BP	62	60.8	47	62.7	-1.9	48	76.2	-15.4

Figure 2-8: Sample Report, Diabetes: Blood Pressure Control Topic

***** CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT *****

DU

May 25, 2011

Page 23

*** IHS 2011 Clinical Performance Measure Patient List ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetes: Blood Pressure Control: List of diabetic patients with BP value, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	UP,AD,AAD	133/82 UNC
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP,AD,AAD	04/21/11 CPT 3080F
DIASTOLIC BP >=90 UNC						
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP,AD,AAD	138/66 UNC
PATIENT4,WINONA	000004	COMMUNITY #1	F	53	UP	unknown
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	UP,AD,AAD	159/86 UNC
PATIENT6,RUTH	000006	COMMUNITY #1	F	64	UP	139/74 UNC

Figure 2-9: Sample Patient List, Diabetes: Blood Pressure Control

2.2.5 Diabetes: LDL Assessment

GPRA Measure Description

During FY 2011, achieve the target rate of 63.3% for the proportion of patients with diagnosed diabetes who are assessed for dyslipidemia (LDL cholesterol).

Denominators

All *User Population* patients diagnosed with diabetes prior to the Report Period.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, and at least two visits during the Report Period, and two DM-related visits ever. (GPRA Denominator)

Active Adult Diabetic patients, defined by meeting the following criteria: (1) who are age 19 or older at the beginning of the Report Period, (2) whose first ever DM diagnosis occurred prior to the Report Period; (3) who had at least two DM related visits ever, (4) at least one encounter with DM POV in a primary clinic with a primary provider during the Report Period; and (5) never have had a creatinine value greater than five.

Numerators

Patients with LDL completed during the Report Period, regardless of result. (GPRA Numerator)

Patients with *LDL results* less than (<) 130.

- a. Patients with LDL results less than or equal to (<=) 100
- b. Patients with LDL results 101–129

Logic Description

Diabetes definition: First DM POV 250.00–250.93 recorded in the V POV file prior to the Report Period.

LDL definition: Searches for most recent LDL test with a result during the Report Period. If more than one LDL test is found on the same day and/or the same visit and one test has a result and the other does not, the test with the result will be used. If an LDL test with a result is not found, CRS searches for the most recent LDL test without a result.

CRS uses the following to define the tests:

	CPT Codes	LOINC Codes	Taxonomy
Creatinine (for Active Adult Diabetic denominator)	CPT codes are not included since they do not store the result, which is used in this topic	Yes	DM AUDIT CREATININE TAX
LDL Done	80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F	Yes	DM AUDIT LDL CHOLESTEROL TAX

	CPT Codes	LOINC Codes	Taxonomy
LDL <130	3048F, 3049F		Tests in above taxonomy with LDL <130
LDL =<100	3048F		Tests in above taxonomy with LDL =<100

In the LOINC Codes column, specific LOINC codes used by CRS are located in the *CRS Technical Manual*.

Key Logic Changes from CRS Version 11.0

1. Added code 43396-1 to LOINC taxonomy for LDL.

Patient List Description

List of diabetic patients with documented LDL cholesterol test, if any.

Measure Source

HP 2010 12–15

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	67.0%
IHS FY 2009 Performance	65.0%
IHS FY 2008 Performance	63.0%
IHS FY 2007 Performance	61.0%
IHS FY 2006 Performance	60.0%
IHS FY 2005 Performance	53.0%
IHS FY 2004 Performance	53.0%
IHS FY 2003 Performance	47.5%
IHS FY 2002 Performance	43.7%

DU	May 25, 2011						Page 15	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes: LDL Assessment								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
# w/ LDL done	84	36.7	48	23.8	+12.9	23	12.8	+23.9
# w/LDL <130	58	25.3	40	19.8	+5.5	15	8.3	+17.0

A. # w/LDL =<100	37	16.2	31	15.3	+0.8	8	4.4	+11.7
B. # w/LDL 101-129	17	7.4	9	4.5	+3.0	7	3.9	+3.5
Active Diabetic Pts (GPRA)	144		97			87		
# w/ LDL done (GPRA)	78	54.2	46	47.4	+6.7	23	26.4	+27.7
# w/LDL <130	54	37.5	38	39.2	-1.7	15	17.2	+20.3
A. # w/LDL =<100	35	24.3	30	30.9	-6.6	8	9.2	+15.1
B. # w/LDL 101-129	15	10.4	8	8.2	+2.2	7	8.0	+2.4
Active Adult Diabetic Patients	102		75			63		
# w/ LDL done	60	58.8	43	57.3	+1.5	21	33.3	+25.5
# w/LDL <130	41	40.2	35	46.7	-6.5	13	20.6	+19.6
A. # w/LDL =<100	26	25.5	26	34.7	-9.2	8	12.7	+12.8
B. # w/LDL 101-129	12	11.8	9	12.0	-0.2	5	7.9	+3.8

Figure 2-10: Sample Report, Diabetes: LDL Assessment

***** CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT *****

DU

May 25, 2011

Page 30

*** IHS 2011 Clinical Performance Measure Patient List ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetes: LDL Assessment: List of diabetic patients with documented LDL
cholesterol test, if any

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	UP,AD,AAD	LDL DONE: 03/28/11
119						
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP,AD,AAD	LDL DONE: 02/20/11
86						
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP,AD,AAD	
PATIENT4,WINONA	000004	COMMUNITY #1	F	53	UP	
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	UP,AD,AAD	LDL DONE: 02/06/11
CPT 3048F LDL<100						
PATIENT6,RUTH	000006	COMMUNITY #1	F	64	UP	LDL DONE: 05/21/11
107						

Figure 2-11: Sample Patient List, Diabetes: LDL Assessment

2.2.6 Diabetes: Nephropathy Assessment

GPRA Measure Description

During FY 2011, achieve the target rate of 51.9% for the proportion of patients with diagnosed diabetes who are assessed for nephropathy.

Denominators

All User Population patients diagnosed with diabetes prior to the Report Period.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, and at least two visits during the Report Period, and two DM-related visits ever. (GPRA Denominator)

Active Adult Diabetic patients, defined by meeting the following criteria: (1) who are aged 19 or older at the beginning of the Report Period; (2) whose first ever DM diagnosis occurred prior to the Report Period; (3) who had at least two DM related visits ever; (4) at least one encounter with DM POV in a primary clinic with a primary provider during the Report Period; and (5) never have had a creatinine value greater than five.

Numerator

Patients with nephropathy assessment, defined as an estimated GFR with result and a quantitative urinary protein assessment during the Report Period or with evidence of diagnosis and/or treatment of ESRD at any time before the end of the Report Period. (GPRA Numerator)

Logic Description

Diabetes definition: First DM Purpose of Visit 250.00-250.93 recorded in the V POV file prior to the Report period.

Nephropathy assessment definition:

- Estimated GFR with result during the Report Period and Quantitative Urinary Protein Assessment during the Report Period, *or*
- ESRD diagnosis/treatment defined as any diagnosis ever.

CRS uses the following to define the tests/diagnoses:

	CPT Codes	LOINC Codes	Taxonomy
Creatinine (for Active Adult Diabetic Denominator)	CPT codes are not included since they do not store the result, which is used in this topic	Yes	DM AUDIT CREATININE TAX
Estimated GFR		Yes	BGP GPRA ESTIMATED GFR TAX
Quantitative Urinary Protein Assessment	CPT: 82042-82043, 84156	Yes	BGP QUANT URINE PROTEIN TAX Note: Be sure and check with your laboratory supervisor that the names you add to your taxonomy reflect quantitative test values
End Stage Renal Disease	V CPT: 36145, 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831-36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951-90970 or old codes 90918-90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, G0257, G0308-G0327 (old codes), G0392 (old code), G0393 (old code), or S9339 V POV: 585.5, 585.6, V42.0, V45.1 (old code), V45.11, V45.12, or V56* V Procedure: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93-39.95, 54.98, or 55.6*		

In the LOINC Codes column, specific LOINC codes used by CRS are located in the *CRS Technical Manual*.

Key Logic Changes from CRS Version 11.0

1. Added CPT code 36147 to ESRD definition.
2. Added codes 62238-1 and 50110-4 to LOINC taxonomy for Estimated GFR.
3. Added codes 56553-1, 57369-1, 58448-2, 58992-9, and 59159-4 to LOINC taxonomy for Quantitative Urine Protein.

Patient List Description

List of diabetic patients with nephropathy assessment, if any.

Measure Source

HP 2010 5-11

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	55.0%
IHS FY 2009 Performance	50.0%
IHS FY 2008 Performance	50.0%
IHS FY 2007 Performance (new baseline established; revised standards of care resulted in revised measure definition)	40.0%
IHS FY 2006 Performance (measure definition was different from current definition)	55.0%
IHS FY 2005 Performance (measure definition was different from current definition)	47.0%
IHS FY 2004 Performance (measure definition was different from current definition)	42.0%
IHS FY 2003 Performance (measure definition was different from current definition)	37.5%
IHS FY 2002 Performance (measure definition was different from current definition)	35.0%

DU	May 25, 2011						Page 17	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes: Nephropathy Assessment								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
User Pop w/ DM DX prior to Report Period	229		202			180		
# w/ est GFR & quant UP assmt or w/ESRD	55	24.0	16	7.9	+16.1	8	4.4	+19.6
Active Diabetic Pts (GPRA)	144		97			87		
# w/ est GFR & quant UP assmt or w/ESRD (GPRA)	53	36.8	11	11.3	+25.5	6	6.9	+29.9

Active Adult Diabetic Patients	102		75			63		
# w/ est GFR & quant UP assmt or w/ESRD	41	40.2	6	8.0	+32.2	4	6.3	+33.8

Figure 2-12: Sample Report, Diabetes: Nephropathy Assessment

***** CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT *****

DU

May 25, 2011

Page 37

*** IHS 2011 Clinical Performance Measure Patient List ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetes: Nephropathy Assessment: List of diabetic patients with nephropathy
assessment, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	UP,AD,AAD	
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP,AD,AAD	
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP,AD,AAD	;GFR: 05/09/11 &
QUANT UP: QUANT URINE PROTEIN-03/31/11						
PATIENT4,WINONA	000004	COMMUNITY #1	F	53	UP	
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	UP,AD,AAD	ESRD: ESRD 36145-01/30/11
PATIENT6,RUTH	000006	COMMUNITY #1	F	64	UP	
PATIENT7,DANIELLE	000007	COMMUNITY #1	F	79	UP	ESRD: ESRD V45.1-12/09/10

Figure 2-13: Sample Patient List, Diabetes: Nephropathy Assessment

2.2.7 Diabetic Retinopathy

GPRA Measure Description

During FY 2011, achieve the target rate of 50.1% for the proportion of patients with diagnosed diabetes who receive an annual retinal examination.

Denominators

All User Population patients diagnosed with diabetes prior to the Report Period.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, and at least two visits during the Report Period, and two DM-related visits ever. (GPRA Denominator)

Active Adult Diabetic patients, defined by meeting the following criteria: (1) who are 19 or older at the beginning of the Report Period, (2) whose first ever DM diagnosis occurred prior to the Report Period; (3) who had at least two DM related visits ever, (4) at least one encounter with DM POV in a primary clinic with a primary provider during the Report Period; and (5) never have had a creatinine value greater than five.

Numerators

Patients receiving a qualified retinal evaluation during the Report Period.

Note: This numerator does not include refusals. (GPRA Numerator)

- a. Patients receiving diabetic retinal exam during the Report Period
- b. Patients receiving other eye exams during the Report Period
- c. Patients who refused a diabetic retinal exam during the Report Period.

Logic Description

Diabetes definition: First DM Purpose of Visit 250.00–250.93 recorded in the V POV file prior to the Report Period.

Serum creatinine definition (used with Active Adult Diabetic denominator): Site-populated taxonomy DM AUDIT CREATININE TAX; or LOINC taxonomy (NOTE: CPT codes are not included since they do not store the result, which is used in this topic.).

Qualified retinal evaluation definition: (1) Diabetic retinal exam or (2) other eye exam, as shown below. The following methods are qualifying for this measure:

- Dilated retinal evaluation by an optometrist or ophthalmologist
- Seven standard fields stereoscopic photos (ETDRS) evaluated by an optometrist or ophthalmologist
- Any photographic method validated to seven standard fields (ETDRS).

CRS searches in the following order for:

Exam	CPT Codes	Other Codes
<i>Diabetic Retinal Exam (any of the following during the report period)</i>		
Diabetic Retinal Exam	2022F, 2024F, 2026F, S0620, S0621, S3000	VExam: 03 (dilated retinal examination or validated photographic equivalent)

Exam	CPT Codes	Other Codes
Other Eye Exam (any of the following during the report period)		
Non-Did Not Keep Appointment (DNKA) visit to ophthalmology or optometry or validated tele-ophthalmology retinal evaluation clinics		Clinic codes: A2, 17, 18, 64
Non-DNKA visit to an optometrist or ophthalmologist	67028, 67038 (old code), 67039, 67040, 92002, 92004, 92012, 92014	Provider Codes: 24, 79, 08 POV Code: V72.0 Procedure: 95.02
Refusal of a diabetic retinal exam. Refusals are only counted if the patient did not have a diabetic retinal exam or other eye exam. If a patient had both a diabetic retinal exam/other eye exam and a refusal, only the diabetic retinal exam/other eye exam will be counted.		Refusals Exam: 03

Key Logic Changes from CRS Version 11.0

1. For Other Eye Exam, noted that CPT code 67038 is an old code.

Patient List Description

List of diabetic patients with qualified retinal evaluation or refusal, if any.

Measure Source

HP 2020 D-10

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	53.0% (National rate)
IHS FY 2009 Performance	51.0% (National rate)
IHS FY 2008 Performance	50.0% (National rate)
IHS FY 2007 Performance	49.0% (National rate)
IHS FY 2006 Performance	49.0% (National Rate) 52.0% (Designated Sites Rate)
IHS FY 2005 Performance	50.0% (National Rate) 50.0% (Designated Sites Rate)
IHS FY 2004 Performance	47.0% (National Rate) 55.0% (Designated Sites Rate)
IHS FY 2003 Performance	49.0%
IHS FY 2002 Performance	49.0%
<i>HP 2020 Goal</i>	<i>58.7%</i>

DU	May 25, 2011						Page 19		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Diabetic Retinopathy									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
User Pop w/ DM DX prior to report period	229		202			180			
# w/Retinal Evaluation									
-No Refusals	65	28.4	47	23.3	+5.1	54	30.0	-1.6	
A. # w/ DM Retinal exam	7	3.1	6	3.0	+0.1	6	3.3	-0.3	
B. # w/Other Eye Exams	58	25.3	41	20.3	+5.0	48	26.7	-1.3	
# w/Retinal Exam Refusal	3	1.3	0	0.0	+1.3	0	0.0	+1.3	
Active Diabetic Pts (GPRA)	117		95			87			
# w/Retinal Evaluation									
-No Refusals (GPRA)	54	44.6	39	41.1	+3.6	44	50.6	-5.9	
A. # w/ DM Retinal exam	7	5.8	6	6.3	-0.5	6	6.9	-1.1	
B. # w/Other Eye Exams	47	38.8	33	34.7	+4.1	38	43.7	-4.8	
# w/Retinal Exam Refusal	3	2.5	0	0.0	+2.5	0	0.0	+2.5	
Active Diabetic Pts (GPRA)	144		97			87			
# w/Retinal Evaluation									
-No Refusals (GPRA)	54	37.5	39	40.2	-2.7	44	50.6	-13.1	
A. # w/ DM Retinal exam	7	4.9	6	6.2	-1.3	6	6.9	-2.0	
B. # w/Other Eye Exams	47	32.6	33	34.0	-1.4	38	43.7	-11.0	
# w/Retinal Exam Refusal	3	2.1	0	0.0	+2.1	0	0.0	+2.1	
Active Adult Diabetic Patients	102		75			63			
# w/Retinal Evaluation									
-No Refusals	39	38.2	32	42.7	-4.4	39	61.9	-23.7	
A. # w/ DM Retinal exam	6	5.9	4	5.3	+0.5	6	9.5	-3.6	
B. # w/Other Eye Exams	33	32.4	28	37.3	-5.0	33	52.4	-20.0	

# w/Retinal Exam Refusal	2	2.0	0	0.0	+2.0	0	0.0	+2.0
-----------------------------	---	-----	---	-----	------	---	-----	------

Figure 2-14: Sample Report, Diabetic Retinopathy

***** CONFIDENTIAL PATIENT INFORMATION, COVERED BY THE PRIVACY ACT *****

DU

May 25, 2011

Page 50

*** IHS 2011 Clinical Performance Measure Patient List ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Entire Patient List

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Diabetic Retinopathy: List of diabetic patients with qualified retinal
evaluation or refusal, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DEBORAH	000001	COMMUNITY #1	F	45	UP,AD,AAD	01/07/11 Cl: 18
PATIENT2,TARA	000002	COMMUNITY #1	F	51	UP,AD,AAD	
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	UP,AD,AAD	06/30/11 Cl: 18
PATIENT4,WINONA	000004	COMMUNITY #1	F	53	UP	
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	UP,AD,AAD	05/22/11 Refused
PATIENT6,RUTH	000006	COMMUNITY #1	F	64	UP	
PATIENT7,JONELLE	000007	COMMUNITY #1	F	69	UP,AD,AAD	03/29/11 Diab Eye

Ex

Figure 2-15: Sample Patient List, Diabetic Retinopathy

2.2.8 ACEI/ARB Use in Diabetic Patients

Denominator

Active Diabetic patients with HTN, defined as all Active Clinical patients diagnosed with diabetes and hypertension prior to the Report Period, AND at least 2 visits during the Report Period, AND 2 DM-related visits ever.

Numerators

Patients not receiving an ACEI or ARB medication during the Report Period.

- a. Patients with contraindication/previous adverse reaction to ACEI/ARB therapy.

Logic Description

Diabetes definition: First Purpose of Visit 250.00–250.93 recorded in the V POV file prior to the Report Period.

Hypertension definition: Diagnosis (POV or problem list) 401.* prior to the Report Period, and at least one hypertension POV during the Report Period.

ACEI/ARB Numerator Logic

Ace Inhibitor (ACEI) and ARB (Angiotensin Receptor Blocker) medication codes defined with medication taxonomy BGP PQA ACEI ARB MEDS.

ACEI medications are: Angiotensin Converting Enzyme Inhibitors (Benazepril, Captopril, Enalapril, Fosinopril, Lisinopril, Moexipril, Perindopril, Quinapril, Ramipril, Trandolopril).

Antihypertensive Combinations (Amlodipine-benazepril, Benazepril-hydrochlorothiazide, Captopril-hydrochlorothiazide, Enalapril-hydrochlorothiazide, Enalapril-Felodipine, Fosinopril-hydrochlorothiazide, Lisinopril-hydrochlorothiazide, Moexipril-hydrochlorothiazide, Quinapril-hydrochlorothiazide, Trandolapril-verapamil).

ARB medications: Angiotensin II Inhibitors (Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan).

Antihypertensive Combinations (Aliskiren-valsartan, Amlodipine-valsartan, Amlodipine-hydrochlorothiazide-valsartan, Amlodipine-olmesartan, Candesartan-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Irbesartan-hydrochlorothiazide, Losartan-hydrochlorothiazide, Olmesartan-amlodipine-hydrochlorothiazide, Olmesartan-hydrochlorothiazide, Telmisartan-amlodipine, Telmisartan-hydrochlorothiazide, Valsartan-hydrochlorothiazide).

CRS uses the following codes to define contraindications to ACE inhibitors/ARBs.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0-V23.9, V72.42, 640.*-649.*, or 651.*-676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI (not medically indicated) refusal for any ACE inhibitor/ARB at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to ACE inhibitors/ARBs.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ACE Inhibitors/ARBs (any of the codes occurring ever)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): “ace inhibitor”, “ACEI”, “Angiotensin Receptor Blocker” or “ARB”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “ace i*”, “ACEI”, “Angiotensin Receptor Blocker” or “ARB”

Key Logic Changes from CRS Version 11.0

New topic.

Patient List Description

List of diabetic patients with hypertension, with ACEI/ARB medication, contraindication, or ADR, if any.

Measure Source

PQA (Pharmacy Quality Alliance)

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 24		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

ACEI/ARB Use in Diabetic Patients									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Diabetic Pts									
w/ HTN	76		57			54			
# w/ no ACEI/ARB									
Rx or w/contra/									
ADR	21	27.6	9	15.8	+11.8	6	11.1	+16.5	
A. # w/contra/ADR									
w/ % of Total	13	61.9	2	22.2	+39.7	2	33.3	+28.6	

Figure 2-16: Sample Report, ACEI/ARB Use in Diabetic Patients

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

ACEI/ARB Use in Diabetic Patients: List of diabetic patients with hypertension, with ACEI/ARB medication, contraindication, or ADR, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,DEBORAH pregnant	000001	COMMUNITY #1	F	20	AD	ACEI/ARB contra
PATIENT2,TARA	000002	COMMUNITY #1	F	44	AD	06/05/11
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	45	AD	01/29/11
PATIENT4,WINONA	000004	COMMUNITY #1	F	57	AD	ACEI/ARB Allergy:
05/05/11 ADR POV 995.2 + E942.6						
PATIENT5,NADINE	000005	COMMUNITY #1	F	57	AD	
PATIENT6,RUTH	000006	COMMUNITY #1	F	61	AD	09/22/10
PATIENT7,JONELLE	000007	COMMUNITY #1	M	25	AD	06/01/11 ACEI/ARB
Contra POV 425.1						

Figure 2-17: Sample Patient List: ACEI/ARB Use in Diabetic Patients

2.2.9 Diabetes: Access to Dental Services

Denominator

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever.

Numerators

Patients with documented dental visit during the Report Period.

Note: This numerator does *not* include refusals.

Patients with documented dental exam refusal during the Report Period.

Logic Description

Diabetes definition: First DM Purpose of Visit 250.00–250.93 recorded in the V POV file prior to the Report period.

Dental visit definition: For non-CHS dental visits, searches for V Dental ADA codes 0000 or 0190; CPT Codes D0000 or D0190; VExam 30; or POV V72.2. For CHS dental visits, searches for any visit with an ADA code. CHS visit defined as Type code of C in Visit file.

Refusal definition: Non-CHS dental visit with refusal of ADA Code 0000 or 0190, refusal of CPT Code D0000 or D0190, or refusal of Exam 30. Refusals are only counted if the patient did not have a documented dental visit.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of diabetic patients with documented dental visit or refusal, if any.

Measure Source

HP 2020 D-8

Measure Past Performance and Long-Term Targets:

Past Performance	Percent
IHS FY 2005 Performance	39.0%
IHS FY 2004 Performance	37.0%
IHS FY 2003 Performance	36.0%
IHS FY 2002 Performance	36.0%

Past Performance	Percent
HP 2020 Goal	61.2%

Performance Improvement Tip

If your facility's dental services are paid for with CHS funds, ensure the final payment for each purchase order is posted and the CHS to PCC link is set to the "on" position. Having the CHS to PCC link on will enable the CHS data to be passed from CHS/MIS to PCC, where CRS can find it and include it in your CRS reporting.

DU	May 25, 2011						Page 26	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Diabetes: Access to Dental Services								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Diabetic Pts	144		97			87		
# w/dental visit in past yr-No Refusals	17	11.8	20	20.6	-8.8	18	20.7	-8.9
# w/dental exam refusal	4	2.8	0	0.0	+2.8	0	0.0	+2.8

Figure 2-18: Sample Report, Diabetes and Dental Access

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease								
Diabetes: Access to Dental Services: List of diabetic patients and documented dental visit or refusal, if any.								
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR		

PATIENT1,DEBORAH ADA 0000	000001	COMMUNITY #1	F	45	AD	03/03/11	Refused	
PATIENT2,TARA	000002	COMMUNITY #1	F	51	AD			
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	AD	01/06/11	ADA 0190	
PATIENT4,NADINE	000004	COMMUNITY #1	F	61	AD			
PATIENT5,SHERRY	000005	COMMUNITY #1	F	68	AD			
PATIENT6,JONELLE	000006	COMMUNITY #1	F	69	AD	03/29/11	ADA 0000	

Figure 2-19: Sample Patient List, Diabetes and Dental Access

2.3 Dental Measure Topics

2.3.1 Access to Dental Services

GPRA Measure Description

During FY 2011, achieve the target rate of 23.0% for the proportion of patients who receive dental services.

Denominators

All patients in the *User Population*. Broken down by age groups (0–5, 6–11, 12–19, 20–34, 35–44, 45–54, 55–74, >74). (GPRA Denominator)

Numerators

Patients with documented dental visit during the Report Period.

Note: This numerator does *not* include refusals. (GPRA Numerator)

Patients with documented dental exam refusal during the Report Period.

Logic Description

Dental visit definition: For non-CHS dental visits, searches for V Dental ADA codes 0000 or 0190; V CPT Codes D0000 or D0190; VExam 30; or POV V72.2. For CHS dental visits, searches for any visit with an ADA code. CHS visit defined as Type code of C in Visit file.

Refusal definition: Non-CHS dental visit with refusal of ADA Code 0000 or 0190, refusal of CPT Code D0000 or D0190, or refusal of Exam 30. Refusals are only counted if the patient did not have a documented dental visit.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with documented dental visit or refusal and date.

Measure Source

HP 2020 OH-7

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	25.0%

Performance	Percent
IHS FY 2009 Performance	25.0%
IHS FY 2008 Performance	25.0%
IHS FY 2007 Performance	25.0%
IHS FY 2006 Performance	23.0%
IHS FY 2005 Performance	24.0%
IHS FY 2004 Performance	24.0%
IHS FY 2003 Performance	25.0%
IHS FY 2002 Performance	24.9%
<i>HP 2020 Goal</i>	<i>49.0%</i>

Performance Improvement Tip

If your facility's dental services are paid for with CHS funds, ensure the final payment for each purchase order is posted and the CHS to PCC link is set to the "on" position. Having the CHS to PCC link on will enable the CHS data to be passed from CHS/MIS to PCC, where CRS can find it and include it in your CRS reporting.

DU	May 25, 2011						Page 28	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Access to Dental Services								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
# User Pop (GPRA)	2,896		2,456			2,346		
# w/dental visit in past yr-No Refusals (GPRA)	252	8.7	201	8.2	+0.5	207	8.8	-0.1
# w/dental exam refusal	6	0.2	0	0.0	+0.2	0	0.0	+0.2

Figure 2-20: Sample Report, Access to Dental Services

DU	May 25, 2011	Page 29
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Access to Dental Services (con't)		
TOTAL USER POPULATION		
Age Distribution		

	0-5	6-11	12-19	20-34	35-44	45-54	55-74	>74 yrs
CURRENT REPORT PERIOD								
Total # User Pop	347	255	367	664	389	390	403	81
# w/dental visit in past yr-No Refusals (GPRA)	25	29	33	72	35	32	25	1
% w/dental visit in past yr-No Refusals (GPRA)	7.2	11.4	9.0	10.8	9.0	8.2	6.2	1.2
# w/dental exam refusal	0	0	1	0	0	2	3	0
% w/dental exam refusal	0.0	0.0	0.3	0.0	0.0	0.5	0.7	0.0
PREVIOUS YEAR PERIOD								
Total # User Pop	352	237	349	595	316	277	275	55
# w/dental visit in past yr-No Refusals (GPRA)	19	22	30	53	24	24	25	4
% w/dental visit in past yr-No Refusals (GPRA)	5.4	9.3	8.6	8.9	7.6	8.7	9.1	7.3
# w/dental exam refusal	0	0	0	0	0	0	0	0
% w/dental exam refusal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CHANGE FROM PREV YR %								
w/dental visit in past yr-No Refusals (GPRA)	+1.8	+2.1	+0.4	+1.9	+1.4	-0.5	-2.9	-6.0
w/dental exam refusal	+0.0	+0.0	+0.3	+0.0	+0.0	+0.5	+0.7	+0.0
BASELINE REPORT PERIOD								
Total # User Pop	363	285	347	546	293	228	232	52
# w/dental visit in past yr-No Refusals (GPRA)	17	30	29	50	31	27	20	3
% w/dental visit in past yr-No Refusals (GPRA)	4.7	10.5	8.4	9.2	10.6	11.8	8.6	5.8
# w/dental exam refusal	0	0	0	0	0	0	0	0
% w/dental exam refusal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CHANGE FROM BASE YR %								
w/dental visit in past yr-No Refusals (GPRA)	+2.5	+0.8	+0.6	+1.7	-1.6	-3.6	-2.4	-4.5
w/dental exam refusal	+0.0	+0.0	+0.3	+0.0	+0.0	+0.5	+0.7	+0.0

Figure 2-21: Sample Age Breakdown Report, Access to Dental Services

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Access to Dental Service: List of patients with documented dental visit or refusal and date.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT10, JOHN	000010	COMMUNITY #1	M	17	UP	01/03/11	ADA 0190
PATIENT11, HOWARD	000011	COMMUNITY #1	M	25	UP	01/24/11	ADA 0000
PATIENT12, JAMES	000012	COMMUNITY #1	M	31	UP	02/19/11	ADA 0000
PATIENT13, STEVEN	000013	COMMUNITY #1	M	32	UP	01/24/11	ADA 0000
PATIENT14, EDWARD	000014	COMMUNITY #1	M	32	UP	06/10/11	ADA 0000
PATIENT15, DAVID	000015	COMMUNITY #1	M	33	UP	04/10/11	ADA 0190

Figure 2-22: Sample Patient List, Access to Dental Services

2.3.2 Dental Sealants

GPRA Measure Description

During FY 2011, achieve the target count of 257,261 sealants placed in American Indian and Alaska Native patients.

Denominator

No denominator. This measure is a total count only, not a percentage.

Numerators

For patients meeting the *User Population* definition, the total number of dental sealants during the report period.

Note: This numerator does not include refusals. (GPRA Numerator).

- Dental sealants in patients <12 yrs.
- Dental sealants in patients 12–18 yrs.
- Dental sealants in patients >18 yrs.

Age breakouts are based on Healthy People 2010 age groups for dental sealants.

Logic Description

Age of the patient is calculated at the beginning of the report period.

Sealants definition: V Dental ADA Code 1351 or V CPT Code D1351. Only two sealants per tooth will be counted during the Report Period. Each tooth is identified by the data element Operative Site in RPMS. If both ADA and CPT codes are found on the same visit, only the ADA will be counted.

Refusal definition: Refusal of ADA Code 1351 or refusal of CPT Code D1351. Refusals are only counted if a patient did not have a sealant during the Report Period. If a patient had both a sealant and a refusal, only the sealant will be counted. If a patient has multiple refusals, only one refusal will be counted.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients who received or refused dental sealants during Report Period.

Measure Source

HP 2010 21–8

Measure Past Performance and Long-Term Targets:

Performance	# of Sealants
IHS FY 2010 Performance	275,459
IHS FY 2009 Performance	257,067
IHS FY 2008 Performance	241,207
IHS FY 2007 Performance	245,449
IHS FY 2006 Performance	246,645
IHS FY 2005 Performance	249,882
IHS FY 2004 Performance	230,295 287,158
IHS FY 2003 Performance	232,182
IHS FY 2002 Performance	227,945
IHS FY 2001 Performance	212,612

For the IHS FY 2004 Performance, # of Sealants, please note this was reported by the National Patient Information Reporting System (NPIRS).

Performance Improvement Tip

If your facility's dental visits are paid for with CHS funds, ensure the final payment for each purchase order is posted and the CHS to PCC link is set to the "on" position. Having the CHS to PCC link on will enable the CHS data to be passed from CHS/MIS to PCC, where CRS can find it and include it in your CRS reporting.

DU	May 25, 2011	Page 30
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Dental Sealants					
	REPORT PERIOD	PREV YR PERIOD	CHG from PREV YR	BASE PERIOD	CHG from BASE
Total # of Sealants Documented (GPRA)	50	61	-11	81	-31
A. # Dental Sealants documented pts <12 yrs	35	26	+9	40	-5
B. # Dental Sealants documented pts 12-18 yrs	13	34	-21	40	-27
C. # Dental Sealants documented pts >18 yrs	2	1	+1	1	+1
# refusals	3	0	+3	0	+3

Figure 2-23: Sample Report, Dental Sealants

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Dental Sealants: List of patients who received or refused dental sealants during Report period.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT20,GEORGE	000020	COMMUNITY #1	M	5	UP	4 sealants:
03/28/11 ADA 1351 (1); 03/28/11 ADA 1351 (1); 03/28/11 ADA 1351 (1); 03/28/11 ADA 1351 (1)						
PATIENT21,CODY	000021	COMMUNITY #1	M	7	UP	0 sealants:
03/03/11 Refused ADA 1351						
PATIENT50,DAWN	000050	COMMUNITY #2	F	4	UP	3 sealants:
04/15/11 ADA 1351 (1); 05/19/11 ADA 1351 (1); 05/19/11 ADA 1351 (1)						
PATIENT51,JOY	000051	COMMUNITY #2	F	6	UP	2 sealants:
03/17/11 ADA 1351 (2)						
PATIENT52,DONALD	000052	COMMUNITY #2	M	8	UP	1 sealants:
02/02/11 CPT D1351 (1)						

Figure 2-24: Sample Patient List, Dental Sealants

2.3.3 Topical Fluoride

GPRA Measure Description

During FY 2011, achieve the target count of 135,604 AI/AN patients who receive at least one topical fluoride application.

Denominator

No denominator. This measure is a total count only, not a percentage.

Numerators

For patients meeting the *User Population* definition, the total number of patients with at least one topical fluoride treatment during the Report Period.

Note: This numerator does *not* include refusals. (GPRA Numerator)

For patients meeting the *User Population* definition, the total number of patients with a documented topical fluoride application refusal in past year.

For patients meeting the *User Population* definition, the total number of appropriate topical fluoride applications based on a maximum of four per patient per year.

For patients meeting the *User Population* definition, the total number of documented topical fluoride application refusals during past year.

Logic Description

Topical fluoride application definition: (1) V Dental ADA Codes 1201 (old code), 1203, 1204, 1205 (old code), 1206, or 5986; (2) V CPT Codes D1201 (old code), D1203, D1204, D1205 (old code), D1206, or D5986; 3) V POV V07.31. A maximum of one application per patient per visit is allowed. A maximum of four topical fluoride applications are allowed per patient per year for the applications measure.

Refusal definition: Refusal of ADA Code 1201 (old code), 1203, 1204, 1205 (old code), 1206 or 5986, or refusal of CPT Code D1201 (old code), D1203, D1204, D1205 (old code), D1206, or D5986. Refusals are only counted if a patient did not have a topical fluoride application during the Report Period. If a patient had both an application and a refusal, only the application will be counted. If a patient has multiple refusals, only one refusal will be counted.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients who received or refused at least one topical fluoride application during Report Period.

Measure Source

Not Available

Measure Past Performance and Long-Term Targets:

Performance	Number of Patients
IHS FY 2010 Performance	145,181
IHS FY 2009 Performance	136,794
IHS FY 2008 Performance	120,754
IHS FY 2007 Performance	107,934
IHS FY 2006 Performance	95,439
IHS FY 2005 Performance	85,318
IHS FY 2005 Performance	113,324

For the IHS FY 2005 Performance, Number of Patients (113,324) is the number of applications.

Performance Improvement Tip

If your facility's dental visits are paid for with CHS funds, ensure the final payment for each purchase order is posted and the CHS to PCC link is set to the "on" position. Having the CHS to PCC link on will enable the CHS data to be passed from CHS/MIS to PCC, where CRS can find it and include it in your CRS reporting.

DU	May 25, 2011				Page 33
*** IHS 2011 Selected Measures with Community Specified Report ***					
DEMO INDIAN HOSPITAL					
Report Period: Jan 01, 2011 to Dec 31, 2011					
Previous Year Period: Jan 01, 2010 to Dec 31, 2010					
Baseline Period: Jan 01, 2000 to Dec 31, 2000					

Topical Fluoride					
	REPORT PERIOD	PREV YR PERIOD	CHG from PREV YR	BASE PERIOD	CHG from BASE
Total # of patients w/At Least 1 Topical Fluoride App					
-No Refusals (GPRA)	45	26	+19	15	+30
# Patients					
w/Refusals	7	0	+7	0	+7
Total # of Topical Fluoride Applications					
	50	26	+24	15	+35
# Refusals					
	7	0	+7	0	+7

Figure 2-25: Sample Report, Topical Fluoride

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Topical Fluoride: List of patients who received or refused at least one topical

fluoride application during Report period.						
Topical Fluoride: List of patients who received or refused at least one topical fluoride application during Report period.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT20,GEORGE 06/18/11 CPT D5986	000020	COMMUNITY #1	M	5	UP	1 topical fluoride:
PATIENT21,RYAN 03/03/11 Refused ADA 1201	000021	COMMUNITY #1	M	8	UP	0 topical fluoride:
PATIENT22,MICHAEL 03/03/11 Refused CPT D1203	000022	COMMUNITY #1	M	9	UP	0 topical fluoride:
PATIENT23,MARTY 01/07/11 ADA 1204; 08/27/11 ADA 1204	000023	COMMUNITY #1	M	15	UP	2 topical fluoride:

Figure 2-26: Sample Patient List, Topical Fluoride

2.4 Immunization Measure Topics

2.4.1 Influenza

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 58.5% for the proportion of non-institutionalized adults aged 65 years and older who receive an influenza immunization.

Denominators

All Active Clinical patients. Broken down by age groups.

- Active Clinical patients *younger than age 18*.
- Active Clinical patients *ages 18-49*.
- Active Clinical patients *ages 18-49 and considered high risk for influenza*.
- Active Clinical patients *ages 50-64*.
- Active Clinical patients *ages 65 and older*. (GPRA Denominator)

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever.

All User Population patients. Broken down by age groups.

- User Population patients *younger than age 18*.
- User Population patients *ages 18-49*.
- User Population patients *ages 18-49 and considered high risk for influenza*.
- User Population patients *ages 50-64*.

- e. User Population patients ages 65 and older.

Numerators

Patients with influenza vaccine documented during the report period or with a contraindication documented at any time before the end of the report period.

Note: The only refusals included in this numerator are not medically indicated (NMI) refusals. (GPRA Numerator)

- a. Patients with a contraindication or a documented NMI refusal.

Patients with documented influenza refusal during the report period.

Logic Description

Age of the patient is calculated at the beginning of the report period.

Diabetes: First DM POV 250.00–250.93 recorded in the V POV file prior to the Report period.

Influenza definition: Any of the following documented during the Report Period unless otherwise noted.

1. **Influenza immunization:** Any of the codes in the table below.

	CPT Codes	ICD and Other Codes
Influenza vaccine	90654-90662, 90724 (old code), G0008, G8108 (old code)	Immunization (CVX) Codes: 88-Influenza Virus Vaccine, NOS; 15 Inf Virus Vac SV; 16 Inf Virus Vac WV; 111 Inf Virus Vac Intranasal; 135 Inf High Dose Seasonal; 140 Inf Virus Vac SV Preservative Free; 141 Inf Virus Vac SV POV: V04.8 (old code), V04.81 NOT documented with 90663, 90664, 90666-90668, 90470, G9141 or G9142, V06.6 NOT documented with 90663, 90664, 90666-90668, 90470, G9141 or G9142 ICD Procedure Code: 99.52

2. **Contraindication:** Any of the following documented at any time before the end of the Report Period, defined as: (A) Contraindication in the Immunization Package of "Egg Allergy" or "Anaphylaxis" or (B) PCC NMI Refusal.

3. **Refusal:** Refusal of Immunization Codes 88, 111, 135, 140, 141, 15, or 16, as documented in PCC Refusal File (i.e., REF) or in the Immunization Package as contraindication of "Patient Refusal."
4. **High Risk for Influenza:** Persons considered high risk for influenza are defined as those who have 2 or more visits in the past 3 years with a POV or Problem diagnosis of any of the following: HIV Infection (042, 042.0-044.9 (old codes)); Diabetes (250.00-250.93); Rheumatic Heart Disease (393.-398.99); Hypertensive Heart Disease (402.00-402.91); Hypertensive Heart/Renal Disease (404.00-404.93); Ischemic Heart Disease (410.00-414.9); Pulmonary Heart Disease (415.0-416.9); Other Endocardial Heart Disease (424.0-424.9); Cardiomyopathy (425.0-425.9); Congestive Heart Failure (428.0-428.9, 429.2); Chronic Bronchitis (491.0-491.9); Emphysema (492.0-492.8); Asthma (493.00-493.91); Bronchiectasis, CLD, COPD (494.0-496.); Pneumoconioses (500-505); Chronic Liver Disease (571.0-571.9); Nephrotic Syndrome (581.0-581.9); Renal Failure (585.6, 585.9); Transplant (996.80-996.89); Kidney Transplant (V42.0-V42.89); Chemotherapy (V58.1); Chemotherapy follow-up (V67.2).

Key Logic Changes from CRS Version 11.0

1. Changed the topic name from "Adult Immunizations: Influenza" to "Influenza".
2. Added new age group denominators.
3. Added logic for patients with high risk for Influenza.
4. Added ICD Procedure code 99.52 to Influenza Vaccine definition.
5. For Influenza definition, noted that HCPCS code G8108 is an old code.

Patient List Description

List of patients with Influenza code or refusal, if any.

Measure Source

HP 2020 IID-12.7

Measure Past Performance and Long-Term Targets for Patients => 65 Vaccine Rate:

Performance	Percent
IHS FY 2010 Performance	62.0%
IHS FY 2009 Performance	59.0%
IHS FY 2008 Performance	62.0%
IHS FY 2007 Performance	59.0%
IHS FY 2006 Performance	58.0%

Performance	Percent
IHS FY 2005 Performance	59.0%
IHS FY 2004 Performance	54.0%
IHS FY 2003 Performance	51.0%
IHS FY 2002 Performance	51.4%
<i>HP 2020 Goal</i>	<i>90.0%</i>

Performance Improvement Tips

1. Providers should ask about and record off-site historical immunizations (IZ type, date received and location) on PCC forms. Data entry mnemonic: HIM
2. Providers should document refusals; write "Refused" in Influenza Order box on PCC form. Data entry mnemonic: REF (Immunization, Value, Date Refused).

DU	May 25, 2011						Page 35		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Influenza									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical Pts	1,554		1,209			1,101			
Total # w/Flu vaccine/ contra/NMI Refusal	194	12.5	115	9.5	+3.0	49	4.5	+8.0	
A. # w/ Contraind/ NMI Ref									
w/ % of Total IZ	12	6.2	2	1.7	+4.4	0	0.0	+6.2	
# w/ Influenza Refusal	17	1.1	6	0.5	+0.6	0	0.0	+1.1	
Active Clinical Pts <18	442		416			435			
Total # w/Flu vaccine/ contra/NMI Refusal	54	12.2	21	5.0	+7.2	6	1.4	+10.8	
A. # w/ Contraind/ NMI Ref									
w/ % of Total IZ	3	5.6	0	0.0	+5.6	0	0.0	+5.6	
# w/ Influenza Refusal	6	1.4	0	0.0	+1.4	0	0.0	+1.4	
Active Clinical Pts 18-49	752		572			486			
Total # w/Flu vaccine/ contra/NMI Refusal	48	6.4	33	5.8	+0.6	14	2.9	+3.5	
A. # w/ Contraind/ NMI Ref									
w/ % of Total IZ	3	6.3	2	6.1	+0.2	0	0.0	+6.3	
# w/ Influenza									

Refusal	3	0.4	0	0.0	+0.4	0	0.0	+0.4
Active Clinical Pts 18-49 high risk	166		109			69		
Total # w/Flu vaccine/ contra/NMI Refusal	25	15.1	25	22.9	-7.9	7	10.1	+4.9
A. # w/ Contraind/ NMI Ref								
w/ % of Total IZ	3	12.0	1	4.0	+8.0	0	0.0	+12.0
# w/ Influenza Refusal	3	1.8	0	0.0	+1.8	0	0.0	+1.8
Active Clinical Patients ages 50-64	244		157			115		
Total # w/Flu vaccine/contra/ NMI Refusal	59	24.2	37	23.6	+0.6	14	12.2	+12.0
A. # w/ Contraind/ NMI Ref w/ % of Total IZ	5	8.5	0	0.0	+8.5	0	0.0	+8.5
# w/Influenza Refusal	3	1.2	5	3.2	-2.0	0	0.0	+1.2

Figure 2-27: Sample Report, Adult Immunizations: Influenza

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Influenza: List of patients with Influenza code or refusal, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,DEBORAH	000001	COMMUNITY #1	F	15	AD	01/28/11 Imm 88	
PATIENT2,CRYSTAL	000002	COMMUNITY #1	F	24	UP,AC		
PATIENT3,DEMETRIA	000003	COMMUNITY #1	F	35	UP,AC, HR	02/25/11 Imm	
PATIENT4,JADE	000004	COMMUNITY #1	F	50	UP		
PATIENT5,MARIE	000005	COMMUNITY #1	F	65	UP,AC,AD,HR	01/21/11 NMI	
Refusal							

Figure 2-28: Sample Patient List, Adult Immunization: Influenza

2.4.2 Adult Immunizations

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 79.3% for the proportion of adult patients age 65 years and older who receive a pneumococcal immunization.

Denominators

Active Clinical patients ages 65 or older. (GPRA Denominator)

Active Clinical patients ages 18-64 and considered high risk for pneumococcal.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever.

Active Clinical patients ages 18-64.

All User Population patients ages 65 and older at beginning of Report Period.

User Population patients ages 18-64 and considered high risk for pneumococcal.

User Population patients ages 18-64.

Numerators

Patients with Pneumococcal vaccine or contraindication documented at any time before the end of the Report Period.

Note: The only refusals included in this numerator are NMI refusals. (GPRA Numerator)

- a. Patients with a contraindication or a documented NMI refusal.

Patients with documented Pneumococcal refusal during the Report Period.

Patients with Pneumococcal vaccine or contraindication documented ever and, if patient is older than 65 years, either a dose of pneumovax after the age of 65 or a dose of pneumovax in the past 5 years.

Note: The only refusals included in this numerator are NMI refusals. (GPRA Numerator)

Patients who have received 1 dose of Tdap ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals. (GPRA Numerator)

Patients who have received 1 dose of Tdap/Td in the past 10 years, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals. (GPRA Numerator)

Logic Description

Age of the patient is calculated at the beginning of the Report period.

Diabetes definition: First DM POV 250.00250.93 recorded in the V POV file prior to the Report period.

Pneumococcal Immunization definition: Any of the following documented anytime before the end of the Report Period unless otherwise noted.

1. **Pneumococcal immunization:** Any of the codes in the table below.

Immunization	CPT Codes	ICD and Other Codes
Pneumococcal Vaccine	90669, 90670, 90732, G0009, G8115 (old code)	Immunization (CVX) Codes: 33 – Pneumococcal Polysaccharide Vaccine; 100 – Pneumococcal Conjugate Vaccine; 109 – Pneumo NOS; 133 – Pneumo Conjugate POV: V06.6; V03.82 V Procedure: 99.55

2. **Pneumococcal Contraindication:** (A) Contraindication in the Immunization Package of "Anaphylaxis" or (B) PCC NMI Refusal.
3. **Pneumococcal Refusal:** Any of the following during the Report Period: (A) Immunization codes 33, 100, 109, or 133, as documented in PCC Refusal File (i.e., REF) or (B) Immunization Package as contraindication of "Patient Refusal."
4. **High Risk for Pneumococcal:** Persons considered high risk for pneumococcal are defined as those who have 2 or more visits in the past 3 years with a POV or Problem diagnosis of any of the following: HIV Infection (042, 042.0-043.9 (old codes), 044.9 (old code)); Diabetes (250.00-250.93); Chronic alcoholism (303.90, 303.91); Congestive Heart Failure (428.0-428.9, 429.2); Emphysema (492.0-492.8); Asthma (493.00-493.91); Bronchiectasis, CLD, COPD (494.-496.); Pneumoconioses (501.-505.); Chronic Liver Disease (571.0-571.9); Nephrotic Syndrome (581.0-581.9); Renal Failure (585.6, 585.9); Injury to spleen (865.00-865.19); Transplant (996.80-996.89); Kidney Transplant (V42.0-V42.89); Chemotherapy (V58.1); Chemotherapy follow-up (V67.2).

Tdap/Td Immunization definition: Any of the following documented during the applicable time frame.

1. **Tdap/Td immunization:** Any of the codes in the table below.

Immunization	CPT Codes	ICD and Other Codes
Tdap Vaccine	90715	Immunization (CVX) Codes: 115 – Tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccine, adsorbed

Td Vaccine	90714, 90718	Immunization (CVX) Codes: 9 – Tetanus and diphtheria toxoids, adsorbed, for adult use 113 – Tetanus and diphtheria toxoids, adsorbed, preservative free, for adult use POV: V06.5
------------	--------------	--

2. **Tdap/Td Contraindication:** (A) Contraindication in the Immunization Package of "Anaphylaxis" or (B) PCC NMI Refusal.

Key Logic Changes from CRS Version 11.0

1. Changed the topic name from "Adult Immunizations: Pneumovax" to "Adult Immunizations".
2. Added new denominator and logic for patients with high risk for Pneumovax.
3. Added new measures and logic for Tdap/Td.
4. For Pneumovax definition, noted that HCPCS code G8115 is an old code.

Patient List Description

List of patients =>18 yrs or DM DX with IZ, evidence of disease, contraindication, or refusal, if any.

Measure Source

HP 2020 IID-13.1

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	84.0%
IHS FY 2009 Performance	82.0%
IHS FY 2008 Performance	82.0%
IHS FY 2007 Performance	79.0%
IHS FY 2006 Performance	74.0%
IHS FY 2005 Performance	69.0%
IHS FY 2004 Performance	69.0%
IHS FY 2003 Performance	65.0%
IHS FY 2002 Performance	64.0%
<i>HP 2020 Goal for % of patients => 65</i>	<i>90.0%</i>

Performance Improvement Tips

1. Providers should ask about and record off-site historical immunizations (IZ type, date received and location) on PCC forms. Data entry mnemonic: HIM

2. Providers should document refusals; write "Refused" in Pneumo Vax Order box on PCC form. Data entry mnemonic: REF (Immunization, Value, Date Refused).

DU	May 25, 2011						Page 41	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Adult Immunizations								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
ages 65 & older								
(GPRA)	116		64			65		
Total # w/Pneumovax/								
contra/NMI Refusal								
(GPRA)	50	43.1	44	68.8	-25.6	37	56.9	-13.8
A. # w/ Contraind/ NMI								
Ref w/ % of								
Total IZ	4	8.0	2	4.5	+3.5	0	0.0	+8.0
# w/Pneumovax								
Refusal	3	2.6	0	0.0	+2.6	0	0.0	+2.6
Active Clinical Pts								
18-64 high risk								
	222		164			105		
Total # w/Pneumovax/ contra/								
NMI Refusal								
	55	24.8	47	28.7	-3.9	39	37.1	-12.4
A. # w/ Contra/ NMI Ref								
w/ % of Total IZ								
	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ Pneumovax								
Refusal	0	0.0	0	0.0	+0.0	0	0.0	+0.0
Active Diabetic Pts								
	144		97			87		
Total # w/up to date								
Pneumovax/contra/NMI								
Refusal	56	38.9	49	50.5	-11.6	46	52.9	-14.0
A. # w/ Contraind/ NMI								
Ref w/ % of								
Total IZ	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/Pneumovax								
Refusal	0	0.0	0	0.0	+0.0	0	0.0	+0.0

Figure 2-29: Sample Report, Adult Immunization: Pneumovax

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic						
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Adult Immunizations: List of patients =>18 yrs or DM DX with IZ, evidence						
of disease, contraindication, or refusal, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,DEBORAH	000001	COMMUNITY #1	F	18	AC 18-64,AC 18-64HR,UP 18-64,UP 18-64HR ; TDAP/TD PAST 10 YRS: 12/22/01
PATIENT2,TARA	000002	COMMUNITY #1	F	27	UP 18-64 ; TDAP/TD PAST 10 YRS: 09/02/98
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	41	AC 18-64,UP 18-64 ; TDAP EVER: ; TDAP/TD PAST 10 YRS: 03/03/03
PATIENT4,NADINE	000004	COMMUNITY #1	F	55	AC 18-64,UP 18-64 ; TDAP EVER: 03/03/03; TDAP/TD PAST 10 YRS: 03/03/03
PATIENT5,SHERRY	000005	COMMUNITY #1	F	68	AC 18-64,AC 18-64HR,UP 18-64,UP 18-64HR,AD04/20/94 Imm 33 (ever) (up-to-date); TDAP/TD PAST 10 YRS: 04/20/94

Figure 2-30: Sample Patient List, Adult Immunization: Pneumovax

2.4.3 Childhood Immunizations

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 74.6% for the proportion of American Indian/Alaska Native children ages 19-35 months who have received the recommended immunizations.

Note: In FY 2011, the GPRA measure is changed to the 4:3:1:3:3:1:4 combination, which includes pneumococcal.

Denominators

Active Clinical patients ages 19–35 months at end of Report Period.

User Population patients ages 19–35 months at end of Report Period.

User Population patients *active in the Immunization Package* who are 19–35 months at end of Report Period. (GPRA Denominator)

Note: Only values for the Current Period will be reported for this denominator since currently there is not a way to determine if a patient was active in the Immunization Package during the Previous Year or Baseline Periods.

Numerators

Patients who have received the 4:3:1:3:3 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B), including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received the 4:3:1:3:3:1 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B, 1 Varicella), including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received the 4:3:1:3:3:1:4 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B, 1 Varicella, and 4 Pneumococcal), including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals. (GPRA Numerator)

Patients who have received four doses of DTaP ever, including contraindications.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received three doses of Polio ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received one dose of MMR ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received three doses of HiB ever, including contraindications.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received three doses of Hepatitis B vaccine ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received one dose of Varicella ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received four doses of Pneumococcal conjugate vaccine ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received two doses of Hepatitis A vaccine ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received two or three doses of Rotavirus vaccine ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received two doses of Influenza vaccine ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

For each of the above numerators, the following subnumerators are included:

- a. Patients with either (1) evidence of the disease, (2) a contraindication, or (3) a documented NMI refusal

Patients with documented REF refusal in PCC or Parent or Patient refusal in the IZ program

Immunization Program Numerator: Patients who have received the 4:3:1:3:3 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B), *not* including refusals, contraindications, and patients with evidence of disease.

Immunization Program Numerator: Patients who have received the 4:3:1:3:3:1 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B, and 1 Varicella), *not* including refusals, contraindications, and patients with evidence of disease.

Immunization Program Numerator: Patients who have received the 4:3:1:3:3:1:4 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 HiB, 3 Hepatitis B, 1 Varicella, and 4 Pneumococcal), *not* including refusals, contraindications, and patients with evidence of disease.

Logic Description

Age definition: Age of the patient is calculated at the beginning of the Report Period. Therefore the age range will be adjusted to 7–23 months, which makes the patient between the ages of 19–35 months at the end of the Report Period. Because IZ data comes from multiple sources, any IZ codes documented on dates within 10 days of each other will be considered as the same immunization.

Active Immunization Package Patients denominator definition: Same as User Pop definition except includes only patients flagged as active in the Immunization Package.

Note: Only values for the Current Period will be reported for this denominator since currently there is not a way to determine if a patient was active in the Immunization Package during the Previous Year or Baseline Periods.

Dosage and types of immunization definitions:

- Four doses of DTaP: (1) four DTaP/DTP/Tdap; (2) one DTaP/DTP/Tdap and three DT/td; (3) one DTaP/DTP/Tdap and three each of Diphtheria and Tetanus; (4) four DT and four Acellular Pertussis; (5) four Td and four Acellular Pertussis; or (6) four each of Diphtheria, Tetanus, and Acellular Pertussis.
- Three doses of Polio: (1) v OPV; (2) three IPV; or (3) combination of OPV and IPV totaling 3 doses.
- One dose of MMR: (1) MMR; (2) one M/R and one Mumps; (3) one R/M and one Measles; or (4) one each of Measles, Mumps, and Rubella.
- Three doses of Hep B *or* two doses *if* documented with CPT 90743.
- Three doses of HIB
- One dose of Varicella
- Four doses of Pneumococcal
- Two doses of Hepatitis A
- Two or three doses of Rotavirus, depending on the vaccine administered
- Two doses of Influenza

Except for the Immunization Program Numerators, NMI refusals, evidence of disease, and contraindications for individual immunizations will also count toward meeting the definition, as defined below. Refusals will count toward meeting the definition for refusal numerators only.

Note: NMI refusals are not counted as refusals; rather, they are counted as contraindications.

- For immunizations that allow a different number of doses (e.g. 2 or 3 Rotavirus): To count toward the numerator with the smaller number of doses, all of the patient's vaccinations must be part of the smaller dose series. For example, for a patient to count toward the Rotavirus numerator with only 2 doses, all two doses must be included in the 2-dose series codes listed in the Rotavirus definition. A patient with a mix of 2-dose and 3-dose series codes will need 3 doses to count toward the numerator.

- Each immunization must be refused and documented separately. For example, if a patient has an NMI refusal for Rubella only, then there must be an immunization, contraindication, or separate NMI refusal for the Measles and Mumps immunizations.
- For immunizations where required number of doses is >1, only one NMI refusal is necessary to be counted in the numerator. For example, if there is a single NMI refusal for Hepatitis B, the patient will be included in the numerator.
- For immunizations where required number of doses is >1, only one contraindication is necessary to be counted in the numerator. For example, if there is a single contraindication for HiB, the patient will be included in the numerator.
- Evidence of disease will be checked for at any time in the child's life (prior to the end of the Report Period.)
- To be counted in subnumerator A, a patient must meet the numerator definition *and* have a REF refusal in PCC or a Parent or Patient Refusal in the IZ program for any of the immunizations in the numerator. For example, if a patient refused Rubella only but had immunizations for Measles and Mumps, the patient would be included in subnumerator A.

For the separate numerator for REF refusal (Patient Refusal for Service) in PCC or a Parent or Patient refusal in the IZ program, all conditions shown below must be met:

- Each immunization must be refused and documented separately. For example, if a patient has an REF refusal for Rubella, then there also must be an immunization, contraindication, or separate REF refusal for Measles and Mumps.
- Where the required number of doses is >1, only one REF refusal in PCC or one Parent or Patient refusal in the IZ program is necessary to be counted in the numerator. For example, for the 4 DTaP numerator, only one refusal is necessary to be counted in the refusal numerator.

Childhood immunization refusals are defined as Parent/Patient Refusal in Immunization package or PCC Refusal type REF or NMI for IZ codes:

Immunization	Immunization Codes for Refusals
DTaP	20, 50, 106, 107, 110, 120, 130, 132
DTP	1, 22, 102
Tdap	115
DT (Diphtheria & Tetanus)	28
Td (Tetanus & Diphtheria)	9, 113
Tetanus	35, 112
Acellular Pertussis	11
OPV	2, 89
IPV	10, 89, 110, 120, 130, 132

Immunization	Immunization Codes for Refusals
MMR	3, 94
M/R (Measles/ Rubella)	4
R/M (Rubella/ Mumps)	38
Measles	5
Mumps	7
Rubella	6
HiB	17, 22, 46-49, 50, 51, 102, 120, 132
Hepatitis B	8, 42-45, 51, 102, 104, 110, 132
Varicella	21, 94
Pneumococcal	33, 100, 109, 133
Hepatitis A	1, 52, 83, 84, 85, 104
Rotavirus	74, 116, 119, 122
Influenza	15, 16, 88, 111, 135, 140, 141

Childhood immunizations are defined in the following ways:

Immunization	CPT Codes	ICD and Other Codes NOTE: IZ Program Numerators Do Not Use POV, V Procedure, Evidence of Disease, Contraindication or Refusal Codes
DTaP	90696, 90698, 90700, 90721, 90723	Immunization (CVX) Codes: 20, 50, 106, 107, 110, 120, 130, 132 POV: V06.1 Contraindications: Immunization Package contraindication of "Anaphylaxis."
DTP	90701, 90711 (old code), 90720	Immunization (CVX) Codes: 1, 22, 102 POV: V06.1, V06.2, V06.3 V Procedure: 99.39 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Tdap	90715	Immunization (CVX) Codes: 115 Contraindications: Immunization Package contraindication of "Anaphylaxis."
DT (Diphtheria & Tetanus)	90702	Immunization (CVX) Codes: 28 POV: V06.5 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Td (Tetanus & Diphtheria)	90714, 90718	Immunization (CVX) Codes: 9, 113 POV: V06.5 Contraindications: Immunization Package contraindication of "Anaphylaxis."

Immunization	CPT Codes	ICD and Other Codes NOTE: IZ Program Numerators Do Not Use POV, V Procedure, Evidence of Disease, Contraindication or Refusal Codes
Diphtheria	90719	POV: V03.5 V Procedure: 99.36 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Tetanus	90703	Immunization (CVX) Codes: 35, 112 POV: V03.7 V Procedure: 99.38 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Acellular Pertussis		Immunization (CVX) Codes: 11 POV: V03.6 V Procedure: 99.37 (old code) Contraindications: Immunization Package contraindication of "Anaphylaxis."
OPV	90712	Immunization (CVX) Codes: 2, 89 Contraindications: POV 279, V08, 042, 200-202, 203.0, 203.1, 203.8, 204-208 or Immunization Package contraindication of "Anaphylaxis."
IPV	90696, 90698, 90711 (old code), 90713, 90723	Immunization (CVX) Codes: 10, 89, 110, 120, 130, 132 POV: V04.0, V06.3 V Procedure: 99.41 Evidence of Disease: POV or PCC Problem List (active or inactive) 730.70-730.79 Contraindications: Immunization Package contraindication of "Anaphylaxis" or "Neomycin Allergy."
MMR	90707, 90710	Immunization (CVX) Codes: 3, 94 POV: V06.4 V Procedure: 99.48 Contraindications: POV 279, V08, 042, 200-202, 203.0, 203.1, 203.8, 204-208; or Immunization Package contraindication of "Anaphylaxis," "Immune Deficiency," "Immune Deficient," or "Neomycin Allergy."
M/R (Measles/Rubella)	90708	Immunization (CVX) Codes: 4 Contraindications: Immunization Package contraindication of "Anaphylaxis"
R/M (Rubella/Mumps)	90709 (old code)	Immunization (CVX) Codes: 38 Contraindications: Immunization Package contraindication of "Anaphylaxis"

Immunization	CPT Codes	ICD and Other Codes NOTE: IZ Program Numerators Do Not Use POV, V Procedure, Evidence of Disease, Contraindication or Refusal Codes
Measles	90705	Immunization (CVX) Codes: 5 POV: V04.2 V Procedure: 99.45 Evidence of Disease: POV or PCC Problem List (active or inactive) 055* Contraindications: Immunization Package contraindication of "Anaphylaxis"
Mumps	90704	Immunization (CVX) Codes: 7 POV: V04.6 V Procedure: 99.46 Evidence of Disease: POV or PCC Problem List (active or inactive) 072* Contraindications: Immunization Package contraindication of "Anaphylaxis"
Rubella	90706	Immunization (CVX) Codes: 6 POV: V04.3 V Procedure: 99.47 Evidence of Disease: POV or PCC Problem List (active or inactive) 056*, 771.0 Contraindications: Immunization Package contraindication of "Anaphylaxis"
HiB	90645-90648, 90698, 90720-90721, 90737 (old code), 90748	Immunization (CVX) Codes: 17, 22, 46-49, 50, 51, 102, 120, 132 POV: V03.81 Contraindications: Immunization Package contraindication of "Anaphylaxis"
Hepatitis B	90636, 90723, 90731 (old code), 90740, 90743-90748, G0010, Q3021, (old code) Q3023 (old code)	Immunization (CVX) Codes: 8, 42-45, 51, 102, 104, 110, 132 Evidence of Disease: POV or PCC Problem List (active or inactive) V02.61, 070.2, 070.3 Contraindications: Immunization Package contraindication of "Anaphylaxis"

Immunization	CPT Codes	ICD and Other Codes NOTE: IZ Program Numerators Do Not Use POV, V Procedure, Evidence of Disease, Contraindication or Refusal Codes
Varicella	90710, 90716	Immunization (CVX) Codes: 21, 94 POV: V05.4 Evidence of Disease: 1) POV or PCC Problem List (active or inactive) 052*, 053* or 2) 2) Immunization Package contraindication of "Hx of Chicken Pox" or "Immune." Contraindications: POV 279, V08, 042, 200-202, 203.0, 203.1, 203.8, 204-208; or Immunization Package contraindication of "Anaphylaxis," "Immune Deficiency," "Immune Deficient," or "Neomycin Allergy."
Pneumococcal	90669, 90670, 90732, G0009, G8115 (old code)	Immunization (CVX) Codes: 33, 100, 109, 133 POV: V06.6; V03.82 Contraindications: Immunization Package contraindication of "Anaphylaxis"
Hepatitis A	90632-90634, 90636, 90730 (old code)	Immunization (CVX) Codes: 31, 52, 83, 84, 85, 104 Evidence of Disease: POV or PCC Problem List (active or inactive) 070.0, 070.1 Contraindications: Immunization Package contraindication of "Anaphylaxis"
Rotavirus – 2-dose series	90681	Immunization (CVX) Codes: 119 Contraindications: Immunization Package contraindication of "Anaphylaxis" or "Immune Deficiency"
Rotavirus – 3-dose series	90680	Immunization (CVX) Codes: 74, 116, 122 POV: V05.8 Contraindications: Immunization Package contraindication of "Anaphylaxis" or "Immune Deficiency"
Influenza	90654-90658, 90659 (old code), 90660-90662, 90724 (old code), G0008, G8108 (old code)	Immunization (CVX) Codes: 15, 16, 88, 111, 135, 140, 141 POV: V04.8 (old code), V04.81, V06.6 V Procedure: 99.52 Contraindications: Immunization Package contraindication of "Egg Allergy" or "Anaphylaxis"

Key Logic Changes from CRS Version 11.0

1. Added measures and logic for Hepatitis A, Rotavirus, and Influenza
2. For Pneumococcal, noted that HCPCS code G8115 is an old code.

Patient List Description

List of patients 19–35 months with IZ, if any. If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had two DTaP, no IZ will be listed for DTaP.

Note 1: Because age is calculated at the beginning of the Report Period, the patient's age on the list will be between 7-23 months.

Note 2: The order of the display for the immunizations is: 4 Dtap/Dtp;3 IPV/OPV;MMR;3 HIB;3 HEP;Vari;4 PNEUMO. A blank value in the Numerator column means the patient didn't meet the requirements for any of the immunizations. Another example is "MMR; vari;4 PNEUMO," which means the patient did not have 4 Dtap/Dtp, 3 IPV/OPV, 3 HIB and 3 Hep B immunizations.

Measure Source

CDC; HP 2020 IID-7, IID-8; HEDIS

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 GPRA Performance Active Immunization Package 4:3:1:3:3:1 (rate for children age 19–35 months)	79.0%
IHS FY 2009 GPRA Performance Active Immunization Package 4:3:1:3:3 (rate for children age 19–35 months)	79.0%
IHS FY 2008 GPRA Performance Active Immunization Package 4:3:1:3:3 (rate for children age 19–35 months)	78.0%
<i>IHS FY 2008 Non-GPRA Performance Active Clinical 4:3:1:3:3 (rate for children age 19–35 months)</i>	68.0%
IHS FY 2007 GPRA Performance Active Immunization Package 4:3:1:3:3(rate for children age 19–35 months)	78.0%
IHS FY 2006 Performance (rate for children age 19–35 months)	80.0%
IHS FY 2005 Performance (rate for children age 19–35 months)	75.0%
IHS FY 2004 Performance(baseline rate for children age 19-35 months)	72.0%
IHS FY 2004 Performance(rate for children age 3-27 months)	81.0%
IHS FY 2003 Performance(rate for children age 3-27 months)	80.0%
IHS FY 2002 Performance(rate for children age 3-27 months)	80.0%

Performance	Percent
HP 2020 goal for % of children age 19–35 months with 4:3:1:3:3:4 vaccines	80.0%
HP 2020 goal for % of children age 19–35 months with each individual vaccine	90.0%

For the IHS FY 2006 Performance (rate for children 19-35 months), the Percent (80.0) please consider: All 2002–2006 rates reported on this table were reported by the Immunization Program from the quarterly immunization reports. Effective in 2007, CRS reports the rate and not the Immunization Program. The CRS rate is reported using the CRS Active Immunization Package denominator.

Performance Improvement Tips

1. Providers should ask about and record off-site historical immunizations (IZ type, date received and location) on PCC forms. Data entry mnemonic: HIM
2. Providers should document refusals; write “Refused” in appropriate vaccine order box on PCC form. Data entry mnemonic: REF (Immunization, Value, Date Refused).

DU

May 25, 2011

Page 51

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Childhood Immunizations

	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
Active Clinical Pts								
19-35 months	56		39			55		
# w/ 43133 combo								
or w/ Dx/ Contraind/								
NMI Refusal	11	19.6	3	7.7	+12.0	6	10.9	+8.7
A. # w/ Dx/Contraind/NMI								
Ref w/ % of								
Total 43133	1	9.1	0	0.0	+9.1	0	0.0	+9.1
# w/ 43133								
refusal	2	3.6	0	0.0	+3.6	0	0.0	+3.6
# w/ 431331 combo								
or w/ Dx/ Contraind/								
NMI Refusal	9	16.1	3	7.7	+8.4	5	9.1	+7.0
A. # w/ Dx/Contraind/NMI								
Ref w/ % of								
Total 431331	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ 431331								

Refusal	1	1.8	0	0.0	+1.8	0	0.0	+1.8
# w/ 4313314 combo or w/Dx/Contraind/ NMI Refusal	3	5.4	0	0.0	+5.4	0	0.0	+5.4
A. # w/ Dx/Contraind/NMI Ref w/ % of Total 4313314	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ 4313314 Refusal	2	3.6	0	0.0	+3.6	0	0.0	+3.6
# w/ 4 doses DTaP or w/ Contraind/ NMI Refusal	16	28.6	3	7.7	+20.9	9	16.4	+12.2
A. # w/ Contraind/NMI Ref w/ % of Total DTaP	1	6.3	0	0.0	+6.3	0	0.0	+6.3
# w/ DTaP Refusal	6	10.7	0	0.0	+10.7	0	0.0	+10.7

Figure 2-31: Sample Report, Childhood Immunizations

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Childhood Immunizations: List of patients 19-35 months with IZ, if any. If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had 2 DTaP, no IZ will be listed for DTaP. NOTE: Because age is calculated at the beginning of the Report Period, the patient's age on the list will be between 7-23 months.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,ANDREA	000001	COMMUNITY #1	F	0	UP;AC;IMM	4 Dtap/Dtp;3
OPV;MMR;3 HIB;3 Hep B;vari; 2 Influenza						
PATIENT2,HEATHER	000002	COMMUNITY #1	F	1	UP;AC	4 Dtap/Dtp;3
OPV;MMR;3 HIB;3 Hep B;vari;4 PNEUMO						
PATIENT3,TONYA	000003	COMMUNITY #1	F	1	UP	
PATIENT4,JAMES	000004	COMMUNITY #1	M	0	UP;AC;IMM	4 Dtap/Dtp;3
OPV;MMR;3 HIB;3 Hep B;vari						
PATIENT5,SCOTT	000005	COMMUNITY #1	M	0	UP;AC;IMM	3 HIB; 2 HEP A; 3
Rotavirus; 2 Influenza						

Figure 2-32: Sample Patient List, Childhood Immunizations

2.4.4 Adolescent Immunizations

Denominators

Active Clinical patients age 13.

Female Active Clinical patients age 13.

Active Clinical patients ages 13–17.

Female Active Clinical patients ages 13–17.

Numerators

Patients who have received the 2:3:1 combination (i.e., two MMR, three Hepatitis B, one Varicella), including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patient who have received the 1:3:2:1 combination (i.e., one Td/Tdap, three Hepatitis B, two MMR, one Varicella), including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received one dose of Tdap/Td ever, including contraindications and evidence of disease.

- a. Patients with (1) evidence of the disease, (2) a contraindication, or (3) a documented NMI refusal.
- b. Patients who have received one dose of Tdap ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received two doses of MMR ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received three doses of Hepatitis B ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received one dose of Varicella ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received one dose of meningococcal ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.

Patients who have received three doses of HPV ever, including contraindications and evidence of disease.

Note: The only refusals included in this numerator are NMI refusals.
Note: Included for Female Active Clinical age 13 and Female Active Clinical ages 13–17 only.

For each of the above numerators, the following subnumerators are included:

- a. Patients with (1) evidence of the disease, (2) a contraindication, or (3) a documented NMI refusal.

Patients with documented REF refusal in PCC or Parent or Patient refusal in the IZ program.

Logic Description

Age definition: Age of the patient is calculated at the beginning of the Report Period. Because IZ data comes from multiple sources, any IZ codes documented on dates within ten days of each other will be considered as the same immunization.

Dosage and types of immunization definitions:

- One dose of Td or Tdap
- Two doses of MMR: (1) two MMRs; (2) two M/R and two Mumps; (3) two R/M and two Measles; or (4) two each of Measles, Mumps, and Rubella
- Three doses of Hep B *or* two doses *if* documented with CPT 90743
- One dose of Varicella
- One dose of Meningococcal
- Three doses of HPV

Not Medically Indicated (NMI) refusals, evidence of disease and contraindications for individual immunizations will also count toward meeting the definition, as defined below. Refusals will count toward meeting the definition for refusal numerators only. NOTE: NMI refusals are not counted as refusals; rather, they are counted as contraindications.

- Each immunization must be refused and documented separately. For example, if a patient has an NMI refusal for Rubella only, then there must be an immunization, contraindication, or separate refusal for the Measles and Mumps immunizations.
- For immunizations where required number of doses is >1, only one NMI refusal is necessary to be counted in the numerator. For example, if there is a single NMI refusal for Hepatitis B, the patient will be included in the numerator.
- For immunizations where required number of doses is >1, only one contraindication is necessary to be counted in the numerator. For example, if there is a single contraindication for HiB, the patient will be included in the numerator.

- Evidence of disease will be checked for at any time in the child's life (prior to the end of the Report Period).
- To be counted in subnumerator A, a patient must have evidence of disease, a contraindication, or an NMI refusal for any of the immunizations in the numerator. For example, if a patient was Rubella immune but had a Measles and Mumps immunization, the patient would be included in subnumerator A.

For the separate numerator for REF refusal (Patient Refusal for Service) in PCC or a Parent or Patient refusal in the IZ program, all conditions shown below must be met:

- Each immunization must be refused and documented separately. For example, if a patient has an REF refusal for Rubella, then there also must be an immunization, contraindication, or separate REF refusal for Measles and Mumps.
- Where the required number of doses is >1, only one REF refusal in PCC or one Parent or Patient refusal in the IZ program is necessary to be counted in the numerator. For example, for the 4 DTap numerator, only one refusal is necessary to be counted in the refusal numerator.

Adolescent immunizations are defined in the following ways:

Immunization	CPT Codes	ICD and Other Codes
MMR	90707, 90710	Immunization codes: 3, 94 POV: V06.4 V Procedure: 99.48 Contraindications: POV 279, V08, 042, 200-202, 203.0, 203.1, 203.8, 204-208; or Immunization Package contraindication of "Anaphylaxis," "Immune Deficiency," "Immune Deficient," or "Neomycin Allergy." Refusals: Immunization codes 3, 94
M/R (Measles/Rubella)	90708	Immunization code: 4 Refusals: Immunization code 4 Contraindications: Immunization Package contraindication of "Anaphylaxis."
R/M (Rubella/Mumps)	90709 (old code)	Immunization code: 38 Refusals: Immunization code 38 Contraindications: Immunization Package contraindication of "Anaphylaxis."

Immunization	CPT Codes	ICD and Other Codes
Measles	90705	Immunization code: 5 POV: V04.2 V Procedure: 99.45 Evidence of Disease: POV or PCC Problem List (active or inactive) 055* Refusals: Immunization code 5 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Mumps	90704	Immunization code: 7 POV: V04.6 V Procedure: 99.46 Evidence of Disease: POV or PCC Problem List (active or inactive) 072* Refusals: Immunization code 7 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Rubella	90706	Immunization code: 6 POV: V04.3 V Procedure: 99.47 Evidence of Disease: POV or PCC Problem List (active or inactive) 056*, 771.0 Refusals: Immunization code 6 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Hepatitis B	90736, 90723, 90731 (old code), 90740, 90743-90748, G0010, Q3021, Q3023	Immunization codes: 8, 42-45, 51, 102, 104, 110, 132 Evidence of Disease: POV or PCC Problem List (active or inactive) V02.61, 070.2, 070.3 Refusals: Immunization codes 8, 42-45, 51, 102, 104, 110, 132 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Varicella	90710, 90716	Immunization codes: 21, 94 POV: V05.4 Evidence of Disease: 1) POV or PCC Problem List (active or inactive) 052*, 053*; or 2) Immunization Package contraindication of "Hx of Chicken Pox" or "Immune." Refusals: Immunization codes 21, 94 Contraindications: POV 279, V08, 042, 200-202, 203.0, 203.1, 203.8, 204-208; or Immunization Package contraindication of "Anaphylaxis," "Immune Deficiency," "Immune Deficient," or "Neomycin Allergy."

Immunization	CPT Codes	ICD and Other Codes
Tdap	90715	Immunization code: 115 Refusals: Immunization code 115 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Td	90714, 90718	Immunization codes: 9, 113 POV: V06.5 Refusals: Immunization codes 9, 113 Contraindications: Immunization Package contraindication of "Anaphylaxis."
Meningococcal	90733, 90734	Immunization codes: 32, 108, 114, 136 Refusals: Immunization codes 32, 108, 114, 136 Contraindications: Immunization Package contraindication of "Anaphylaxis."
HPV	90649, 90650	Immunization codes: 62, 118, 137 Refusals: Immunization codes 62, 118, 137 Contraindications: Immunization Package contraindication of "Anaphylaxis."

Key Logic Changes from CRS Version 11.0

1. Removed non-NMI refusals from main numerators. Refusal numerator is a separate numerator, and no longer a subnumerator.

Patient List Description

List of patients 13–17 with IZ, if any. If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had two Hep B, no IZ will be listed for Hep B.

Note: An absent value in the Numerator column means the patient did not meet the requirements for any of the immunizations. An example for a female patient age 13 with a value of “;2 MMR” which means the patient did not have one Td/Tdap, three Hepatitis B, one Varicella, one Meningococcal, and three HPV immunizations.

Measure Source

HEDIS, HP 2020 IID-11

Measure Past Performance and Long-Term Targets:

Target	Percent
HP 2020 goal for each individual IZ: Tdap, Meningococcal, HPV	80.0%

Target	Percent
HP 2020 goal for each individual IZ: varicella	90.0%

Performance Improvement Tips

1. Providers should ask about and record off-site historical immunizations (IZ type, date received and location) on PCC forms. Data entry mnemonic: HIM
2. Providers should document refusals; write "Refused" in appropriate vaccine Order box on PCC form. Data entry mnemonic: REF (Immunization, Value, Date Refused).

DU	May 25, 2011						Page 66	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Adolescent Immunizations								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical patients								
age 13	20		17			28		
# w/2:3:1 Combo or w/ Dx/Contraind/ NMI Refusal	2	10.0	0	0.0	+10.0	0	0.0	+10.0
A. # w/ Dx/ Contraind/ NMI Ref w/ % of Total 2:3:1	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ 2:3:1 Refusal	4	20.0	0	0.0	+20.0	0	0.0	+20.0
# w/1:3:2:1 Combo or w/ Dx/Contraind/ NMI Refusal	1	5.0	0	0.0	+5.0	0	0.0	+5.0
A. # w/ Dx/ Contraind/ NMI Ref w/ % of Total 1:3:2:1	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ 1:3:2:1 Refusal	4	20.0	0	0.0	+20.0	0	0.0	+20.0
# w/ 1 dose Tdap/Td or w/ Dx/ Contraind/ NMI Refusal	3	15.0	4	23.5	-8.5	6	21.4	-6.4
A. # w/ Dx/ Contraind/ NMI Ref w/ % of Total Tdap/Td	0	0.0	0	0.0	+0.0	0	0.0	+0.0
B. # w/ Tdap or w/ Dx/ Contraind/ NMI Refusal	0	0.0	0	0.0	+0.0	0	0.0	+0.0

# w/ Tdap/Td Refusal	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ 2 doses MMR or w/ DX/ Contraind/ NMI Refusal	6	30.0	0	0.0	+30.0	0	0.0	+30.0
A. # w/ Dx/ Contraind/ NMI Ref w/ % of Total MMR	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ MMR Refusal	2	10.0	0	0.0	+10.0	0	0.0	+10.0

Figure 2-33: Sample Report, Adolescent Immunizations

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Adolescent Immunizations: List of patients 13-17 with IZ, if any. If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had 2 Hep B, no IZ will be listed for Hep B.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,LINDA	000001	COMMUNITY #3	F	13	AC		;3 HPV
PATIENT2,SHERRY	000002	COMMUNITY #3	F	13	AC		;meningococcal
PATIENT22,JESSICA	000022	COMMUNITY #4	F	13	AC		;2 MMR; evid var
PATIENT23,SAMANTHA	000023	COMMUNITY #4	F	13	AC		;2 MMR;3 HEP
PATIENT24,NINA	000024	COMMUNITY #4	F	13	AC		;contra mmr;contra var
PATIENT25,RHONDA	000025	COMMUNITY #4	F	13	AC		;3 HEP;vari
PATIENT26,SARA	000026	COMMUNITY #4	F	13	AC		;3 HEP;Td
PATIENT27,AMANDA	000027	COMMUNITY #4	F	14	AC		;Tdap

Figure 2-34: Sample Patient List, Adolescent Immunizations

2.5 Childhood Diseases Group

2.5.1 Appropriate Treatment for Children with Upper Respiratory Infection

Denominators

Active Clinical patients who were ages 3 months through 18 years of age who were diagnosed with an *upper respiratory infection* during the period 6 months (180 days) prior to the report period through the first six months of the report period.

User Population patients who were ages 3 months through 18 years of age who were diagnosed with an *upper respiratory infection* during the period 6 months (180 days) prior to the report period through the first 6 months of the report period.

Numerator

Patients who were *not* prescribed an antibiotic on or within three days after diagnosis. In this measure, appropriate treatment is *not* to receive an antibiotic.

Logic Description

Age is calculated as follows: Children three months as of 6 months (180 days) of the year prior to the Report Period to 18 years as of the first 6 months of the Report Period.

In order to be included in the denominator, *all* of the following conditions must be met:

1. Patient's diagnosis of an upper respiratory infection (URI) must have occurred at an outpatient visit. Upper Respiratory Infection defined as POV 460 or 465.*. Outpatient visit defined as Service Category A, S, or O.
2. If outpatient visit was to Clinic Code 30 (Emergency Medicine), it must not have resulted in a hospitalization, defined as Service Category H, either on the same day or the next day with URI diagnosis.
3. Patient's visit must only have a diagnosis of URI. If any other diagnosis exists, the visit will be excluded.
4. The patient did not have a new or refill prescription for antibiotics within 30 days prior to the URI visit date.
5. The patient did not have an active prescription for antibiotics as of the URI visit date. "Active" prescription defined as:

$$\text{Rx Days Supply} \geq (\text{URI Visit Date} - \text{Prescription Date})$$

If multiple visits exist that meet the above criteria, the first visit will be used.

Antibiotic medications defined with medication taxonomy BGP HEDIS ANTIBIOTIC MEDS or V Procedure 99.21. (Medications are: Amoxicillin, Amox/Clavulanate, Ampicillin, Azithromycin, Cefaclor, Cefadroxil hydrate, Cefazolin, Cefdinir, Cefixime, Cefditoren, Ceftibuten, Cefpodoxime proxetil, Cefprozil, Ceftriaxone, Cefuroxime, Cephalexin, Cephradine, Ciprofloxacin, Clindamycin, Dicloxacillin, Doxycycline, Erythromycin, Ery E-Succ/Sulfisoxazole, Gatifloxacin, Levofloxacin, Lomefloxacin, Loracarbef, Minocycline, Ofloxacin, Penicillin VK, Penicillin G, Sparfloxacin, Sulfisoxazole, Tetracycline, Trimethoprim, Trimethoprim-Sulfamethoxazol.) Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients 3 months to 18 years of age with upper respiratory infection, with antibiotic prescription, if any.

Measure Source

HEDIS

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 72

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Appropriate Treatment for Children with Upper Respiratory Infection (con't)

	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical 3 months-18 yrs								
w/Upper Respiratory Infection	38		36			30		
# w/o Antibiotic Rx	37	97.4	35	97.2	+0.1	27	90.0	+7.4
User Pop 3 months-18 yrs								
w/Upper Respiratory Infection	43		38			35		
# w/o Antibiotic Rx	42	97.7	37	97.4	+0.3	32	91.4	+6.2

Figure 2-35: Sample Report, Appropriate Treatment for Children with Upper Respiratory Infection

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic						
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Appropriate Treatment for Children with Upper Respiratory Infection: List of patients 3 months to 18 years with upper respiratory infection, with antibiotic prescription, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,PAMELA	000001	COMMUNITY	#3	F	3	UP;AC	MEETS MEASURE
PATIENT2,ALICIA	000002	COMMUNITY	#3	F	7	UP;AC	MEETS MEASURE
PATIENT3,JAMES	000003	COMMUNITY	#3	M	0	UP;AC	MEETS MEASURE
PATIENT4,HENRY	000004	COMMUNITY	#3	M	12	UP;AC	MEETS MEASURE
PATIENT25,HEATHER	000025	COMMUNITY	#4	F	7	UP;AC	MEETS MEASURE
PATIENT26,DYLAN	000026	COMMUNITY	#4	M	3	UP;AC	MEETS MEASURE
PATIENT27,CODY	000027	COMMUNITY	#4	M	4	UP;AC	MEETS MEASURE
PATIENT28,KAREN	000028	COMMUNITY	#5	F	0	UP;AC	antibiotic
injection: 01/06/11 DOES NOT MEET MEASURE							

Figure 2-36: Sample Patient List, Appropriate Treatment for Children with Upper Respiratory Infection

2.5.2 Appropriate Testing for Children with Pharyngitis

Denominators

Active Clinical patients who were ages 2–18 years who were diagnosed with *pharyngitis* and prescribed an antibiotic during the period 6 months (180 days) prior to the Report Period through the first 6 months of the Report Period.

User Population patients who were ages 2–18 years who were diagnosed with *pharyngitis* and prescribed an antibiotic during the period 6 months (180 days) prior to the Report Period through the first 6 months of the Report Period.

Numerator

Patients who received a Group A strep test.

Logic Description

Age is calculated as follows: Children two years as of 6 months (180 days) of the year prior to the Report Period to 18 years as of the first 6 months of the Report Period.

In order to be included in the denominator, *all* of the following conditions must be met:

1. Patient's diagnosis of pharyngitis must have occurred at an outpatient visit. Pharyngitis defined as POV 462, 463, or 034.0. Outpatient visit defined as Service Category A, S, or O.
2. If outpatient visit was to Clinic Code 30 (Emergency Medicine), it must not have resulted in a hospitalization, defined as Service Category H, either on the same day or the next day with pharyngitis diagnosis.
3. Patient's visit must only have a diagnosis of pharyngitis. If any other diagnosis exists, the visit will be excluded.
4. The patient did not have a new or refill prescription for antibiotics within 30 days prior to the pharyngitis visit date.

5. The patient did not have an active prescription for antibiotics as of the pharyngitis visit date. "Active" prescription defined as:
6. Rx Days Supply \geq (URI Visit Date - Prescription Date)
7. The patient filled a prescription for antibiotics on or within three days after the pharyngitis visit.

If multiple visits exist that meet the above criteria, the first visit will be used.

Antibiotic medications defined with medication taxonomy BGP HEDIS ANTIBIOTIC MEDS or V Procedure 99.21. (Medications are: Amoxicillin, Amox/Clavulanate, Ampicillin, Azithromycin, Cefaclor, Cefadroxil hydrate, Cefazolin, Cefdinir, Cefixime, Cefditoren, Ceftibuten, Cefpodoxime proxetil, Cefprozil, Ceftriaxone, Cefuroxime, Cephalexin, Cephadrine, Ciprofloxacin, Clindamycin, Dicloxacillin, Doxycycline, Erythromycin, Ery E-Succ/Sulfisoxazole, Gatifloxacin, Levofloxacin, Lomefloxacin, Loracarbef, Minocycline, Ofloxacin, Penicillin VK, Penicillin G, Sparfloxacin, Sulfisoxazole, Tetracycline, Trimethoprim, Trimethoprim-Sulfamethoxazol.) Medications must not have a comment of RETURNED TO STOCK.

To be included in the numerator, a patient must have received a Group A Streptococcus test within the seven-day period beginning three days prior through three days after the Pharyngitis visit date.

Group A Streptococcus test defined as: CPT 87430 (by enzyme immunoassay), 87650-87652 (by nucleic acid), 87880 (by direct optical observation), 87081 (by throat culture), 3210F (Group A Strep Test); site-populated taxonomy BGP GROUP A STREP TESTS; and LOINC taxonomy.

Key Logic Changes from CRS Version 11.0

1. Added CPT code 3210F to Group A Strep Test definition.

Patient List Description

List of patients 2–18 years of age with pharyngitis and a Group A Strep test, if any.

Measure Source

HEDIS

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 75

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Appropriate Testing for Children with Pharyngitis (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical 2-18 yrs w/ Pharyngitis and Antibiotic Rx	10		5			8			
# w/Group A Strep Test	9	90.0	3	60.0	+30.0	2	25.0	+65.0	
User Pop 2-18 yrs w/ Pharyngitis and Antibiotic Rx	11		7			10			
# w/Group A Strep Test	9	81.8	4	57.1	+24.7	2	20.0	+61.8	

Figure 2-37: Sample Report, Appropriate Testing for Children with Pharyngitis

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Appropriate Testing for Children with Pharyngitis: List of patients 2-18 years with pharyngitis and a Group A Strep test, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT1,MICHAEL ANTIGEN (STREP A)	000001	COMMUNITY #1	M	9	UP;AC	03/19/11	RAPID
PATIENT2,JOSEPH ANTIGEN (STREP A)	000002	COMMUNITY #1	M	12	UP;AC	05/01/11	RAPID
PATIENT3,LESTER	000003	COMMUNITY #1	M	13	UP		
PATIENT24,MONICA ANTIGEN (STREP A)	000024	COMMUNITY #2	F	5	UP;AC	01/23/11	RAPID
PATIENT25,MICHAEL JAMES ANTIGEN (STREP A)	000025	COMMUNITY #2	M	7	UP;AC	03/12/11	RAPID

Figure 2-38: Sample Patient List, Appropriate Testing for Children with Pharyngitis

2.6 Cancer Related Measure Topics

2.6.1 Cancer Screening: Pap Smear Rates

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 55.7% for the proportion of female patients ages 21 through 64 without a documented history of hysterectomy who have had a Pap screen within the previous three years.

Denominators

Female Active Clinical patients ages 21 through 64 without documented history of Hysterectomy. (GPRA Denominator)

Female User Population patients ages 21 through 64 without a documented history of Hysterectomy.

Numerators

Patients with a Pap smear documented in the past three years.

Note: This numerator does *not* include refusals. (GPRA Denominator)

Patients with documented Pap smear refusal in past year.

Logic Description

Age of the patient is calculated at the beginning of the report period. Patients must be at least 21 years of age at the beginning of the report period and less than 65 years of age as of the end of the report period.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Hysterectomy	51925, 56308 (old code), 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200-58294, 58548, 58550-58554, 58570-58573, 58951, 58953-58954, 58956, 59135	V Procedure: 68.4-68.8 V POV: 618.5, V88.01, V88.03 Women's Health: Procedure called Hysterectomy.		

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Pap Smear	88141-88167, 88174-88175, G0123, G0124, G0141, G0143-G0145, G0147, G0148, P3000, P3001, Q0091	V Lab: PAP SMEAR POV: V67.01 Follow-up Vaginal Pap Smear V76.2 Screen Mal Neop-Cervix V72.32 Encounter for Pap Cervical Smear to Confirm Findings of Recent Normal Smear Following Initial Abnormal Smear V72.3 Gynecological Examination, Pap Cervical Smear as Part of General Gynecological Exam, Pelvic Exam (annual) (periodic) (old code, to be counted for visits prior to 10/1/04 only) V76.47 Vaginal Pap Smear for Post-Hysterectomy Patients 795.0*, 795.10-16, 795.19 V Procedure: 91.46 Women's Health: Procedure called Pap Smear and where the result does NOT have "ERROR/DISREGARD"	Yes	BGP PAP SMEAR TAX
Refusal (in past year)	Refusal of codes: 88141-88167, 88174-88175, G0123, G0124, G0141, G0143-G0145, G0147, G0148, P3000, P3001, Q0091	Refusals: Lab Test Value Pap Smear		

Key Logic Changes from CRS Version 11.0

1. Added CPT codes 57540, 57545, 57550, 57555, 57556 to Hysterectomy definition.
2. Added refusal of CPT/ HCPCS codes 88141-88167, 88174-88175, G0123, G0124, G0141, G0143-G0145, G0147, G0148, P3000, P3001, Q0091 to Pap Smear refusal definition.
3. Removed POV code V72.31 from Pap Smear definition.

Patient List Description

List of women 21–64 with documented Pap Smear or refusal, if any.

Measure Source

HP 2020 C-15

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	59.0%
IHS FY 2009 Performance	59.0%
IHS FY 2008 Performance	59.0%
IHS FY 2007 Performance	59.0%
IHS FY 2006 Performance	59.0%
IHS FY 2005 Performance	60.0%
IHS FY 2004 Performance	58.0%
IHS FY 2003 Performance	61.0%
IHS FY 2002 Performance	62.0%
<i>HP 2020 Goal</i>	<i>93.0%</i>

Performance Improvement Tips

1. Providers should ask about and record off-site tests (date received and location) on PCC forms. Data entry mnemonic: HPAP
2. Providers should document refusals; write “Refused” in Pap Order box on PCC form. Data entry mnemonic: REF (Lab Test Value, Date Refused).

DU	May 25, 2011						Page 77	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Cancer Screening: Pap Smear Rates (con't)								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Female Active Clinical								
21-64 yrs (GPRA)	500		377			320		
# w/Pap Smear recorded								
w/in 3 years-No Refusals								
(GPRA)	202	40.4	179	47.5	-7.1	147	45.9	-5.5
# w/ Pap Smear								
Refusal	8	1.6	0	0.0	+1.6	0	0.0	+1.6
# Female User Pop								

21-64 years	833		699			614		
# w/Pap Smear recorded w/in 3 years								
-No Refusals	220	26.4	196	28.0	-1.6	159	25.9	+0.5
# w/ Pap Smear Refusal	8	1.0	0	0.0	+1.0	0	0.0	+1.0

Figure 2-39: Sample Report, Cancer Screening: Pap Smear Rates

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Cancer Screening: Pap Smear Rates: List of women 21-64 with documented Pap smear or refusal, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,EVELYN	000001	COMMUNITY #1	F	21	UP	05/05/10	795.06
PATIENT2,MICHELLE	000002	COMMUNITY #1	F	22	UP,AC	05/31/11	Lab
PATIENT3,KAITLYN	000003	COMMUNITY #1	F	22	UP,AC	04/03/11	V67.01
PATIENT4,BRITNEY	000004	COMMUNITY #1	F	22	UP,AC	01/10/11	V72.3
PATIENT5,KATY	000005	COMMUNITY #1	F	22	UP,AC	05/08/10	88150

Figure 2-40: Sample Patient List, Cancer Screening: Pap Smear rates

2.6.2 Cancer Screening: Mammogram Rates

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 46.9% for the proportion of female patients ages 52 through 64 who have had mammography screening within the last 2 years.

Denominators

Female Active Clinical patients ages 52 through 64 without a documented history of bilateral mastectomy or 2 separate unilateral mastectomies. (GPRA Denominator)

Female Active Clinical patients ages 42 and older without a documented history of bilateral mastectomy or 2 separate unilateral mastectomies.

Female User Population patients ages 52 through 64 without a documented history of bilateral mastectomy or 2 separate unilateral mastectomies.

Female User Population patients ages 42 and older without a documented history of bilateral mastectomy or 2 separate unilateral mastectomies. (GPRA Developmental Denominator)

Numerators

All patients who had a Mammogram documented in the past two years.

Note: This numerator does *not* include refusals. (GPRA Numerator)

Patients with documented mammogram refusal in the past year

Logic Description

Age of the patient is calculated at the beginning of the Report Period. For all denominators, patients must be at least the minimum age as of the beginning of the Report Period. For the 52–64 denominators, the patients must be less than 65 years of age as of the end of the Report Period.

	CPT Codes	ICD and Other Codes
Bilateral Mastectomy	19300.50-19307.50 or old code 19180, 19200, 19220 OR 19300-19307 w/modifier 09950 (.50 and 09950 modifiers indicate bilateral) OR 19240, with modifier of .50 or 09950	V Procedure: 85.42, 85.44, 85.46, 85.48
Unilateral Mastectomy	Must have 2 separate occurrences on 2 different dates of service. 19300-19307, or old codes 19180, 19200, 19220, 19240	Must have 2 separate occurrences on 2 different dates of service. 85.41, 85.43, 85.45, 85.47
Mammogram	V Rad or VCPT: 77052-77059, 76090 (old code), 76091 (old code), 76092 (old code), G0206, G0204, G0202	POV: V76.11, V76.12, 793.80 Abnormal mammogram, unspecified; 793.81 Mammographic microcalcification; 793.89 Other abnormal findings on radiological exam of breast V Procedure: 87.36-87.37 Women's Health: Mammogram Screening, Mammogram Dx Bilat, Mammogram Dx Unilat and where the mammogram result does NOT have "ERROR/DISREGARD"
Refusal (in past year)	V Rad Mammogram for CPT: 77052-77059, 76090 (old code), 76091 (old code), 76092 (old code), G0206, G0204, G0202	

Key Logic Changes from CRS Version 11.0

1. Added CPT code 77052 to Mammogram Screening and Mammogram refusal definitions.

Patient List Description

List of women 42+ with mammogram/refusal, if any.

Measure Source

HP 2020 C-17

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	48.0%
IHS FY 2009 Performance	45.0%
IHS FY 2008 Performance	45.0%
IHS FY 2007 Performance	43.0%
IHS FY 2006 Performance	41.0%
IHS FY 2005 Performance	41.0%
IHS FY 2004 Performance	40.0%
IHS FY 2003 Performance	40.0%
IHS FY 2002 Performance	42.0%
<i>HP 2020 Goal</i>	<i>81.1%</i>

Performance Improvement Tips

1. Providers should ask about and record off-site mammogram procedures (date received and location) on PCC forms. Data entry mnemonic: HRAD.
2. Providers should document refusals; write “Refused” in Mammogram Order box on PCC form. Data entry mnemonic: REF (Mammogram, Procedure (CPT) Code, Date Refused).

DU

May 25, 2011

Page 79

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Cancer Screening: Mammogram Rates (con't)

	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Female Active Clinical								
52-64 (GPRA)	97		60			47		
# w/Mammogram recorded w/in								
2 years-No Refusals								
(GPRA)	31	32.0	22	36.7	-4.7	22	46.8	-14.8
# w/ Mammogram								
Refusal	7	7.2	0	0.0	+7.2	0	0.0	+7.2
# Female Active Clinical								
42+ (GPRA Dev.)	293		188			163		
# w/Mammogram recorded								
w/in 2 years-No Refusals								
(GPRA Dev.)	63	21.5	62	33.0	-11.5	54	33.1	-11.6
# w/ Mammogram								

Refusal	9	3.1	0	0.0	+3.1	0	0.0	+3.1
# Female User Pop 52-64	175		122			102		
# w/Mammogram recorded w/in 2 years-No Refusals	34	19.4	26	21.3	-1.9	23	22.5	-3.1
# w/ Mammogram Refusal	7	4.0	0	0.0	+4.0	0	0.0	+4.0
# Female User Pop 42+	513		380			333		
# w/Mammogram recorded w/in 2 years-No Refusals	67	13.1	69	18.2	-5.1	58	17.4	-4.4
# w/ Mammogram Refusal	9	1.8	0	0.0	+1.8	0	0.0	+1.8

Figure 2-41: Sample Report, Cancer Screening: Mammogram rates

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Cancer Screening: Mammogram Rates: List of women 42+ with mammogram/refusal, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,CARLA	000001	COMMUNITY #1	F	43	UP;AC - >41 10/01/02 CPT 77052	
PATIENT2,CRYSTAL	000002	COMMUNITY #1	F	42	UP - >41	
PATIENT3,ALEXA	000003	COMMUNITY #1	F	45	UP;AC - >41 04/24/10 76090	
PATIENT4,HANNAH	000004	COMMUNITY #1	F	42	UP - >41	
PATIENT5,MARTHA	000005	COMMUNITY #1	F	43	UP - >41	
PATIENT6,TARA	000006	COMMUNITY #1	F	44	UP;AC - >41 01/15/11 ref CPT	
PATIENT7,CAROL LYNN	000007	COMMUNITY #1	F	44	UP;AC - >41 03/05/11 76092	
PATIENT8,MARY ANN	000008	COMMUNITY #1	F	52	UP;AC - >41, 52-64	
PATIENT9,BARBARA	000009	COMMUNITY #1	F	52	UP;AC - >41, 52-64 04/22/11 77057	

Figure 2-42: Sample Patient List, Cancer Screening: Mammogram rates

2.6.3 Colorectal Cancer Screening

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 36.7% for the proportion of clinically appropriate patients ages 51–80 who have received colorectal screening.

Denominators

All Active Clinical patients ages 51–80 without a documented diagnosis of colorectal cancer or total colectomy. Broken down by gender. (GPRA Denominator)

All *User Population patients* ages 51–80 without any documented diagnosis of colorectal cancer or total colectomy.

Numerators

Patients who have had any CRC screening, defined as any of the following: (1) Fecal Occult Blood Test (FOBT) or Fecal Immunochemical Test (FIT) during the Report period; (2) flexible sigmoidoscopy or double contrast barium enema in the past 5 years; or (3) colonoscopy in the past 10 years.

Note: This numerator does *not* include refusals. (GPRA Numerator)

Patients with documented CRC screening refusal in the past year.

Patients with FOBT or FIT during the Report Period.

Patients with a flexible sigmoidoscopy or double contrast barium enema in the past 5 years or a colonoscopy in the past 10 years.

Patients with a flexible sigmoidoscopy in the past 5 years or a colonoscopy in the past 10 years.

Patients with a flexible sigmoidoscopy and double contrast barium enema in the past 5 years or a colonoscopy in the past 10 years.

Logic Description

Age is calculated at the beginning of the Report Period.

Denominator Exclusions

Any diagnosis ever of one of the following:

1. **Colorectal Cancer:** POV: 153.*, 154.0, 154.1, 197.5, V10.05; CPT G0213–G0215 (old codes), G0231 (old code).
2. **Total Colectomy:** CPT 44150–44151, 44152 (old code), 44153 (old code), 44155–44158, 44210–44212; V Procedure 45.8 (old code).

Colorectal cancer screening definition: The most recent of any of the following tests and procedures during applicable timeframes. CRS identifies the tests and procedures described in the numerators above with the following codes:

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Colorectal Cancer Screening (CRS looks for the most recent of any of the following during timeframes specified in numerator section above)				
Fecal Occult Blood lab test (FOBT) or Fecal Immuno-chemical Test (FIT)	82270, 82274, 89205 (old code), G0107 (old code), G0328, G0394 (old code)		Yes	BGP GPRA FOB TESTS
Flexible Sigmoidoscopy	45330-45345, G0104	V Procedure: 45.24		
Double contrast barium enema	VRad 74280, G0106, G0120			
Colonoscopy	44388-44394, 44397, 45355, 45378-45387, 45391, 45392, G0105, G0121	V POV: V76.51 Colon screening V Procedure: 45.22, 45.23, 45.25, 45.42, 45.43		

Refusal definition: Any of the following in the past year:

	CPT Codes	ICD and Other Codes	Taxonomy
Refusals	FOBT or FIT: 82270, 82274, 89205 (old code), G0107 (old code), G0328, or G0394 (old code) Flexible Sigmoidoscopy: 45330-45345, G0104 DCBE: 74280, G0106, G0120 Colonoscopy: 44388-44394, 44397, 45355, 45378-45387, 45391, 45392, G0105, G0121	Flexible Sigmoidoscopy V Procedure: 45.24, 45.42 Colonoscopy V Procedure: 45.22, 45.23, 45.25, 45.42, 45.43	V Lab Fecal Occult Blood Test

Key Logic Changes from CRS Version 11.0

1. Added codes 56490-6, 56491-4, 57905-2, and 58453-2 to LOINC taxonomy for FOBT.

Patient List Description

List of patients 51–80 with CRC screening or refusal, if any.

Measure Source

HEDIS, HP 2020 C-16

Measure Past Performance and Long-Term Targets:

Performance	Percent
IHS FY 2010 Performance	37.0%
IHS FY 2009 Performance	33.0%
IHS FY 2008 Performance	29.0%
IHS FY 2007 Performance	26.0%
IHS FY 2006 Performance	22.0%
HP 2020 Goal	70.5%

Performance Improvement Tip

Providers should ask about and record off-site historical tests (test type, date received and location) on PCC forms. Data entry mnemonics: HBE (barium enema); HCOL (colonoscopy); HFOB (Fecal Occult Blood); HSIG (sigmoidoscopy). Providers should also enter as a refusal if the patient refuses the colorectal cancer screening. Refusals may be entered with the data entry mnemonic of REF (refusal).

DU	May 25, 2011						Page 82	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Colorectal Cancer Screening (con't)								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
AC Pts 51-80 w/o colorectal cancer or total colectomy (GPRA)	316		195			152		
# w/ CRC screening -No Refusals (GPRA)	64	20.3	49	25.1	-4.9	28	18.4	+1.8
# w/ CRC Screening Refusal	10	3.2	0	0.0	+3.2	0	0.0	+3.2
# w/FOBT/FIT during Report period	12	3.8	11	5.6	-1.8	0	0.0	+3.8
# w/Flex Sig, DCBE, or Colonoscopy	55	17.4	40	20.5	-3.1	28	18.4	-1.0
# w/Flex Sig or Colonoscopy	49	15.5	31	15.9	-0.4	20	13.2	+2.3
# w/Flex Sig & DCBE or Colonoscopy	46	14.6	29	14.9	-0.3	18	11.8	+2.7
Male Active Clinical 51-80	151		93			65		
# w/ CRC screening -No Refusals	27	17.9	18	19.4	-1.5	9	13.8	+4.0
# w/ CRC Screening								

Refusal	5	3.3	0	0.0	+3.3	0	0.0	+3.3
# w/FOBT/FIT during Report period	7	4.6	3	3.2	+1.4	0	0.0	+4.6
# w/Flex Sig, DCBE, or Colonoscopy	22	14.6	15	16.1	-1.6	9	13.8	+0.7
# w/Flex Sig or Colonoscopy	20	13.2	14	15.1	-1.8	8	12.3	+0.9
# w/Flex Sig & DCBE or Colonoscopy	20	13.2	14	15.1	-1.8	8	12.3	+0.9

Figure 2-43: Sample Report, Colorectal Cancer Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Colorectal Cancer Screening: List of patients 51-80 with CRC screening or refusal, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DANIELLE G0107	000001	COMMUNITY #1	F	51	UP	FOB: 08/19/11 CPT
PATIENT2,MARIE cpt	000002	COMMUNITY #1	F	51	UP,AC	02/12/11 ref DCBE
PATIENT3,MARY ANN	000003	COMMUNITY #1	F	52	UP,AC	
PATIENT4,BOBBIE 74280	000004	COMMUNITY #1	F	52	UP,AC	BE: 01/25/11 RAD
PATIENT5,WINONA	000005	COMMUNITY #1	F	53	UP,AC	
PATIENT6,DARLENE 45.24; BE: 03/31/09 RAD 74280	000006	COMMUNITY #1	F	54	UP,AC	SIG: 04/07/08
PATIENT7,JOYCE V76.51	000007	COMMUNITY #1	F	57	UP,AC	COLO: 07/07/11 DX
PATIENT8,LOUISE	000008	COMMUNITY #1	F	62	UP	03/19/11 ref COLO proc

Figure 2-44: Sample Patient List, Colorectal Cancer Screening

2.6.4 Comprehensive Cancer Screening

GPRA Measure Description

During FY 2011, establish the proportion of patients ages 21–80 who received a comprehensive cancer screening.

Denominators

Active Clinical patients ages 21–80 who are eligible for cervical cancer, breast cancer, and/or colorectal cancer screening. (GPRA Developmental Denominator)

- Active Clinical female patients* ages 21–80.
- Active Clinical male patients* ages 51–80.

Numerators

Patients who have had all screenings for which they are eligible.

Note: This numerator does *not* include refusals. (GPRA Developmental Numerator)

- a. Female patients with cervical cancer, breast cancer, and/or colorectal cancer screening.
- b. Male patients with colorectal cancer screening.

Logic Description

Age is calculated at the beginning of the Report Period.

Cervical Cancer Screening definition: To be eligible for this screening, patients must be female Active Clinical ages 21 through 64 and not have a documented history of hysterectomy. Patients must be at least 21 years of age at the beginning of the Report Period and less than 65 years of age as of the end of the Report Period. To be counted as having the screening, the patient must have had a Pap Smear documented in the past three years.

CRS identifies the tests and procedures described in the numerators above with the following codes:

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Hysterectomy	51925, 56308 (old code), 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200-58294, 58548, 58550-58554, 58570-58573, 58951, 58953-58954, 58956, 59135	V Procedure: 68.4-68.8 V POV: 618.5, V88.01, V88.03 Women's Health: Procedure called Hysterectomy.		

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Pap Smear	88141-88167, 88174-88175, G0123, G0124, G0141, G0143- G0145, G0147, G0148, P3000, P3001, Q0091	V Lab: PAP SMEAR POV: V67.01 Follow-up Vaginal Pap Smear V76.2 Screen Mal Neop-Cervix V72.32 Encounter for Pap Cervical Smear to Confirm Findings of Recent Normal Smear Following Initial Abnormal Smear V72.3 Gynecological Examination, Pap Cervical Smear as Part of General Gynecological Exam, Pelvic Exam (annual) (periodic) (old code, to be counted for visits prior to 10/1/04 only) V76.47 Vaginal Pap Smear for Post-Hysterectomy Patients 795.0*, 795.10-16, 795.19 V Procedure: 91.46 Women's Health: Procedure called Pap Smear	Yes	BGP PAP SMEAR TAX

Breast Cancer Screening definition: To be eligible for this screening, patients must be female Active Clinical ages 52 through 64 and not have a documented history ever of bilateral mastectomy or two separate unilateral mastectomies. Patients must be at least age 52 as of the beginning of the Report Period and must be less than 65 years of age as of the end of the Report Period. To be counted as having the screening, the patient must have had a Mammogram documented in the past two years.

CRS identifies the tests and procedures described in the numerators above with the following codes:

	CPT Codes	ICD and Other Codes
Bilateral Mastectomy	19300.50-19307.50 or old code 19180, 19200, 19220 OR 19300-19307 w/modifier 09950 (.50 and 09950 modifiers indicate bilateral) OR 19240, with modifier of .50 or 09950	V Procedure: 85.42, 85.44, 85.46, 85.48
Unilateral Mastectomy	Must have 2 separate occurrences on 2 different dates of service. 19300-19307, or old codes 19180, 19200, 19220, 19240	Must have 2 separate occurrences on 2 different dates of service. 85.41, 85.43, 85.45, 85.47

	CPT Codes	ICD and Other Codes
Mammogram	V Rad or VCPT: 77052-77059, 76090 (old code), 76091 (old code), 76092 (old code), G0206, G0204, G0202	POV: V76.11, V76.12, 793.80 Abnormal mammogram, unspecified; 793.81 Mammographic microcalcification; 793.89 Other abnormal findings on radiological exam of breast V Procedure: 87.36-87.37 Women's Health: Mammogram Screening, Mammogram Dx Bilat, Mammogram Dx Unilat and where the mammogram result does NOT have "ERROR/DISREGARD".

Colorectal cancer screening definition: To be eligible for this screening, patients must be Active Clinical ages 51–80 and not have a documented history ever of colorectal cancer or total colectomy. To be counted as having the screening, patients must have had any of the following: (1) FOBT or FIT during the Report Period; (2) flexible sigmoidoscopy or double contrast barium enema in the past 5 years; or (3) colonoscopy in the past 10 years.

The most recent of any of the following tests and procedures during applicable timeframes. CRS identifies the tests and procedures described in the numerators above with the following codes:

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Colorectal Cancer	G0213-G0215, G0231	V POV: 153.*, 154.0, 154.1, 197.5, V10.05		
Total Colectomy	44150-44151, 44152 (old code), 44153 (old code), 44155-44158, 44210-44212	V Procedure: 45.8 (old code)		
Fecal Occult Blood lab test (FOBT) or Fecal Immuno-chemical Test (FIT)	82270, 82274, 89205 (old code), G0107 (old code), G0328, G0394 (old code)		Yes	BGP GPRA FOB TESTS
Flexible Sigmoidoscopy	45330-45345, G0104	V Procedure: 45.24		
Double contrast barium enema	VRad 74280, G0106, G0120			
Colonoscopy	44388-44394, 44397, 45355, 45378-45387, 45391, 45392, G0105, G0121	V POV: V76.51 Colon screening V Procedure: 45.22, 45.23, 45.25, 45.42, 45.43		

Key Logic Changes from CRS Version 11.0

1. Added CPT codes 57540, 57545, 57550, 57555, and 57556 to Hysterectomy definition.
2. Added CPT code 77052 to Mammogram Screening definition.
3. Added codes 56490-6, 56491-4, 57905-2, and 58453-2 to LOINC taxonomy for FOBT.
4. Removed POV code V72.31 from Pap Smear definition.

Patient List Description

List of patients 21–80 with comprehensive cancer screening, if any.

Measure Source

Not Available

Measure Past Performance and Long-Term Targets

None

Performance Improvement Tip

1. Providers should ask about and record off-site Pap tests (date received and location) on PCC forms. Data entry mnemonic: HPAP
2. Providers should ask about and record off-site mammogram procedures (date received and location) on PCC forms. Data entry mnemonic: HRAD.
3. Providers should ask about and record off-site historical colorectal cancer tests (test type, date received and location) on PCC forms. Data entry mnemonics: HBE (barium enema); HCOL (colonoscopy); HFOB (Fecal Occult Blood); HSIG (sigmoidoscopy).

DU	May 25, 2011					Page 87		
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Comprehensive Cancer Screening								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical 21-80								
(GPRA Dev.)	721		509			424		

# w/ Comprehensive Cancer Screening-No Refusals (GPRA Dev.)	226	31.3	195	38.3	-7.0	153	36.1	-4.7
A. Female 21-80	570		416			359		
A. # Female w/all Screens	199	34.9	177	42.5	-7.6	144	40.1	-5.2
B. Male 51-80	151		93			65		
B. # Male w/CRC Screen	27	17.9	18	19.4	-1.5	9	13.8	+4.0

Figure 2-45: Sample Report, Comprehensive Cancer Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Comprehensive Cancer Screening: List of patients 21-80 with comprehensive cancer screening, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,DANIELLE POV 795.0	000001	COMMUNITY #1	F	21	AC;PAP	PAP: 05/05/02	
PATIENT2,MARIE	000002	COMMUNITY #1	F	51	AC;PAP		
PATIENT3,MARY ANN CPT 77055	000003	COMMUNITY #1	F	52	AC;MAM	MAM: 07/06/02	
PATIENT4,BOBBIE DX V76.51	000004	COMMUNITY #1	F	53	AC,PAP,MAM,CRCS	CRCS: 07/20/03	
PATIENT5,WINONA CPT 77052	000005	COMMUNITY #1	F	57	AC,PAP,MAM,CRCS	MAM: 10/01/02	
PATIENT6,HARRY 45.24	000006	COMMUNITY #1	M	56	AC;CRCS	CRCS: 04/07/99	
PATIENT7,LARRY	000007	COMMUNITY #1	M	57	AC,CRCS		
PATIENT8,BARRY CPT SIG 45330	000008	COMMUNITY #1	M	63	AC,CRCS	CRCS: 02/18/03	

Figure 2-46: Sample Patient List, Comprehensive Cancer Screening

2.6.5 Tobacco Use and Exposure Assessment

Denominators

Active Clinical patients ages five and older. Broken down by gender and age groups (5–13, 14–17, 18–24, 25–44, 45–64, and 65 and older), based on HP 2010 age groups.

All Pregnant female User Population patients with no documented miscarriage or abortion.

All User Population patients ages five and older. Broken down by gender.

Numerators

Patients who have been screened for tobacco use during the Report Period.

Patients identified as current tobacco users during the Report Period, both smokers and smokeless users.

- a. Patients identified as current smokers during the Report Period.
- b. Patients identified as current smokeless tobacco users during the Report Period.

Patients identified as exposed to environmental tobacco smoke (ETS) (second-hand smoke) during the Report Period.

Logic Description

Ages are calculated at beginning of Report Period.

For screening, an additional eight months is included for patients who were pregnant during the Report Period but who had their tobacco assessment prior to that.

CRS uses the following codes to define the denominators and numerators:

	CPT Codes	ICD and Other Codes
Pregnancy (At least two visits during the past 20 months. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the past 20 months, CRS will use the first two visits in the 20-month period. The patient must not have a documented miscarriage or abortion occurring after the second pregnancy-related visit. In addition, the patient must have at least one pregnancy-related visit occurring during the reporting period.) An additional 8 months is included for patients who were pregnant during the Report period but who had their tobacco assessment prior to that.		V POV: V22.0-V23.9, V72.42, 640.*-649.*, 651.*-676.*
Miscarriage (after second pregnancy POV in past 20 months)	59812, 59820, 59821, 59830	V POV: 630, 631, 632, 633*, 634*
Abortion (after second pregnancy POV in past 20 months)	59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	V POV: 635*, 636* 637* Procedure: 69.01, 69.51, 74.91, 96.49
Screened for Tobacco Use (time frame for pregnant patients is past 20 months)	D1320, 99406, 99407, G0375 (old code), G0376 (old code), G8453 (old code), G8455-G8457 (old codes), G8402 (old code), 1034F (Current Tobacco Smoker), 1035F (Current Smokeless Tobacco User), 1036F (Current Tobacco Non-User), 1000F (Tobacco Use Assessed)	V POV or current Active Problem List: 305.1, 305.1* (old codes), 649.00-649.04, V15.82 Patient Education codes: "TO-", "-TO", "-SHS", 305.1, 305.1* (old codes), 649.00-649.04, V15.82, D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455-G8457 (old codes), G8402 (old code), or G8453 (old code) Dental code: 1320
Tobacco Users (time frame for pregnant patients is past 20 months)	99406, 99407, G0375 (old code), G0376 (old code), 1034F (Current Tobacco Smoker), 1035F (Current Smokeless Tobacco User), G8455 (old code), G8456 (old code), G8402 (old code), G8453 (old code)	V POV or current Active Problem List: 305.1, 305.10-305.12 (old codes), or 649.00-649.04 Dental code: 1320

	CPT Codes	ICD and Other Codes
Current Smokers (time frame for pregnant patients is past 20 months)	99406, 99407, G0375 (old code), G0376 (old code), 1034F (Current Tobacco Smoker), G8455 (old code), G8402 (old code), G8453 (old code)	V POV or current Active Problem List: 305.1, 305.10-305.12 (old codes), or 649.00-649.04 Dental code: 1320
Current Smokeless (time frame for pregnant patients is past 20 months)	1035F (Current Smokeless Tobacco User), G8456 (old code)	

For numerator definitions, all existing national Tobacco, TOBACCO (SMOKING), TOBACCO (SMOKELESS–CHEWING/DIP), and TOBACCO (EXPOSURE) Health Factors are listed below with the numerator to which they apply.

Health Factor	Numerator
Ceremonial	Screened (does NOT count as Smoker)
Cessation-Smokeless	Screened; Tobacco Users; Smokeless User
Cessation-Smoker	Screened; Tobacco Users; Smoker
Current Smokeless	Screened; Tobacco Users; Smokeless User
Current Smoker	Screened; Tobacco Users; Smoker
Current Smoker, status unknown	Screened; Tobacco Users; Smoker
Current smoker, every day	Screened; Tobacco Users; Smoker
Current smoker, some day	Screened; Tobacco Users; Smoker
Non-Tobacco User	Screened
Previous Smokeless	Screened
Previous (Former) Smokeless	Screened
Previous Smoker	Screened
Previous (Former) Smoker	Screened
Smoke Free Home	Screened
Smoker In Home	Screened; ETS
Current Smoker & Smokeless	Screened; Tobacco Users; Smoker; Smokeless User
Exposure To Environmental Tobacco Smoke	Screened; ETS

Key Logic Changes from CRS Version 11.0

1. For Tobacco Screening, Tobacco Users, Smokers, and Smokeless, noted that G8453, G8455, G8456, G8457, and G8402 are old codes.
2. Updated patient list.

Patient List Description

List of patients 5 and older with documented tobacco screening, if any.

Measure Source

HP 2020 TU-1.1 Cigarette smoking 18 and older; TU-1.2 Smokeless tobacco use 18 and older; TU-11 Exposure to ETS-nonsmokers 3 and older

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2008 Performance (Screening)	54.0%
IHS FY 2005 Performance (Screening)	34.0%
IHS FY 2004 Performance (Screening)	27.0%

Performance	Percent
IHS FY 2008 Performance (Tobacco Users)	29.0%
HP 2020 Goals: TU-1.1 (Cigarette smoking 18 and older): 12%; TU-1.2 (Smokeless tobacco use 18 and older): 0.3%; TU-11 (Exposure to ETS-non smokers 18 and older): 68%	

DU	May 25, 2011							Page 90	
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Tobacco Use and Exposure Assessment (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
%	PREV YR	%	CHG from	BASE	%	CHG from	PREV YR %	BASE %	
		PERIOD		PERIOD		PERIOD			
# Active Clinical Pts									
=> 5		1,357		1,031			911		
# w/Tobacco									
Screening		624	46.0	426	41.3	+4.7	328	36.0 +10.0	
# Tobacco Users w/ % of									
Total Screened		294	47.1	165	38.7	+8.4	130	39.6 +7.5	
A. # Smokers w/ % of									
Total Tobacco Users		277	94.2	164	99.4	-5.2	130	100.0 -5.8	
B. # Smokeless Tobacco									
Users w/ % of Total									
Tobacco Users		29	9.9	5	3.0	+6.8	3	2.3 +7.6	
# exposed to ETS/									
smoker in home w/ % of									
Total Screened		4	0.6	2	0.5	+0.2	1	0.3 +0.3	
# Male Active Clinical									

ages => 5	570		430			374		
# w/Tobacco Screening	230	40.4	152	35.3	+5.0	128	34.2	+6.1
# Tobacco Users w/ % of Total Screened	136	59.1	69	45.4	+13.7	59	46.1	+13.0
A. # Smokers w/ % of Total Tobacco Users	124	91.2	69	100.0	-8.8	59	100.0	-8.8
B. # Smokeless Tobacco Users w/ % of Total Tobacco Users	22	16.2	2	2.9	+13.3	3	5.1	+11.1
# exposed to ETS/ smoker in home w/ % of Total Screened	1	0.4	0	0.0	+0.4	1	0.8	-0.3
# Female Active Clinical ages => 5	787		601			537		
# w/Tobacco Screening	394	50.1	274	45.6	+4.5	200	37.2	+12.8
# Tobacco Users w/ % of Total Screened	158	40.1	96	35.0	+5.1	71	35.5	+4.6
A. # Smokers w/ % of Total Tobacco Users	153	96.8	95	99.0	-2.1	71	100.0	-3.2
B. # Smokeless Tobacco Users w/ % of Total Tobacco Users	7	4.4	3	3.1	+1.3	0	0.0	+4.4
# exposed to ETS/ smoker in home w/ % of Total Screened	3	0.8	2	0.7	+0.0	0	0.0	+0.8

Figure 2-47: Sample Report, Tobacco Use Assessment Tobacco Use and Exposure Assessment

DU	May 25, 2011						Page 95
*** IHS 2011 Selected Measures with Community Specified Report ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Previous Year Period: Jan 01, 2010 to Dec 31, 2010							
Baseline Period: Jan 01, 2000 to Dec 31, 2000							

Tobacco Use and Exposure Assessment (con't)							
TOTAL ACTIVE CLINICAL POPULATION							
Age Distribution							
	5-13	14-17	18-24	25-44	45-64	65 and older	
CURRENT REPORT PERIOD							
# Active Clinical	171	74	177	448	371	116	
# w/Tobacco Screening	10	19	102	243	199	51	
% w/Tobacco Screening	5.8	25.7	57.6	54.2	53.6	44.0	
# Tobacco Users	3	8	46	118	104	15	
% Tobacco Users w/ % of							
Total Screened	30.0	42.1	45.1	48.6	52.3	29.4	
# Smokers	2	8	43	108	102	14	

% Smokers w/ % of Total Tobacco Users	66.7	100.0	93.5	91.5	98.1	93.3
# Smokeless	1	0	3	15	8	2
% Smokeless w/ % of Total Tobacco Users	33.3	0.0	6.5	12.7	7.7	13.3
# ETS/Smk Home	0	0	0	3	1	0
% ETS/Smk Home w/ % of Total Screened	0.0	0.0	0.0	1.2	0.5	0.0
PREVIOUS YEAR PERIOD						
# Active Clinical	176	62	168	323	238	64
# w/Tobacco Screening	11	14	87	147	128	39
% w/Tobacco Screening	6.3	22.6	51.8	45.5	53.8	60.9
# Tobacco Users	0	5	41	60	50	9
% Tobacco Users w/ % of Total Screened	0.0	35.7	47.1	40.8	39.1	23.1
# Smokers	0	5	40	60	50	9
% Smokers w/ % of Total Tobacco Users	0.0	100.0	97.6	100.0	100.0	100.0
# Smokeless	0	0	1	3	1	0
% Smokeless w/ % of Total Tobacco Users	0.0	0.0	2.4	5.0	2.0	0.0
# ETS/Smk Home	0	0	0	2	0	0
% ETS/Smk Home w/ % of Total Screened	0.0	0.0	0.0	1.4	0.0	0.0
CHANGE FROM PREV YR %						
# w/Tobacco Screening	-0.4	+3.1	+5.8	+8.7	-0.1	-17.0
Tobacco Users	+30.0	+6.4	-2.0	+7.7	+13.2	+6.3
Smokers	+66.7	+0.0	-4.1	-8.5	-1.9	-6.7
Smokeless	+33.3	+0.0	+4.1	+7.7	+5.7	+13.3
ETS	+0.0	+0.0	+0.0	-0.1	+0.5	+0.0

Figure 2-48: Sample Age Breakdown Report, Tobacco Use Assessment

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Tobacco Use and Exposure Assessment: List of patients 5 and older with documented tobacco screening, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,CHESTER	000001	COMMUNITY #1	F	7	UP,AC	01/10/11	SCREEN
PATIENT2,JUAN	000002	COMMUNITY #1	F	19	UP		
PATIENT3,BEN	000003	COMMUNITY #1	F	22	UP		
PATIENT4,STUART	000004	COMMUNITY #1	F	35	UP,AC,PREG	04/10/11	SCREEN,
04/10/11 USER, 04/10/11		SMOKELESS					
PATIENT5,HARRY B	000005	COMMUNITY #1	M	13	UP	03/15/11	SCREEN
PATIENT6,EMERSON	000006	COMMUNITY #1	M	15	UP,AC	05/21/11	SCREEN,
05/21/11 USER, 05/21/11		SMOKER, 05/21/11					
PATIENT7,EUGENE JAY	000007	COMMUNITY #1	M	29	UP		

PATIENT8,ROGER	000008	COMMUNITY #1	M	31	UP,AC	01/21/11	SCREEN,
01/21/11 USER,	01/21/11	SMOKER					
PATIENT9,ANDREW	000009	COMMUNITY #1	M	42	UP		

Figure 2-49: Sample Patient List, Tobacco Use Assessment

2.6.6 Tobacco Cessation

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 23.7% for the proportion of tobacco-using patients who receive tobacco cessation intervention.

Denominators

Active Clinical patients identified as *current tobacco users* prior to the Report Period. Broken down by gender and age groups (<12, 12–17, 18 and older). (GPRA Denominator)

User Population patients identified as *current tobacco users* prior to the report period. Broken down by gender.

Numerators

Patients who have received tobacco cessation counseling or received a prescription for a smoking cessation aid during the Report Period.

Note: This numerator does *not* include refusals. (GPRA Numerator)

Patients who refused tobacco cessation counseling during the Report Period.

Patients identified during the Report Period as having quit their tobacco use.

Patients who have received tobacco cessation counseling, received a prescription for a smoking cessation aid, or who quit their tobacco use during the Report Period.

Note: This numerator does *not* include refusals.

Logic Description

Age is calculated at the beginning of the Report Period.

	ICD and Other Codes
Tobacco Users (documented prior to the Report Period)	<p>Tobacco Health Factors (looks at the last documented health factor in the Tobacco, TOBACCO (SMOKING) and TOBACCO (SMOKELESS–CHEWING/DIP) categories): Current Smoker, Current Smokeless, Current Smoker and Smokeless, Cessation-Smoker, Cessation-Smokeless, Current Smoker, status unknown, Current smoker, every day, or Current smoker, some day</p> <p>CPT code (looks at the last documented): 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, G8455 (old code), G8456 (old code), G8402 (old code) or G8453 (old code)</p> <p>V POV or current Active Problem List (looks at the last documented): 305.1, 305.10-305.12 (old codes), or 649.00-649.04</p>
Tobacco Cessation Counseling (documented during the Report Period)	<p>Patient education codes containing: "TO-", "-TO", "-SHS", 305.1, 305.1* (old codes), 649.00-649.04, D1320, 99406, 99407, G0375 (old code), G0376 (old code), 4000F, G8402 (old code) or G8453 (old code)</p> <p>Dental code: 1320</p> <p>Clinic code: 94 (tobacco cessation clinic)</p> <p>CPT code: D1320, 99406, 99407, G0375 (old code), G0376 (old code), 4000F, G8402 (old code) or G8453 (old code)</p>
Prescription for Tobacco Cessation Aid (documented during the Report Period)	<p>Taxonomy: Medication in the site-populated BGP CMS SMOKING CESSATION MEDS taxonomy</p> <p>Medication Name: Any medication with name containing "NICOTINE PATCH", "NICOTINE POLACRILEX", "NICOTINE INHALER", or "NICOTINE NASAL SPRAY"</p> <p>CPT Code: 4001F</p>
Quit Tobacco User (documented during the Report Period)	<p>V POV or current Active Problem List: 305.13 Tobacco use in remission (old code) or V15.82</p> <p>Health Factors documented during the Report Period (looks at the last documented health factor): Previous Smoker, Previous Smokeless, Previous (Former) Smoker, Previous (Former) Smokeless.</p>

Tobacco Cessation Counseling Refusal Definition

Documented refusal of patient education code containing "TO-", "-TO", or "-SHS" or CPT code D1320, 99406, 99407, G0375 (old code), G0376 (old code), 4000F, G8402 (old code) or G8453 (old code). Refusals will only be counted if a patient did not receive counseling or a prescription for tobacco cessation aid.

Key Logic Changes from CRS Version 11.0

1. For Tobacco Users and Tobacco Cessation Counseling, noted that G8453, G8455, G8456, and G8402 are old codes.

- Added refusal of CPT/ HCPCS codes D1320, 99406, 99407, G0375 (old code), G0376 (old code), 4000F, G8402 (old code) or G8453 (old code) to Tobacco Cessation Counseling refusal definition.

Patient List Description

List of tobacco users with tobacco cessation intervention, if any, or who have quit tobacco use.

Measure Source

Smoking Cessation Attempts: HP 2020 TU-4; Smoking Cessation Counseling: HP 2020 TU-10

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	25.0%
IHS FY 2009 Performance	24.0%
IHS FY 2008 Performance	21.0%
IHS FY 2007 Performance	16.0%
IHS FY 2006 Performance	12.0%
<i>HP 2020 goal for increasing smoking cessation attempts for adult smokers</i>	<i>80.0%</i>

DU	May 25, 2011							Page 101	
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Tobacco Cessation (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical Tobacco Users (GPRA)	301		236			182			
# w/tobacco cessation counseling or Rx for cessation-No Refusals (GPRA)	54	17.9	46	19.5	-1.6	48	26.4	-8.4	
# w/refusal of counseling	8	2.7	0	0.0	+2.7	0	0.0	+2.7	
# who quit	13	4.3	3	1.3	+3.0	1	0.5	+3.8	
# w/ cessation counseling, cessation aid, or quit -No Refusals	66	21.9	48	20.3	+1.6	49	26.9	-5.0	
Male Active Clinical Tobacco Users	140		117			94			

# w/tobacco cessation counseling or RX for cessation-No Refusals	29	20.7	20	17.1	+3.6	25	26.6	-5.9
# w/refusal of counseling	2	1.4	0	0.0	+1.4	0	0.0	+1.4
# who quit	5	3.6	1	0.9	+2.7	1	1.1	+2.5
# w/ cessation counseling, cessation aid, or quit -No Refusals	34	24.3	21	17.9	+6.3	26	27.7	-3.4
Female Active Clinical Tobacco Users	161		119			88		
# w/tobacco cessation counseling or RX for cessation-No Refusals	25	15.5	26	21.8	-6.3	23	26.1	-10.6
# w/refusal of counseling	6	3.7	0	0.0	+3.7	0	0.0	+3.7
# who quit	8	5.0	2	1.7	+3.3	0	0.0	+5.0
# w/ cessation counseling, cessation aid, or quit -No Refusals	32	19.9	27	22.7	-2.8	23	26.1	-6.3

Figure 2-50: Sample Report, Tobacco Cessation

Tobacco Cessation (con't)			
ACTIVE CLINICAL TOBACCO USERS			
Age Distribution			
	<12	12-17	=>18
CURRENT REPORT PERIOD			
Active Clin Tobacco Users	0	6	295
# w/tobacco cessation counseling or Rx for cessation aid -No Refusals	0	0	54
% w/ tobacco cessation counseling or Rx for cessation aid -No Refusals	0.0	0.0	18.3
# w/refusal of counseling	0	0	8
% w/refusal of counseling	0.0	0.0	2.7
# who quit	0	1	12
% who quit	0.0	16.7	4.1
# w/tobacco cessation counseling, or Rx for cessation aid or quit -No Refusals	0	1	65
% w/ tobacco cessation counseling, or Rx for cessation aid or quit -No Refusals	0.0	16.7	22.
PREVIOUS YEAR PERIOD			

Active Clin Tobacco Users	1	4	231
# w/tobacco cessation counseling or Rx for cessation aid			
-No Refusals	0	0	46
% w/tobacco cessation counseling or Rx for cessation aid			
-No Refusals	0.0	0.0	19.9
# w/refusal of counseling	0	0	0
% w/refusal of counseling	0.0	0.0	0.0
# who quit	0	0	3
% who quit	0.0	0.0	1.3
# w/tobacco cessation counseling, Rx for cessation aid or quit			
-No Refusals	0	0	48
% w/ tobacco cessation counseling, Rx for cessation aid or quit			
-No Refusals	0.0	0.0	20.8
CHANGE FROM PREV YR %			
w/tobacco cessation counseling or Rx for cessation aid			
-No Refusals	+0.0	+0.0	-1.6
w/refusal of counseling	+0.0	+0.0	+2.7
who quit	+0.0	+16.7	+2.8
w/tobacco cessation counseling, Rx for cessation aid or quit			
-No Refusals	+0.0	+16.7	+1.3

Figure 2-51: Sample Age Breakdown Report, Tobacco Cessation

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Tobacco Cessation: List of tobacco users with tobacco cessation intervention, if any, or who have quit tobacco use.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,BRITNEY CPT G0375	000001	COMMUNITY #1	F	22	UP;AC	COUNSEL: 06/10/11	
PATIENT2,LORETTA 305.1-DP	000002	COMMUNITY #1	F	22	UP;AC	COUNSEL: 01/13/11	
PATIENT3,HALEY TO-LA	000003	COMMUNITY #1	F	25	UP;AC	COUNSEL: 02/19/11	
PATIENT4,ANGEL CPT 4000F	000004	COMMUNITY #1	F	30	UP;AC	COUNSEL: 03/05/11	
PATIENT5,JOYCE (FORMER) SMOKER 05/31/11	000005	COMMUNITY #1	F	31	UP;AC	QUIT: PREVIOUS	
PATIENT6,ESTHER CESSATION MED - NICOTINE 14MG TRANSDERMAL PATCH	000006	COMMUNITY #1	F	32	UP;AC	COUNSEL: 03/05/11	
PATIENT7,SARAH	000007	COMMUNITY #1	F	33	UP;AC		

PATIENT8, PAULA TO-QT	000008 COMMUNITY #1 F 34 UP;AC	COUNSEL: 03/17/11
--------------------------	--------------------------------	-------------------

Figure 2-52: Sample Patient List Tobacco Cessation

2.7 Behavioral Health Related Performance Measure Topics

2.7.1 Alcohol Screening (FAS Prevention)

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 51.7% for the proportion of female patients ages 15 to 44 who receive screening for alcohol use.

Denominators

Female Active Clinical patients ages 15 to 44. (GPRA Denominator)

Female User Population patients ages 15 to 44.

Numerators

Patients screened for alcohol use, had an alcohol-related diagnosis or procedure, or received alcohol-related patient education during the report period.

Note: This numerator does *not* include refusals. (GPRA Numerator)

- Patients with alcohol screening during the report period.
- Patients with alcohol-related diagnosis or procedure during the report period
- Patients with alcohol-related patient education during the report period.
- Patients with documented alcohol screening refusal in past year.

Logic Description

Ages are calculated at beginning of Report Period.

Alcohol screening definition: Any of the following during the Report Period: (a) Alcohol Screening Exam, any CAGE Health Factor, or Screening Diagnosis; (b) Alcohol-related diagnosis in POV, Current PCC or BHS Problem List; (c) Alcohol-related procedure; or (d) Patient education.

	ICD and Other Codes
Alcohol Screening	PCC Exam Code: 35 CPT code: 99408, 99409, G0396, G0397, H0049, H0050, 3016F Any CAGE Health Factor V POV: V11.3 (history of alcoholism), V79.1 (screening for alcoholism) BHS Problem Code: 29.1 (Screening for Alcoholism) V Measurement in PCC or BHS: AUDT, AUDC, or CRFT
Alcohol-related Diagnosis	V POV, Current PCC or BHS Problem List: 303.*, 305.0*, 291.*, 357.5* BHS POV: 10, 27, 29
Alcohol-related Procedure	V Procedure: 94.46, 94.53, 94.61-94.63, 94.67-94.69
Alcohol-related Education	Patient Education codes: "AOD-" or "-AOD", "CD-" or "-CD" (old codes), or containing V11.3, V79.1, 303.*, 305.0*, 291.* 357.5*, 99408, 99409, G0396, G0397, H0049, or H0050, or 3016F

Alcohol screening may be documented with either an exam code or the CAGE health factor in PCC or BHS. BHS problem codes can also currently be used.

Refusal definition: Refusal of PCC Exam Code 35 in the past year

Recommended Brief Screening Tool

Single Alcohol Screening Question (SASQ) (below).

For Women:

When was the last time you had more than four drinks in one day?

For Men:

When was the last time you had more than five drinks in one day?

Any time in the past three months is a positive screen; further evaluation indicated. Provider should note the screening tool used was the SASQ in the COMMENT section of the Exam Code.

Alcohol Health Factors

The existing Health Factors for alcohol screening are based on the CAGE questionnaire, which asks the following four questions:

1. Have you ever felt the need to **C**ut down on your drinking?
2. Have people **A**nnoyed you by criticizing your drinking?
3. Have you ever felt bad or **G**uilty about your drinking?

4. Have you ever needed an **E**ye opener the first thing in the morning to steady your nerves or get rid of a hangover?
5. Based on how many YES answers are received, document Health Factor:
 - HF-CAGE 0/4 (all “No” answers)
 - HF-CAGE 1/4
 - HF-CAGE 2/4
 - HF-CAGE 3/4
 - HF-CAGE 4/4

Optional values:

- Level/Severity: Mild, Moderate, or Severe
- Quantity: # of drinks daily

Key Logic Changes from CRS Version 11.0

1. Added CPT code 3016F to Alcohol Screening definition.
2. Updated patient list.

Patient List Description

List of female patients with documented alcohol screening or refusal, if any.

Measure Source

HP 2010 16–17a

Measure Past Performance and Long-term Targets

Performance	Percent
IHS FY 2010 Performance	55.0%
IHS FY 2009 Performance	52.0%
IHS FY 2008 Performance	47.0%
IHS FY 2007 Performance	41.0%
IHS FY 2006 Performance	28.0%
IHS FY 2005 Performance	11.0%
IHS FY 2004 Performance	7.0%

DU	May 25, 2011	Page 114
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Alcohol Screening (FAS Prevention)									
	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %	
Female Active Clinical ages 15-44 (GPRA)	434		348			304			
# w/ alcohol screening/Dx/Proc/Pt Ed									
-No Refusals (GPRA)	50	11.5	2	0.6	+10.9	1	0.3	+11.2	
A. # w/alcohol screening	43	9.9	1	0.3	+9.6	0	0.0	+9.9	
B. # w/alcohol related Dx or procedure	3	0.7	1	0.3	+0.4	1	0.3	+0.4	
C. # w/alcohol related patient education	10	2.3	0	0.0	+2.3	0	0.0	+2.3	
# w/alcohol screening refusal	3	0.7	0	0.0	+0.7	0	0.0	+0.7	
Female User Population ages 15-44	725		636			588			
# w/ alcohol screening/Dx/Proc/Pt Ed									
-No Refusals	54	7.4	2	0.3	+7.1	2	0.3	+7.1	
A. # w/alcohol screening	46	6.3	1	0.2	+6.2	0	0.0	+6.3	
B. # w/alcohol related Dx or procedure	4	0.6	1	0.2	+0.4	2	0.3	+0.2	
C. # w/alcohol related patient education	11	1.5	0	0.0	+1.5	0	0.0	+1.5	
# w/alcohol screening refusal	3	0.4	0	0.0	+0.4	0	0.0	+0.4	

Figure 2-53: Sample Report, Alcohol Screening (FAS Prevention)

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Alcohol Screening (FAS Prevention): List of female patients with documented alcohol screening or refusal, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,CHRISTINE S	000001	COMMUNITY #1	F	15	UP;	
PATIENT2,RITA A	000002	COMMUNITY #1	F	15	UP;AC	SCREEN: 03/06/03
POV V11.3						
PATIENT3,DIANE L	000003	COMMUNITY #1	F	15	UP;	Refused: 02/02/03
exam 35						
PATIENT4,ALICIA	000004	COMMUNITY #1	F	15	UP;AC	
PATIENT5,MELISSA	000005	COMMUNITY #1	F	16	UP;AC	PT ED: 02/13/03
99408-P						
PATIENT6,LISA MARIE	000006	COMMUNITY #1	F	16	UP;AC	SCREEN: 10/13/03
HF: CAGE 1/4						
PATIENT7,RUTH NELLIE	000007	COMMUNITY #1	F	16	UP;	

PATIENT8,ALISHA DAWN CPT 3016F	000008 COMMUNITY #1 F 16 UP;AC	SCREEN: 03/03/03
-----------------------------------	-----------------------------------	------------------

Figure 2-54: Sample Patient List, Alcohol Screening (FAS Prevention)

2.7.2 Alcohol Screening and Brief Intervention (ASBI) in the ER

Denominators

Number of visits for *Active Clinical patients* age 15–34 seen in the ER for injury during the report period. Broken down by gender and age groups (15–24 and 25–34).

Number of visits for *Active Clinical patients* age 15–34 seen in the ER for injury and screened positive for hazardous alcohol use during the report period. Broken down by gender and age groups (15–24 and 25–34).

Number of visits for *User Population patients* age 15–34 seen in the ER for injury during the report period. Broken down by gender and age groups (15–24 and 25–34).

Number of visits for *User Population patients* age 15–34 seen in the ER for injury and screened positive for hazardous alcohol use during the report period. Broken down by gender and age groups (15–24 and 25–34).

Numerators

Number of visits where patients were screened in the ER for hazardous alcohol use.

- a. Number of visits where patients were screened positive.

Number of visits where patients were provided a brief negotiated interview (BNI) at or within seven days of the ER visit.

- a. Number of visits where patients were provided a BNI at the ER visit.
- b. Number of visits where patients were provided a BNI not at the ER visit but within 7 days of the ER visit.

Logic Description

Age of the patient is calculated as of the beginning of the Report Period.

Emergency room visit definition: Clinic Code 30.

Multiple visits definition: If a patient has multiple ER visits for injury during the Report Period, each visit will be counted in the denominator. For the screening numerator, each ER visit with injury at which the patient was screened for hazardous alcohol use will be counted. For the positive alcohol use screen numerator, each ER visit with injury at which the patient screened positive for hazardous alcohol use will be counted. For the BNI numerators, each visit where the patient was either provided a BNI at the ER or within seven days of the ER visit will be counted. An example of this logic is shown below.

ER Visit w/Injury	Denom Count	Scrn Num	Pos Scrn Num Count	BNI Num Count

John Doe, 07/17/09, Screened Positive at ER, BNI at ER				
John Doe, 09/01/09, Screened Positive at ER, No BNI				
John Doe, 11/15/09, No Screen				

COUNTS:	3	2	2	1

CRS uses the following codes:

	ICD and Other Codes
Injury	V POV (primary or secondary): 800.0–999.9 or E800.0–E989.
ER Screening for Hazardous Alcohol Use	<i>Any conducted during an ER visit:</i> PCC Exam Code: 35 Any Alcohol Health Factor (i.e., CAGE) V POV: V79.1 Screening for Alcoholism CPT: G0396, G0397, H0049, 99408, 99409, 3016F V Measurement in PCC: AUDT, AUDC, or CRFT
Positive Screen for Hazardous Alcohol Use	<i>Any of the following for the screening conducted during an ER visit:</i> PCC Exam Code: 35 Alcohol Screening result of “Positive” Health Factor: CAGE result of 1/4, 2/4, 3/4 or 4/4 CPT: G0396, G0397, 99408, 99409 V Measurement Result in PCC: AUDT result of => 8, AUDC result of => 4 for men and =>3 for women, CRFT result of 2-6
BNI	<i>Any of the following documented at the ER visit or within seven days of the ER visit at a face-to-face visit, which excludes chart reviews and telecommunication visits:</i> CPT: G0396, G0397, H0050, 99408, 99409 Patient Education Code: AOD-BNI or containing G0396, G0397, H0050, 99408, or 99409

Recommended Brief Screening Tool

SASQ (below).

For Women:

When was the last time you had more than four drinks in one day?

For Men:

When was the last time you had more than five drinks in one day?

Any time in the past three months is a positive screen; further evaluation indicated. Provider should note the screening tool used was the SASQ in the COMMENT section of the Exam Code.

Alcohol Health Factors

The existing Health Factors for alcohol screening are based on the CAGE questionnaire, which asks the following four questions:

1. Have you ever felt the need to **C**ut down on your drinking?
2. Have people **A**nnoyed you by criticizing your drinking?
3. Have you ever felt bad or **G**uilty about your drinking?
4. Have you ever needed an **E**ye opener the first thing in the morning to steady your nerves or get rid of a hangover?

Based on how many YES answers are received, document Health Factor:

- HF-CAGE 0/4 (all No answers)
- HF-CAGE 1/4
- HF-CAGE 2/4
- HF-CAGE 3/4
- HF-CAGE 4/4

Optional values:

- Level/Severity: Mild, Moderate, or Severe
- Quantity: # of drinks daily

Key Logic Changes from CRS Version 11.0

1. Added CPT code 3016F to ER Screening for Hazardous Alcohol Use definition.
2. Updated patient list.

Patient List Description

List of visits for patients seen in the ER for an injury, with screening for hazardous alcohol use, results of screen and BNI, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 117	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Alcohol Screening and Brief Intervention (ASBI) in the ER								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
# ER Injury Visits for AC Pts 15-34	33		33			32		
# Visits w/ ER Hazardous Alcohol Screening	19	57.6	0	0.0	+57.6	0	0.0	+57.6
A. # Visits w/Positive Screen	15	45.5	0	0.0	+45.5	0	0.0	+45.5
# ER Injury Visits for Male AC Pts 15-34	12		18			20		
# Visits w/ ER Hazardous Alcohol Screening	7	58.3	0	0.0	+58.3	0	0.0	+58.3
A. # Visits w/Positive Screen	5	41.7	0	0.0	+41.7	0	0.0	+41.7
# ER Injury Visits for Female AC Pts 15-34	21		15			12		
# Visits w/ ER Hazardous Alcohol Screening	12	57.1	0	0.0	+57.1	0	0.0	+57.1
A. # Visits w/Positive Screen	10	47.6	0	0.0	+47.6	0	0.0	+47.6
# of ER Injury Visits for AC Pts 15-24	17		16			21		
# Visits w/ ER Hazardous Alcohol Screening	10	58.8	0	0.0	+58.8	0	0.0	+58.8
A. # Visits w/Positive								

Screen	9	52.9	0	0.0	+52.9	0	0.0	+52.9
# ER Injury Visit for AC Pts 25-34	16		17			11		
# Visits w/ ER Hazardous Alcohol Screening	9	56.3	0	0.0	+56.3	0	0.0	+56.3
A. # Visits w/Positive Screen	6	37.5	0	0.0	+37.5	0	0.0	+37.5

Figure 2-55: Sample Report, Alcohol Screening and Brief Intervention (ASBI) in the ER

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Alcohol Screening and Brief Intervention (ASBI) in the ER: List of visits
for patients seen in the ER for an injury, with screening for hazardous
alcohol use, results of screen and BNI, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,DARLENE S	000001	COMMUNITY #1	F	33	UP;AC	ER 1) 02/02/11POV
816.02, SCREEN: Neg/No Res screening CPT 3016F						
PATIENT2,RITA A	000002	COMMUNITY #1	F	33	UP;AC	ER 1) 07/12/11POV
875.0, SCREEN: Pos EXAM 35, BNI: No						
PATIENT3,DIANE L	000003	COMMUNITY #1	F	15	UP;	ER 1) 09/08/11POV
815.00, SCREEN: None						
PATIENT4,ALICIA	000004	COMMUNITY #1	F	18	UP;AC	ER 1) 04/20/11POV
959.7, SCREEN: Neg/No Res screening CPT H0049						
PATIENT5,MELISSA	000005	COMMUNITY #1	F	16	UP;AC	
PATIENT6,LISA MARIE	000006	COMMUNITY #1	F	20	UP;AC	ER 1) 05/01/11POV
959.01, SCREEN: None; ER 2) 07/01/11POV 999.9, SCREEN: Pos EXAM 35, BNI: 07/01/11 At ER CPT-H0050						
PATIENT7,RUTH NELLIE	000007	COMMUNITY #1	F	16	UP;	ER 1) 12/10/11POV
920., SCREEN: None; ER 2) 12/11/11POV 851.80, SCREEN: None						

Figure 2-56: Sample Patient List, Alcohol Screening and Brief Intervention (ASBI) in the ER

2.7.3 Intimate Partner (Domestic) Violence Screening

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 52.8% for the proportion of female patients ages 15 to 40 who receive screening for domestic violence.

Denominators

Female Active Clinical patients ages 13 and older.

Female Active Clinical patients ages 15–40. (GPRA Denominator)

Female User Population patients ages 13 and older.

Numerators

Patients screened for intimate partner (domestic) violence at any time during the Report Period.

Note: This numerator does *not* include refusals. (GPRA Numerator)

- a. Patients with documented IPV/DV exam
- b. Patients with IPV/DV related diagnosis
- c. Patients provided with education or counseling about IPV/DV
- d. Patients with documented refusal in past year of an IPV/DV exam or IPV/DV-related education.

Logic Description

Age of the patient is calculated at the beginning of the report period. CRS uses the following codes to define numerators.

	CPT Codes	ICD and Other Codes
IPV/DV Screening		V Exam: Code 34 BHS Exam: IPV/DV
IPV/DV Diagnosis		V POV or current PCC or BHS Problem List: 995.80-995.83, 995.85, V15.41, V15.42, V15.49 BHS POV: 43.*, 44.*
IPV/DV Education		Patient education codes: "DV-" or "-DV", 995.80-83, 995.85, V15.41, V15.42, or V15.49
IPV/DV Counseling		V POV: V61.11
Refusals		V Exam: Code 34 BHS IPV/DV exam Patient education codes containing "DV-" or "-DV"

Key Logic Changes from CRS Version 11.0

1. Updated patient list.

Patient List Description

List of female patients 13 and older with documented IPV/DV screening or refusal, if any.

Measure Source

HP 2010 15–34

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	53.0%
IHS FY 2009 Performance	48.0%
IHS FY 2008 Performance	42.0%
IHS FY 2007 Performance	36.0%
IHS FY 2006 Performance	28.0%
IHS FY 2005 Performance	13.0%
IHS FY 2004 Performance	4.0%

DU	May 25, 2011						Page 128		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Intimate Partner (Domestic) Violence Screening (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
# Female Active									
Clinical ages									
13 and older	712		519			460			
# w/IPV/DV									
Screening									
-No Refusals	13	1.8	1	0.2	+1.6	0	0.0	+1.8	
A. # w/documented									
IPV/DV exam	8	1.1	0	0.0	+1.1	0	0.0	+1.1	
B. # w/ IPV/DV related									
diagnosis	4	0.6	0	0.0	+0.6	0	0.0	+0.6	
C. # provided DV									
education	4	0.6	1	0.2	+0.4	0	0.0	+0.6	
# w/ IPV/DV									
refusal	3	0.4	0	0.0	+0.4	0	0.0	+0.4	
# Female Active									
Clinical ages 15-40									
(GPRA)	380		311			267			
# w/IPV/DV screening									
-No Refusals									
(GPRA)	11	2.9	1	0.3	+2.6	0	0.0	+2.9	
A. # w/ documented									
IPV/DV exam	6	1.6	0	0.0	+1.6	0	0.0	+1.6	
B. # w/ IPV/DV related									
diagnosis	3	0.8	0	0.0	+0.8	0	0.0	+0.8	
C. # provided DV									
education	4	1.1	1	0.3	+0.7	0	0.0	+1.1	
# w/ IPV/DV									
refusal	2	0.5	0	0.0	+0.5	0	0.0	+0.5	

# Female User Pop 13 and older	1,218		1,005			917		
# w/IPV/DV Screening								
-No Refusals	13	1.1	1	0.1	+1.0	1	0.1	+1.0
A. # w/ documented IPV/DV exam	8	0.7	0	0.0	+0.7	0	0.0	+0.7
B. # w/ IPV/DV related diagnosis	4	0.3	0	0.0	+0.3	1	0.1	+0.2
C. # provided DV education	4	0.3	1	0.1	+0.2	0	0.0	+0.3
# w/ IPV/DV refusal	3	0.2	0	0.0	+0.2	0	0.0	+0.2

Figure 2-57: Sample Report, Intimate Partner (Domestic) Violence Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Intimate Partner (Domestic) Violence Screening: List of female patients 13 and older with documented IPV/DV screening or refusal, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,ELVIRA	000001	COMMUNITY #1	F	13	UP;	03/18/11 Refused Ex	
34							
PATIENT2,SHARON KAY	000002	COMMUNITY #1	F	14	UP;		
PATIENT3,KRISTINA	000003	COMMUNITY #1	F	15	UP;		
PATIENT4,RITA	000004	COMMUNITY #1	F	15	UP;AC	EXAM: 05/06/11 Ex	
34							
PATIENT5,DIANE LOUISE	000005	COMMUNITY #1	F	15	UP;	EXAM: 02/24/11 Ex	
34							
PATIENT6ALICE LILA	000006	COMMUNITY #1	F	15	UP;AC		

Figure 2-58: Sample Patient List, Intimate Partner (Domestic) Violence Screening

2.7.4 Depression Screening

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 51.9% for the proportion of adults ages 18 and older who receive annual screening for depression.

Denominators

Active Clinical patients ages 18 and older. Broken down by gender. (GPRA Denominator)

- a. *Active Clinical patients ages 65 and older.* Broken down by gender.

User Population patients ages 18 and older. Broken down by gender.

- a. *User Population patients ages 65 and older.* Broken down by gender.

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever. Broken down by gender.

Active IHD patients, defined as all Active Clinical patients diagnosed with ischemic heart disease (IHD) prior to the Report Period, *and* at least two visits during the Report Period, *and* two IHD-related visits ever. Broken down by gender.

Numerators

Patients screened for depression or diagnosed with a mood disorder at any time during the Report Period.

Note: This numerator does not include refusals. (GPRA Numerator)

- a. Patients screened for depression during the Report Period.
- b. Patients with a diagnosis of a mood disorder during the Report Period.
- c. Patients with documented depression screening refusal in past year.
- d. Patients with depression-related education or refusal of education in past year.

Note: Depression-related patient education does not count toward the GPRA numerator and is included as a separate numerator only.

Logic Description

Age is calculated at beginning of the Report Period.

CRS uses the following codes and taxonomies to define the denominator and numerators.

	ICD and Other Codes
Diabetes	V POV: 250.00–250.93
Ischemic Heart Disease	V POV: 410.0–412.*, 414.0–414.9, 429.2
Depression Screening	V Exam: Exam Code 36 V POV: V79.0 CPT: 1220F BHS Problem Code: 14.1 (Screening for Depression) V Measurement in PCC or BHS: PHQ2 or PHQ9
Mood Disorders	At least 2 visits in PCC or BHS for: Major Depressive Disorder, Dysthymic Disorder, Depressive Disorder NOS, Bipolar I or II Disorder, Cyclothymic Disorder, Bipolar Disorder NOS, Mood Disorder Due to a General Medical Condition, Substance-induced Mood Disorder, or Mood Disorder NOS. V POV: 296.*, 291.89, 292.84, 293.83, 300.4, 301.13, or 311 BHS POV: 14, 15

	ICD and Other Codes
Depression-related Patient Education (does not count toward GPRA numerator)	<i>Documented education of any of the following during the Report Period:</i> Patient education codes: containing "DEP-" (depression), 296.2* or 296.3*, "BH-" (behavioral and social health), 290319, 995.5*, or 995.80–995.85, "SB-" (suicidal behavior) or 300.9, or "PDEP-" (postpartum depression) or 648.44.
Screening Refusals	V Exam: Exam Code 36, in past year
Refusal of Depression-related Patient Education (does not count toward GPRA numerator)	<i>Documented refusal of any of the following during the Report Period:</i> Patient education codes: containing "DEP-" (depression), "BH-" (behavioral and social health), "SB-" (suicidal behavior), or "PDEP-" (postpartum depression).

Recommended Brief Screening Tool

A sample of a Patient Health Questionnaire (PHQ-2 Scaled Version) appears below.

Over the past two weeks, how often have you been bothered by any of the following problems?

1. Little interest or pleasure in doing things

- Not at all Value: 0
- Several days Value: 1
- More than half the days Value: 2
- Nearly every day Value: 3

2. Feeling down, depressed, or hopeless

- Not at all Value: 0
- Several days Value: 1
- More than half the days Value: 2
- Nearly every day Value: 3

Total Possible PHQ-2 Score: Range: 0–6

0–2: Negative

3–6: Positive; further evaluation indicated

Provider should note the screening tool used was the PHQ-2 Scaled in the COMMENT section of the Exam Code.

Key Logic Changes from CRS Version 11.0

1. Removed CPT code range 428.* from Ischemic Heart Disease definition.

2. Added CPT code 1220F to Depression Screening definition.
3. Updated patient list.

Patient List Description

List of patients with documented depression screening or refusal/diagnosed with mood disorder, if any.

Measure Source

USPSTF (US Preventive Services Task Force), HP 2010 developmental indicator 18–6.

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	52.0%
IHS FY 2009 Performance	44.0%
IHS FY 2008 Performance	35.0%
IHS FY 2007 Performance	24.0%
IHS FY 2006 Performance	15.0%

DU	May 25, 2011						Page 130	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Depression Screening (con't)								
	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
Active Clinical Pts =>18 (GPRA)	1,112		793			666		
# w/Depression screening or Mood Disorder DX-No Refusals (GPRA)	81	7.3	42	5.3	+2.0	17	2.6	+4.7
A. # screened for depression	39	3.5	0	0.0	+3.5	0	0.0	+3.5
B. # w/mood disorder DX	43	3.9	42	5.3	-1.4	17	2.6	+1.3
# w/depression screening refusal	5	0.4	0	0.0	+0.4	0	0.0	+0.4
# w/depression education or refusal	11	1.0	3	0.4	+0.6	0	0.0	+1.0
Male Active Clinical Pts >=18	448		308			250		

# w/ Depression screening or Mood Disorder DX-No Refusals	25	5.6	7	2.3	+3.3	1	0.4	+5.2
A. # screened for depression	14	3.1	0	0.0	+3.1	0	0.0	+3.1
B. # w/Mood Disorder DX	11	2.5	7	2.3	+0.2	1	0.4	+2.1
# with depression screening refusal	3	0.7	0	0.0	+0.7	0	0.0	+0.7
# w/depression education or refusal	3	0.7	1	0.3	+0.3	0	0.0	+0.7
Female Active Clinical Pts >=18	664		485			416		
# w/ Depression screening or Mood Disorder DX-No Refusals	56	8.4	35	7.2	+1.2	16	3.8	+4.6
A. # screened for depression	25	3.8	0	0.0	+3.8	0	0.0	+3.8
B. # w/Mood Disorder DX	32	4.8	35	7.2	-2.4	16	3.8	+1.0
# with depression screening refusal	2	0.3	0	0.0	+0.3	0	0.0	+0.3
# w/depression education or refusal	8	1.2	2	0.4	+0.8	0	0.0	+1.2
A. Active Clinical Pts => 65	116		64			65		
# w/ Depression screening or Mood Disorder DX-No Refusals	9	7.8	6	9.4	-1.6	2	3.1	+4.7
A. # screened for depression	3	2.6	0	0.0	+2.6	0	0.0	+2.6
B. # w/mood disorder DX	6	5.2	6	9.4	-4.2	2	3.1	+2.1
# with depression screening refusal	1	0.9	0	0.0	+0.9	0	0.0	+0.9
# w/depression education or refusal	2	1.7	2	3.1	-1.4	0	0.0	+1.7

Figure 2-59: Sample Report, Depression Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Depression Screening: List of patients with documented depression screening or refusal/diagnosed with mood disorder, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT55, LORETTA LYNN	000055	COMMUNITY #1	F	78	UP		
PATIENT56, TINA MARIE	000056	COMMUNITY #1	F	78	UP;AC;AD;IHD	SCREEN: 05/22/0	
PATIENT57, DANIELLE	000057	COMMUNITY #1	F	79	UP;AC	PT ED: 02/06/11	
296.20-DP							
PATIENT58, LESLIE ANN	000058	COMMUNITY #1	F	80	UP;AC	SCREEN: 04/15/11	
POV V79.0							

PATIENT59,DONNA SUE	000059	COMMUNITY #1	F	86	UP;AC	SCREEN: 01/15/11
POV V79.0						
PATIENT60,TAYLOR OLIVIA	000060	COMMUNITY #1	F	87	UP;AC	
PATIENT61,DENNIS GERALD	000061	COMMUNITY #1	M	18	UP	EDUC: 296.20-
DP:02/01/11						
PATIENT62,JOSHUA DALE	000062	COMMUNITY #1	M	18	UP;AC	

Figure 2-60: Sample Patient List, Depression Screening

2.7.5 Antidepressant Medication Management

Denominators

As of the 120th day of the Report Period, *Active Clinical* patients 18 years of age and older who were diagnosed with a new episode of depression and treated with antidepressant medication in the past year.

As of the 120th day of the Report Period, *User Population* patients 18 years of age and older who were diagnosed with a new episode of depression and treated with antidepressant medication in the past year.

Numerators

Optimal Practitioner Contacts: Patients with at least three mental health visits with a non-mental health or mental health provider within 12 weeks (84 days) after diagnosis, 2 of which must be face-to-face visits and one of which must be with a prescribing provider.

Effective Acute Phase Treatment: Patients who filled a sufficient number of separate prescriptions/refills of antidepressant medication for continuous treatment of at least 84 days (12 weeks).

Effective Continuation Phase Treatment: Patients who filled a sufficient number of separate prescriptions/refills of antidepressant medication treatment to provide continuous treatment for at least 180 days (6 months).

Logic Description

Age is calculated at the beginning of the report period. To be included in the denominator, patient must meet *both* of the following conditions:

1. One of the following from the 121st day of the year prior to the Report Period to the 120th day of the Report Period:
 - a. One visit in any setting with major depression DX (see list of codes below) as primary POV
 - b. Two outpatient visits occurring on different dates of service with secondary POV of major depression

c. An inpatient visit with secondary POV of major depression.

For example, if Report Period is July 1, 2010–June 30, 2011, the patient must have one of the three scenarios above during 11/1/2009–10/29/2010.

Major depression is defined as POV 296.2*, 296.3*, 298.0, 300.4, 309.1, 311. The Index Episode Start Date is date of the patient's earliest visit during this period. For inpatient visits, the discharge date will be used.

2. Filled a prescription for an antidepressant medication (see list of medications below) within 30 days before the Index Episode Start Date or 14 days on or after that date. In V Medication, Date Discontinued must not be equal to the prescription (i.e., visit) date. The Index Prescription Date is the date of the earliest prescription for antidepressant medication filled during that time period.

Denominator Exclusions

1. Patients who have had any diagnosis of depression within the previous 120 days (four months) of the Index Episode Start Date. The POVs to be checked for prior depressive episodes is more comprehensive and include the following: POV 296.2*–296.9*, 298.0, 300.4, 309.0, 309.1, 309.28, 311, or
2. Patients who had a new or refill prescription for antidepressant medication (see the list of medications below) within 90 days (3 months) prior to the Index Prescription Date are excluded as they do not represent new treatment episodes, or
3. Patients who had an acute mental health or substance abuse inpatient stay during the 245 days after the Index Episode Start Date treatment period. Acute mental health stays are defined as Service Category of H and primary POV 290*, 293*–302*, 306*–316*. Substance abuse inpatient stays are defined as Service Category of H and primary POV 291*–292*, 303*–305* or primary POV 960*–979* and secondary POV of 291*–292*, 303*–305*.

Optimal Practitioner Contacts Numerator

Patient must have one of the following:

1. Three face-to-face follow-up outpatient, non-ER visits (clinic code not equal to 30) or intermediate treatment with either a non-mental-health or mental health provider within 84 days after the Index Episode Start Date, *or*...
2. Two face-to-face outpatient, non-ER visits (clinic code not equal to 30) and one telephone visit (Service Category T), with either a non-mental-health or mental health provider within 84 days after the Index Episode Start Date.

For either option, one of the visits must be to a prescribing provider, defined as Provider Codes 00, 08, 11, 16-18, 21, 24-25, 30, 33, 41, 44-45, 47, 49, 64, 67-68, 70-83, 85-86, A1, A9, or B1-B6.

Note: If patient was diagnosed with two secondary diagnoses of depression, the second visit may be counted toward the numerator.

Outpatient mental health provider visits are defined as BHS or PCC visit with primary provider code of 06, 12, 19, 48, 49, 50, 62, 63, 81, or 92–96, *and*:

- a. (1) Service category A, S, or O, and (2A) CPT 90801, 90802, 90804–90819, 90821–90824, 90826–90829, 90845, 90847, 90849, 90853, 90857, 90862, 90870, 90871 (old code), 90875, 90876, 99384–99387, 99394–99397, 99401–99404, G0155, G0176, G0177, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013–H2020, M0064, S9484, S9485 or (2B) POV 290*, 293*–302*, 306*–316, or
- b. (1) Service category of A, S, or O and (2A) Location of Encounter = Home (as designated in Site Parameters) or (2B) Clinic Code = 11, or
- c. Service category of T.

Outpatient non-mental-health provider visits are defined as BHS or PCC visits with:

- a. (1) Service category A, S, or O, a (2) CPT 90801, 90802, 90804–90819, 90821–90824, 90826–90829, 90845, 90847, 90849, 90853, 90857, 90862, 90870, 90871 (old code), 90875, 90876, G0155, G0176, G0177, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013–H2020, M0064, S9484, S9485, or...
- b. (1) Service category A, S, O, or T or (2) Location of Encounter = Home (as designated in Site Parameters) or (3A) clinic code 11 and (3B) POV 290*, 293*–302*, 306*–316*, or...
- c. (1) Service category A, S, or O, and (2) CPT 99384–99387, 99394–99397, 99401–99404 and (3) POV 290*, 293*–302*, 306*–316*.

Effective Acute Phase Treatment Numerator

For all antidepressant medication prescriptions filled (see list of medications below) within 114 days of the Index Prescription Date, from V Medication CRS counts the days prescribed (i.e., treatment days) from the Index Prescription Date until a total of 84 treatment days has been established. If the patient had a total gap exceeding 30 days or if the patient does not have 84 treatment days within the 114-day time frame, the patient is not included in the numerator.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2010, Discontinued Date=11/19/2010, Recalculated # Days Prescribed=4.

Example of Patient Included in Numerator:

- First RX is Index Rx Date: 11/1/2010, # Days Prescribed=30
- Rx covers patient through 12/1/2010
- Second RX: 12/15/2010, # Days Prescribed=30
- Gap #1 = (12/15/2010-12/1/2010) = 14 days
- Rx covers patient through 1/14/2011
- Third RX: 1/10/2011, # Days Prescribed=30
- No gap days
- Rx covers patient through 2/13/2011
- Index Rx Date 11/1/2010 + 114 days = 2/23/2011
- Patient's 84th treatment day occurs on 2/7/2011, which is \leq 2/23/2011 and # gap days of 14 is less than 30

Example of Patient Not Included in Numerator:

- First Rx is Index Rx Date: 11/1/2010, # Days Prescribed=30
- Rx covers patient through 12/1/2010
- Second Rx: 12/15/2010, # Days Prescribed=30
- Gap #1 = (12/15/2010-12/1/2010) = 14 days
- Rx covers patient through 1/14/2011
- Third Rx: 2/01/2011, # Days Prescribed=30
- Gap #2 = (2/01/2011-1/14/2011) = 18, total # gap days = 32, so patient is not included in the numerator

Effective Continuation Phase Treatment Numerator

For all antidepressant medication prescriptions (see list of medications below) filled within 231 days of the Index Prescription Date, CRS counts the days prescribed (i.e., treatment days) (from V Medication) from the Index Prescription Date until a total of 180 treatment days has been established. If the patient had a total gap exceeding 51 days or if the patient does not have 180 treatment days within the 231 day time frame, the patient is not included in the numerator.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2010, Discontinued Date=11/19/2010, Recalculated # Days Prescribed=4.

Antidepressant medications defined with medication taxonomy BGP HEDIS ANTIDEPRESSANT MEDS. (Medications are: Tricyclic antidepressants (TCA) and other cyclic antidepressants, Selective serotonin reuptake inhibitors (SSRI), Monoamine oxidase inhibitors (MAOI), Serotonin-norepinephrine reuptake inhibitors (SNRI), and other antidepressants.) Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with new depression DX and optimal practitioner contact (OPC), acute phase treatment (APT) and continuation phase treatment (CONPT), if any.

Measure Source

HEDIS, HP 2010 18-9b

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 141

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Antidepressant Medication Management (con't)

	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts =>18 w/new depression DX and antidepressant meds	17		6			2		
# w/3 outpt mental health visits within 12 weeks	5	29.4	1	16.7	+12.7	0	0.0	+29.4
# w/12 week treatment meds	9	52.9	4	66.7	-13.7	0	0.0	+52.9
# w/180 day treatment								

meds	4	23.5	3	50.0	-26.5	0	0.0	+23.5
User Pop Pts =>18 w/new depression DX and antidepressant meds	18		7			3		
# w/3 outpt mental health visits within 12 weeks	5	27.8	1	14.3	+13.5	0	0.0	+27.8
# w/12 week treatment meds	9	50.0	4	57.1	-7.1	0	0.0	+50.0
# w/180 day treatment meds	4	22.2	3	42.9	-20.6	0	0.0	+22.2

Figure 2-61: Sample Report, Antidepressant Medication Management

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Antidepressant Medication Management: List of patients with new depression DX and optimal practitioner contact (OPC), acute phase treatment (APT) and continuation phase treatment (CONPT), if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,MICHELLE D	000001	COMMUNITY #1	F	22	UP,AC IESD: 06/06/11	NOT OPC;NOT APT: 06/07/11(30);07/08/11(30) ;DAYS=60; GAP=1;NOT CONPT: 06/07/11(30);07/08/11(30) ;DAYS=60; GAP=1
PATIENT2,PAULA KAY	000002	COMMUNITY #1	F	34	UP,AC IESD: 02/05/11	NOT OPC; APT;CONPT
PATIENT3,RHONDA SUE	000003	COMMUNITY #1	F	35	UP,AC IESD: 08/09/11	PC; APT;NOT CONPT: 04/01/11(100);05/23/11(24);05/23/11(24);06/16/11(21);06/16/11(30);07/07/11(30);07/25/11(23);07/25/11(23) ;DAYS=298; GAP=111
PATIENT4,KATHLEEN	000004	COMMUNITY #1	F	38	UP,AC IESD: 01/29/11	NOT OPC;NOT APT: 01/16/11(7);01/16/11(6);01/29/11(20);03/25/11(35) ;DAYS=68; GAP=28;CONPT

Figure 2-62: Sample Patient List, Antidepressant Medication Management

2.8 Cardiovascular Disease Related Measure Topics

2.8.1 Obesity Assessment

Denominators

Active Clinical patients ages 2 through 74. Broken down by gender and age groups (2-5, 6-11, 12-19, 20-24, 25-34, 35-44, 45-54, 55-74).

All User Population patients ages 2 through 74. Broken down by gender.

Numerators

Patients for whom a Body Mass Index (BMI) could be calculated.

Note: This numerator does *not* include refusals.

1. For those with a BMI calculated, those considered overweight but not obese using BMI and standard tables
2. For those with a BMI calculated, those considered obese using BMI and standard tables
3. Total of overweight and obese

Patients with documented refusal in past year.

Logic Description

Age is calculated at beginning of the Report Period.

BMI calculation definition: CRS calculates BMI at the time the report is run, using NHANES II. For age 18 and under, a height and weight must be taken on the same day any time during the Report Period. For 19 through 50, height and weight must be recorded within last 5 years, not required to be on the same day. For over 50, height and weight within last 2 years, not required to be recorded on same day. Overweight but not obese is defined as BMI of 25 through 29 for adults 19 and older. Obese is defined as BMI of 30 or more for adults 19 and older. For ages 2-18, definitions are based on standard tables. Refusals include REF, NMI, and UAS (unable to screen) and must be documented during the past year. For ages 18 and under, both the height and weight must be refused on the same visit at any time during the past year. For ages 19 and older, the height and weight must be refused during the past year and are not required to be on the same visit.

Patients whose BMI either is greater or less than the Data Check Limit range shown in the BMI Standard Reference Data Table in PCC will not be included in the report counts for Overweight or Obese.

Key Logic Changes from CRS Version 11.0

1. Updated patient list.

Patient List Description

List of patients with current BMI, if any.

Measure Source

HP 2020: NWS-9 Obesity in Adults 20+, NWS-10.1 (Obesity in Children 2-5), NWS-10.2 Overweight or Obesity in Children 6-11, NWS-10.3 Overweight or Obesity in Adolescents 12-19, NWS-10.4 Overweight or Obesity in Children 2-19

Measure Past Performance and Long-Term Targets

Performance	Percent
Assessed as Obese—IHS FY 2008 Performance	46.0%
BMI Measured—IHS FY 2008 Performance	74.0%
BMI Measured— FY 2005 Performance	64.0%
BMI Measured—IHS FY 2004 Performance	60.0%
<i>HP 2020 Goal: Obesity in Adults 20+ (NWS-9)</i>	<i>30.6%</i>
<i>HP 2020 Goal: Overweight or Obesity in Children 2–5 (NWS-10.1)</i>	<i>9.6%</i>
<i>HP 2020 Goal: Overweight or Obesity in Children 6–11 (NWS-10.2)</i>	<i>15.7%</i>
<i>HP 2020 Goal: Overweight or Obesity in Adolescents 12–19 (NWS-10.3)</i>	<i>16.1%</i>
<i>HP 2020 Goal: Overweight or Obesity in Children 2–19 (NWS-10.4)</i>	<i>14.6%</i>

Performance Improvement Tips

1. A Body Mass Index report can be run from your PCC Management Reports menu. This report can be run for all patients or for a specific template of patients that has been pre-defined with a QMan search. The BMI report will provide you with patient height, weight, date weight taken, BMI and NHANES percentile.
2. Recent guidelines indicate that height for adults must be taken at least once every five years, rather than once after age 18. Your BMI rates may be lower than anticipated because of height data that is over five years old.
3. If height and weight measurements are being recorded as cm/kg vs. in/lbs ensure providers are *noting* they are cm/kg *and* that data entry is entering the measurements correctly in PCC, as shown below.
 - Use mnemonics of CHT and KWT (vs. HT and WT), or
 - Add “c” after height value and “k” after weight value (e.g. 100c, 50k)

DU	May 25, 2011	Page 143
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Obesity Assessment (con't)		
REPORT	%	PREV YR %
		CHG from BASE %
		CHG from

	PERIOD		PERIOD		PREV YR %		PERIOD		BASE %
Active Clinical Pts ages 2-74	1,400		1,096				982		
# w/BMI calculated									
-No Refusals	880	62.9	824	75.2	-12.3	712	72.5	-9.6	
A. # Overweight w/ % of Total BMI	242	27.5	237	28.8	-1.3	191	26.8	+0.7	
B. # Obese w/ % of Total BMI	372	42.3	339	41.1	+1.1	267	37.5	+4.8	
C. # Overweight/Obese w/ % of Total BMI	614	69.8	576	69.9	-0.1	458	64.3	+5.4	
# w/BMI refusal (No BMI)	4	0.3	0	0.0	+0.3	0	0.0	+0.3	
Male Active Clinical Pts 2-74	595		465				410		
# w/BMI calculated									
-No Refusals	341	57.3	331	71.2	-13.9	283	69.0	-11.7	
A. # Overweight w/ % of Total BMI	103	30.2	98	29.6	+0.6	73	25.8	+4.4	
B. # Obese w/ % of Total BMI	155	45.5	141	42.6	+2.9	117	41.3	+4.1	
C. #Overweight/Obese w/ % of Total BMI	258	75.7	239	72.2	+3.5	190	67.1	+8.5	
# w/BMI refusal (no BMI)	2	0.3	0	0.0	+0.3	0	0.0	+0.3	
Female Active Clinical Pts 2-74	805		631				572		
# w/BMI calculated									
-No Refusals	539	67.0	493	78.1	-11.2	429	75.0	-8.0	
A. # Overweight w/ % of Total BMI	139	25.8	139	28.2	-2.4	118	27.5	-1.7	
B. # Obese w/ % of Total BMI	217	40.3	198	40.2	+0.1	150	35.0	+5.3	
C. #Overweight/Obese w/ % of Total BMI	356	66.0	337	68.4	-2.3	268	62.5	+3.6	
# w/BMI refusal (No BMI)	2	0.2	0	0.0	+0.2	0	0.0	+0.2	

Figure 2-63: Sample Report, Obesity Assessment

Obesity Assessment (con't)								
TOTAL ACTIVE CLINICAL POPULATION								
Age Distribution								
	2-5	6-11	12-19	20-24	25-34	35-44	45-54	55-74
CURRENT REPORT PERIOD								
Total # Active Clin	109	112	157	133	233	215	216	225
# w/BMI calculated								
-No Refusals	52	44	89	117	183	146	128	121
% w/BMI calculated								
-No Refusals	47.7	39.3	56.7	88.0	78.5	67.9	59.3	53.8

# A. Overweight	9	10	21	33	44	39	38	48
% A. Overweight w/ % Total BMI	17.3	22.7	23.6	28.2	24.0	26.7	29.7	39.7
# B. Obese	7	13	28	39	86	86	61	52
% B. Obese w/ % of Total BMI	13.5	29.5	31.5	33.3	47.0	58.9	47.7	43.0
# C. Overweight or Obese	16	23	49	72	130	125	99	100
% C. Overweight or Obese w/ % Total BMI	30.8	52.3	55.1	61.5	71.0	85.6	77.3	82.6
# w/BMI refusal (No BMI)	1	0	0	0	0	1	1	1
% w/BMI refusal (No BMI)	1.9	0.0	0.0	0.0	0.0	0.7	0.8	0.8
PREVIOUS YEAR PERIOD								
Total # Active Clin	111	120	137	128	172	151	140	137
# w/BMI calculated -No Refusals	49	56	88	114	155	129	113	120
% w/BMI calculated -No Refusals	44.1	46.7	64.2	89.1	90.1	85.4	80.7	87.6
# A. Overweight	7	11	20	38	47	33	36	45
% A. Overweight w/ % Total BMI	14.3	19.6	22.7	33.3	30.3	25.6	31.9	37.5
# B. Obese	14	14	26	35	65	77	56	52
% B. Obese w/ % of Total BMI	28.6	25.0	29.5	30.7	41.9	59.7	49.6	43.3
# C. Overweight or Obese	21	25	46	73	112	110	92	97
% C. Overweight or Obese w/ % Total BMI	42.9	44.6	52.3	64.0	72.3	85.3	81.4	80.8
# w/BMI refusal (No BMI)	0	0	0	0	0	0	0	0
% w/BMI refusal (No BMI)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CHANGE FROM PREV YR %								
w/BMI calculated -No Refusals	+3.6	-7.4	-7.5	-1.1	-11.6	-17.5	-21.5	-33.8
A. Overweight	+3.0	+3.1	+0.9	-5.1	-6.3	+1.1	-2.2	+2.2
B. Obese	-15.1	+4.5	+1.9	+2.6	+5.1	-0.8	-1.9	-0.4
C. Overweight or Obese	-12.1	+7.6	+2.8	-2.5	-1.2	+0.3	-4.1	+1.8
w/BMI refusal (No BMI)	+1.9	+0.0	+0.0	+0.0	+0.0	+0.7	+0.8	+0.8

Figure 2-64: Sample Report, Age Breakout, Obesity Assessment

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Obesity Assessment: List of patients with current BMI, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,PAMELA	000001	COMMUNITY #1	F	3	UP;AC	16.03	
PATIENT2,GLENDA	000002	COMMUNITY #1	F	3	UP;AC	17.49	
PATIENT3,SHIRLEY	000003	COMMUNITY #1	F	5	UP		
PATIENT4,MARY ANNE	000004	COMMUNITY #1	F	5	UP;AC	Refused	
PATIENT5,JACKIE	000005	COMMUNITY #1	F	9	UP		
PATIENT6,ZINNIA	000006	COMMUNITY #1	F	15	UP	29.41 [OVERWEIGHT]	
PATIENT7,MARY RYAN	000007	COMMUNITY #1	F	15	UP;AC	33.69 [OBESE]	

Figure 2-65: Sample Patient List, Obesity Assessment

2.8.2 Childhood Weight Control

GPRA Description

During FY 2011, achieve the tentative long-term target rate of 24% for the proportion of children with a BMI of 95% or higher.

Denominators

Active Clinical patients aged 2–5 for whom a BMI could be calculated. Broken down by gender and age groups (2, 3, 4, 5).

Numerators

Patients with BMI in the 85th to 94th percentile.

Patients with a BMI at or above the 95th percentile.

Patients with a BMI at or above the 85th percentile.

Logic Description

BMI calculation definition: All patients for whom a BMI could be calculated and who are between the ages of two and five at the beginning of the Report Period and who do not turn age six during the Report Period are included in this measure. Age in the age groups is calculated based on the date of the most current BMI found. For example, a patient may be two years of age at the beginning of the time period, but is three years old at the time of the most current BMI found. That patient will fall into the age three group. CRS looks for the most recent BMI in the Report Period. CRS calculates BMI at the time the report is run using NHANES II. A height and weight must be taken on the same day any time during the Report Period. The BMI values for this measure are reported differently than in Obesity Assessment since this age group is children ages 2-5, whose BMI values are age-dependent. The BMI values are categorized as Overweight for patients with a BMI in the 85th to 94th percentile and Obese for patients with a BMI at or above the 95th percentile.

Patients whose BMI either is greater or less than the Data Check Limit range shown below will not be included in the report counts for Overweight or Obese.

BMI Standard Reference Data

Low-High Ages	Sex	BMI >= (OVERWT)	BMI >= (OBESE)	Data Check Limits BMI >	Data Check Limits BMI <
2-2	MALE	17.7	18.7	36.8	7.2
	FEMALE	17.5	18.6	37.0	7.1
3-3	MALE	17.1	18.0	35.6	7.1
	FEMALE	17.0	18.1	35.4	6.8
4-4	MALE	16.8	17.8	36.2	7.0
	FEMALE	16.7	18.1	36.0	6.9
5-5	MALE	16.9	18.1	36.0	6.9
	FEMALE	16.9	18.5	39.2	6.8

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients ages 2–5, with current BMI.

Measure Source

CDC, National Center for Health Statistics, HP 2020 NWS-10.1

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	25.0%
IHS FY 2009 Performance	25.0%
IHS FY 2008 Performance	24.0%
IHS FY 2007 Performance	24.0%
IHS FY 2006 Performance	24.0%
<i>HP 2020 Goal</i>	<i>9.6%</i>

DU	May 25, 2011						Page 156	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Childhood Weight Control (con't)								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
2-5 w/BMI	44		39			40		
# w/BMI 85-94%	7	15.9	5	12.8	+3.1	10	25.0	-9.1
# w/BMI =>95%	5	11.4	9	23.1	-11.7	5	12.5	-1.1
# w/BMI =>85%	12	27.3	14	35.9	-8.6	15	37.5	-10.2
Active Clinical Pts								
Age 2	2		8			5		
# w/BMI 85-94%	1	50.0	0	0.0	+50.0	1	20.0	+30.0
# w/BMI =>95%	0	0.0	2	25.0	-25.0	0	0.0	+0.0
# w/BMI =>85%	1	50.0	2	25.0	+25.0	1	20.0	+30.0
Active Clinical Pts								
Age 3	23		15			8		
# w/BMI 85-94%	2	8.7	2	13.3	-4.6	3	37.5	-28.8
# w/BMI =>95%	3	13.0	3	20.0	-7.0	2	25.0	-12.0
# w/BMI =>85%	5	21.7	5	33.3	-11.6	5	62.5	-40.8
Active Clinical Pts								
Age 4	12		10			17		
# w/BMI 85-94%	1	8.3	2	20.0	-11.7	3	17.6	-9.3
# w/BMI =>95%	1	8.3	2	20.0	-11.7	2	11.8	-3.4
# w/BMI =>85%	2	16.7	4	40.0	-23.3	5	29.4	-12.7
Active Clinical Pts								
Age 5	7		6			10		
# w/BMI 85-94%	3	42.9	1	16.7	+26.2	3	30.0	+12.9
# w/BMI =>95%	1	14.3	2	33.3	-19.0	1	10.0	+4.3

# w/BMI =>85%	4	57.1	3	50.0	+7.1	4	40.0	+17.1
---------------	---	------	---	------	------	---	------	-------

Figure 2-66: Sample Report, Childhood Weight Control

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Childhood Weight Control: List of patients ages 2-5, with current BMI.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,MELISSA ANN 08/20/11 16.03	000001	COMMUNITY #1	F	4	AC	Age at BMI: 4;	
PATIENT2,RANDY 17.96:05/06/11 [OVERWEIGHT]	000002	COMMUNITY #1	M	2	AC	Age at BMI: 2;	
PATIENT3,PAUL BARRY 19.87:08/05/11 [OBESE]	000003	COMMUNITY #1	M	2	AC	Age at BMI: 2;	
PATIENT4,TYLER 15.67:02/19/11	000004	COMMUNITY #1	M	4	AC	Age at BMI: 4;	
PATIENT5,SAMUEL III 19.07:12/29/11 [OBESE]	000005	COMMUNITY #1	M	5	AC	Age at BMI: 5;	
PATIENT21,JOSEPHINE 15.71:05/30/11	000021	COMMUNITY #2	F	4	AC	Age at BMI: 4;	

Figure 2-67: Sample Patient List, Childhood Weight Control

2.8.3 Nutrition and Exercise Education for At Risk Patients

Denominators

Active Clinical patients ages six and older considered overweight (including obese). Broken down by gender.

- Active Clinical patients ages 6 and older *considered obese*. Broken down by gender and age groups (6-11, 12-19, 20-39, 40-59, 60+).

Active Diabetic patients, defined as all Active Clinical patients diagnosed with diabetes at least one year prior to the end of the Report Period, *and* at least two visits in the past year, *and* two diabetes-related visits ever.

Numerators

Patients provided with medical nutrition therapy during the Report Period.

Patients provided specific nutrition education during the Report Period.

Patients provided specific exercise education during the Report Period.

Patients provided with other related exercise and nutrition (lifestyle) education.

Logic Description

Age of the patient is calculated at beginning of Report Period.

Diabetes: First DM Purpose of Visit 250.00–250.93 recorded in the V POV file prior to the Report period.

Overweight: Ages 19 and older, BMI equal to or greater than (\Rightarrow) 25. Overweight is defined as including both obese and overweight categories calculated by BMI.

Obese: Ages 19 and older, BMI equal to or greater than (\Rightarrow) 30. For ages 18 and under, the definition is based on standard tables. CRS calculates BMI at the time the report is run, using NHANES II. For 18 and under, a height and weight must be taken on the same day any time in the year prior to the end of the Report Period. For 19 through 50, height and weight must be recorded within last 5 years, not required to be on the same day. For over 50, height and weight within last 2 years; not required to be recorded on same day.

CRS uses any of the following codes to define the numerators.

	CPT Codes	ICD and Other Codes
Medical nutrition therapy	97802-97804, G0270, G0271	Primary or secondary provider codes: 07, 29 Clinic codes: 67 (dietary) or 36 (WIC)
Nutrition education		V POV: V65.3 dietary surveillance and counseling Patient education codes: ending “-N” (nutrition), “-MNT” (medical nutrition therapy), (or old code “-DT” (diet)) or containing V65.3, 97802-97804, G0270, or G0271.
Exercise education		V POV: V65.41 exercise counseling Patient education codes: ending “-EX” (exercise) or containing V65.41.
Related exercise and nutrition education		Patient education codes: ending “-LA” (lifestyle adaptation) or containing “OBS-” (obesity) or 278.00 or 278.01.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

A list of at risk patients with education, if any.

Measure Source

HP 2010 19–17

Measure Past Performance and Long-Term Targets for Diabetic Education

Performance	Percent
HP 1997 data	42.0%

DU	May 25, 2011						Page 160	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Nutrition and Exercise Education for At Risk Patient								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
# Overweight Active Clinical patients =>6	598		555			442		
# w/medical nutrition therapy	44	7.4	23	4.1	+3.2	27	6.1	+1.2
# specific nutrition education provided	83	13.9	79	14.2	-0.4	78	17.6	-3.8
# w/exercise educ	34	5.7	28	5.0	+0.6	35	7.9	-2.2
# w/ other exec or nutrition educ	75	12.5	59	10.6	+1.9	24	5.4	+7.1
# Male Overweight Active Clinical pts =>6	251		230			182		
# w/medical nutrition therapy	18	7.2	8	3.5	+3.7	10	5.5	+1.7
# specific nutrition education provided	36	14.3	32	13.9	+0.4	28	15.4	-1.0
# w/exercise educ	16	6.4	12	5.2	+1.2	16	8.8	-2.4
# w/ other exec or nutrition educ	41	16.3	22	9.6	+6.8	11	6.0	+10.3
# Female Overweight Active Clinical pts =>6	347		325			260		
# w/medical nutrition therapy	26	7.5	15	4.6	+2.9	17	6.5	+1.0
# specific nutrition education provided	47	13.5	47	14.5	-0.9	50	19.2	-5.7
# w/exercise educ	18	5.2	16	4.9	+0.3	19	7.3	-2.1
# w/ other exec or nutrition educ	34	9.8	37	11.4	-1.6	13	5.0	+4.8

Figure 2-68: Sample Report, Nutrition and Exercise Education for At Risk Patients

Nutrition and Exercise Education for At Risk Patient (con't)					
TOTAL OBESE ACTIVE CLINICAL POPULATION					
Age Distribution					
# Obese Active Clinical	6-11	12-19	20-39	40-59	=>60
CURRENT REPORT PERIOD					
# Obese Active Clinical	13	28	167	125	32
# w/medical nutrition therapy	0	2	15	10	3
% w/medical nutrition therapy	0.0	7.1	9.0	8.0	9.4
# # w/specific nutrition education provided	0	3	22	28	7
% # w/specific nutrition education provided	0.0	10.7	13.2	22.4	21.9
# w/exercise educ	0	1	8	14	5
% w/exercise educ	0.0	3.6	4.8	11.2	15.6
# w/other exec or nutrition educ	0	3	18	16	7
% w/other exec or nutrition educ	0.0	10.7	10.8	12.8	21.9
PREVIOUS YEAR PERIOD					
# Obese Active Clinical	14	26	137	116	32
# w/medical nutrition therapy	0	3	8	3	2
% w/medical nutrition therapy	0.0	11.5	5.8	2.6	6.3
# # w/specific nutrition education provided	0	2	19	22	7
% # w/specific nutrition education provided	0.0	7.7	13.9	19.0	21.9
# w/exercise educ	0	0	4	14	4
% w/exercise educ	0.0	0.0	2.9	12.1	12.5
# w/other exec or nutrition educ	0	2	13	23	3
% w/other exec or nutrition educ	0.0	7.7	9.5	19.8	9.4
CHANGE FROM PREV YR %					
medical nutrition therapy	+0.0	-4.4	+3.1	+5.4	+3.1
Spec nutr ed	+0.0	+3.0	-0.7	+3.4	+0.0
w/exercise educ	+0.0	+3.6	+1.9	-0.9	+3.1
w/other exec or nutrition educ	+0.0	+3.0	+1.3	-7.0	+12.5

Figure 2-69: Sample Age Breakout Report, Nutrition and Exercise Education for At Risk Patients

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Nutrition and Exercise Education for At Risk Patients: List of at risk patients, with education if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,SANDRA KAY	000001	COMMUNITY #1	F	21	AC-OW,AC-OB	LIFE: 05/15/11	TO-
LA							
PATIENT2,CAITLYN	000002	COMMUNITY #1	F	22	AC-OW	NUTR: 03/15/11	
UTI-N SN;							
PATIENT3,BRITNEY	000003	COMMUNITY #1	F	22	AC-OW,AC-OB	MNT: 03/04/11	Prv:
29;							
PATIENT4,LORETTA	000004	COMMUNITY #1	F	22	AC-OW,AC-OB	NUTR: 05/07/11	
HTN-N SN; EXER ED: 05/07/11							HTN-EX;
PATIENT5,HALEY	000005	COMMUNITY #1	F	25	AC-OW,AC-OB		
PATIENT6,BRITTANY	000006	COMMUNITY #1	F	25	AC-OW,AC-OB	EXER ED: 01/15/11	
278.00-EX;LIFE: 01/15/11							278.00-EX

Figure 2-70: Sample Patient List, Nutrition, and Exercise Education for At Risk Patients

2.8.4 Physical Activity Assessment

Denominators

Active Clinical patients ages 5 and older. Broken down by gender and age groups (5-11, 12-19, 20-24, 25-34, 35-44, 45-54, 55-74, >75).

Numerator 1 (Active Clinical Patients assessed for physical activity during the Report Period). Broken down by gender and age groups (5-11, 12-19, 20-24, 25-34, 35-44, 45-54, 55-74, >75).

User Population patients ages 5 and older. Broken down by gender.

Numerator 1 (User Population Patients assessed for physical activity during the Report Period). Broken down by gender.

Numerators

Patients assessed for physical activity during the Report Period.

Patients from Numerator 1 who have received exercise education following their physical activity assessment.

Logic Description

Age of the patient is calculated at beginning of Report Period.

CRS uses any of the following codes to define the numerators.

	ICD and Other Codes
Physical Activity Assessment	Health Factors: Any health factor for category Activity Level documented during the Report Period.
Exercise education	V POV: V65.41 exercise counseling Patient education codes: ending "-EX" (exercise) or containing V65.41.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with physical activity assessment and any exercise education.

DU	May 25, 2011						Page 171	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Physical Activity Assessment								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
5 and older	1,357		1,031			911		
# w/ physical activity assessment	19	1.4	0	0.0	+1.4	0	0.0	+1.4
# w/ exercise educ w/ % of physical activity assessment	4	21.1	0	0.0	+21.1	0	0.0	+21.1
Male Active Clinical =>5	570		430			374		
# w/ physical activity assessment	7	1.2	0	0.0	+1.2	0	0.0	+1.2
# w/ exercise educ w/ % of physical activity assessment	1	14.3	0	0.0	+14.3	0	0.0	+14.3
Female Active Clinical =>5	787		601			537		
# w/ physical activity assessment	12	1.5	0	0.0	+1.5	0	0.0	+1.5
# w/ exercise educ w/ % of physical activity assessment	3	25.0	0	0.0	+25.0	0	0.0	+25.0

User Pop ages 5 and older	2,592		2,142			2,025		
# w/ physical activity assessment	19	0.7	0	0.0	+0.7	0	0.0	+0.7
# w/ exercise educ w/ % of physical activity assessment	4	21.1	0	0.0	+21.1	0	0.0	+21.1
Male User Pop Pts =>5	1,211		988			945		
# w/ physical activity assessment	7	0.6	0	0.0	+0.6	0	0.0	+0.6
# w/ exercise educ w/ % of physical activity assessment	1	14.3	0	0.0	+14.3	0	0.0	+14.3
Female User Pop Pts =>5	1,381		1,154			1,080		
# w/ physical activity assessment	12	0.9	0	0.0	+0.9	0	0.0	+0.9
# w/ exercise educ w/ % of physical activity assessment	3	25.0	0	0.0	+25.0	0	0.0	+25.0

Figure 2-71: Sample Report, Physical Activity Assessment

Physical Activity Assessment (con't)								
	TOTAL ACTIVE CLINICAL5 AND OLDER Age Distribution							
	5-11	12-19	20-24	25-34	35-44	45-54	55-74	>74 yrs
CURRENT REPORT PERIOD								
Total # AC Pts =>5	132	157	133	233	215	216	225	46
# w/ physical activity assessment	2	6	3	1	2	0	3	2
% w/ physical activity assessment	1.5	3.8	2.3	0.4	0.9	0.0	1.3	4.3
# w/ exercise educ w/ % of physical activity assessment	1	2	1	0	0	0	0	0
% w/ exercise educ w/ % of physical activity assessment	50.0	33.3	33.3	0.0	0.0	0.0	0.0	0.0
PREVIOUS YEAR PERIOD								
Total # AC Pts =>5	141	137	128	172	151	140	137	25
# w/ physical activity assessment	0	0	0	0	0	0	0	0
% w/ physical activity assessment	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
# w/ exercise educ w/ % of physical activity								

assessment	0	0	0	0	0	0	0	0
% w/ exercise educ w/ % of physical activity assessment	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
CHANGE FROM PREV YR % # w/ physical activity assessment	+1.5	+3.8	+2.3	+0.4	+0.9	+0.0	+1.3	+4.3
w/ exercise educ w/ % of physical activity assessment	+50.0	+33.3	+33.3	+0.0	+0.0	+0.0	+0.0	+0.0

Figure 2-72: Sample Age Breakout Report, Physical Activity Assessment

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Physical Activity Assessment: List of patients with physical activity assessment and any exercise education.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,MISTY DAWN VERY ACTIVE; EXER ED: 08/08/11 OBS-EX	000001	COMMUNITY #1	F	5	UP;AC	PHYS ACT: 08/08/11	
PATIENT2,RITA ANN SOME ACTIVITY; EXER ED: 03/06/11 TO-EX	000002	COMMUNITY #1	F	15	UP;AC	PHYS ACT: 03/06/11	
PATIENT3,RHONDA SUE ACTIVE; EXER ED: 04/02/11 V65.41	000003	COMMUNITY #1	F	22	UP;AC	PHYS ACT: 04/02/11	
PATIENT4,MARY SOME ACTIVITY;	000004	COMMUNITY #1	F	28	UP;AC	PHYS ACT: 12/12/11	
PATIENT5,JOSEPH HENRY SOME ACTIVITY;	000005	COMMUNITY #1	M	12	UP;AC	PHYS ACT: 08/02/11	
PATIENT6,BOB INACTIVE; EXER ED: 05/05/11 OBS-EX	000006	COMMUNITY #1	M	17	UP;AC	PHYS ACT: 05/05/11	

Figure 2-73: Sample Patient List, Physical Activity Assessment

2.8.5 Comprehensive Health Screening

Denominators

Active Clinical patients ages 2 and older.

Active Clinical patients ages 12 to 75.

Active Clinical patients ages 18 and older.

Female Active Clinical patients ages 15–40.

Active Clinical patients ages 5 and older.

Active Clinical patients ages 2 through 74.

Active Clinical patients ages 20 and over.

Active Clinical patients ages 5 and older.

Numerators

ALL Comprehensive Health Screening: Patients with Comprehensive Health Screening for which they are eligible, defined as having alcohol, depression, and Intimate Partner Violence/Domestic Violence (IPV/DV) screening, BMI calculated, and tobacco use, BP, and physical activity assessed.

Note: This does *not* include refusals.

Comprehensive Health Screening: Patients with Comprehensive Health Screening minus physical activity assessment for which they are eligible, defined as having alcohol, depression, and IPV/DV screening, BMI calculated, and tobacco use and BP assessed.

Note: This does *not* include physical activity assessment and does *not* include refusals.

Alcohol Screening: Patients screened for alcohol use or had an alcohol-related diagnosis or procedure during the Report Period.

Note: This numerator does *not* include refusals or alcohol-related patient education.

Depression Screening: Patients screened for depression or diagnosed with a mood disorder at any time during the Report Period.

Note: This numerator does *not* include refusals.

IPV/DV Screening: Patients screened for IPV/DV at any time during the Report Period.

Note: This numerator does *not* include refusals.

Tobacco Use Assessed: Patients who have been screened for tobacco use during the Report period.

BMI Available: Patients for whom a BMI could be calculated.

Note: This numerator does *not* include refusals.

BP Assessed: Patients with BP value documented at least twice in prior two years.

Physical Activity Assessed: Patients assessed for physical activity during the Report Period.

Logic Description

Age of the patient is calculated at beginning of Report Period.

Alcohol screening definition: Any of the following during the Report Period: (a) Alcohol Screening Exam, any CAGE Health Factor, or Screening Diagnosis; (b) Alcohol-related diagnosis in POV, Current PCC or BHS Problem List; (c) Alcohol-related procedure; or (d) Patient education.

	ICD and Other Codes
Alcohol Screening	PCC Exam Code: 35 CPT code: 99408, 99409, G0396, G0397, H0049, H0050, 3016F Any CAGE Health Factor V POV: V11.3 (history of alcoholism), V79.1 (screening for alcoholism) BHS Problem Code: 29.1 (Screening for Alcoholism) V Measurement in PCC or BHS: AUDT, AUDC, or CRFT
Alcohol-related Diagnosis	V POV, Current PCC or BHS Problem List: 303.*, 305.0*, 291.*, 357.5* BHS POV: 10, 27, 29
Alcohol-related Procedure	V Procedure: 94.46, 94.53, 94.61-94.63, 94.67-94.69

Alcohol screening may be documented with either an exam code or the CAGE health factor in PCC or BHS. BHS problem codes can also currently be used.

Depression screening definition: CRS uses the following codes to define the numerator.

	ICD and Other Codes
Depression Screening	V Exam: Exam Code 36 V POV: V79.0 CPT: 1220F BHS Problem Code: 14.1 (Screening for Depression) V Measurement in PCC or BHS: PHQ2 or PHQ9
Mood Disorders	At least 2 visits in PCC or BHS for: Major Depressive Disorder, Dysthymic Disorder, Depressive Disorder NOS, Bipolar I or II Disorder, Cyclothymic Disorder, Bipolar Disorder NOS, Mood Disorder Due to a General Medical Condition, Substance-induced Mood Disorder, or Mood Disorder NOS. V POV: 296.*, 291.89, 292.84, 293.83, 300.4, 301.13, or 311 BHS POV: 14, 15

IPV/DV screening definition: CRS uses the following codes to define the numerator.

	ICD and Other Codes
IPV/DV Screening	V Exam: Code 34 BHS Exam: IPV/DV
IPV/DV Diagnosis	V POV or current PCC or BHS Problem List: 995.80-995.83, 995.85, V15.41, V15.42, V15.49 BHS POV: 43.*, 44.*
IPV/DV Education	Patient education codes: "DV-" or "-DV", 995.80-83, 995.85, V15.41, V15.42, or V15.49
IPV/DV Counseling	V POV: V61.11

Tobacco screening definition: CRS uses the following codes to define the numerator.

	CPT Codes	ICD and Other Codes
Screened for Tobacco Use	D1320, 99406, 99407, G0375 (old code), G0376 (old code), G8455-G8457 (old codes), G8402 (old code), G8453 (old code), 1034F (Current Tobacco Smoker), 1035F (Current Smokeless Tobacco User), 1036F (Current Tobacco Non-User), 1000F (Tobacco Use Assessed)	V POV or current Active Problem List: 305.1, 305.1* (old codes), 649.00-649.04, V15.82 Patient Education codes: "TO-", "-TO", "-SHS", 305.1, 305.1* (old codes), 649.00-649.04, V15.82, D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455-G8457 (old codes), G8402 (old code), or G8453 (old code) Dental code: 1320 Health Factor categories: Tobacco, TOBACCO (SMOKING), TOBACCO (SMOKELESS – CHEWING/DIP), or TOBACCO (EXPOSURE)

BMI calculation definition: CRS calculates BMI at the time the report is run, using NHANES II. For 18 and under, a height and weight must be taken on the same day any time during the report period. For 19 through 50, height and weight must be recorded within last 5 years, not required to be on the same day. For over 50, height and weight within last 2 years, not required to be recorded on same day.

Blood pressure definition: CRS uses mean of last three Blood Pressures documented on non-ER visits in the past two years. If three BPs are not available, uses mean of last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) blood pressures and dividing by three (or two). If the systolic and diastolic values do not *both* meet the current category, then the value that is least controlled determines the category.

If CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F–3080F documented on a non-ER visit during the Report Period.

Physical Activity Assessment definition: CRS uses the following codes to define the numerator.

	ICD and Other Codes
Physical Activity Assessment	Health Factors: Any health factor for category Activity Level documented during the Report Period.

Key Logic Changes from CRS Version 11.0

1. Added CPT code 1220F to Depression Screening definition.
2. Added CPT code 3016F to Alcohol Screening definition.
3. Added CPT codes 0001F and 2000F to BP Documented definition.
4. For Tobacco Screening, noted that G8453, G8455, G8456, G8457, and G8402 are old codes.

Patient List Description

List of patients with assessments received, if any.

DU	May 25, 2011						Page 181
*** IHS 2011 Selected Measures with Community Specified Report ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Previous Year Period: Jan 01, 2010 to Dec 31, 2010							
Baseline Period: Jan 01, 2000 to Dec 31, 2000							

Comprehensive Health Screening							
	REPORT	%	PREV YR	%	CHG from	BASE	%
	PERIOD		PERIOD		PREV YR %	PERIOD	CHG from
							BASE %
Active Clinical ages							
2 and older	1,446		1,121			1,007	

# w/ Comprehensive Health Screening-No Refusals	56	3.9	39	3.5	+0.4	40	4.0	-0.1
# w/ Comprehensive Health Screening-No Refusals or Phys Activity	67	4.6	46	4.1	+0.5	53	5.3	-0.6
Active Clinical ages 12-75	1,180		867			753		
# w/ alcohol screening/Dx/Proc/-No Refusals or Pt Ed	95	8.1	12	1.4	+6.7	3	0.4	+7.7
Active Clinical => 18	1,112		793			666		
# w/ Depression screening or Mood Disorder Dx-No Refusals	81	7.3	42	5.3	+2.0	17	2.6	+4.7
# Female Active Clinical ages 15-40	380		311			267		
# w/IPV/DV screening-No Refusals	11	2.9	1	0.3	+2.6	0	0.0	+2.9
# Active Clinical Pts =>5	1,357		1,031			911		
# w/Tobacco Screening	624	46.0	426	41.3	+4.7	328	36.0	+10.0
Active Clinical Pts 2-74	1,400		1,096			982		
# w/BMI calculated-No Refusals	880	62.9	824	75.2	-12.3	712	72.5	-9.6
Active Clinical Patients ages 20 and older	1,068		753			640		
# w/ BPs documented w/in 2 yrs	643	60.2	557	74.0	-13.8	478	74.7	-14.5
Active Clinical Pts 5 and older	1,357		1,031			911		
# w/ physical activity assessment	19	1.4	0	0.0	+1.4	0	0.0	+1.4

Figure 2-74: Sample Report, Comprehensive Health Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Comprehensive Health Screening: List of patients with assessments

received, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,SANDRA KAY	000001	COMMUNITY #1	F	15	AC	ALL COMP HEALTH:
ALC: 03/06/11 POV V11.3; IPV: 03/06/11 Ex 34; TOB: 09/05/11 NEVER SMOKED; BMI:						
17.49; PHYS ACT: 03/06/11 SOME ACTIVITY						
PATIENT2,CAITLYN	000002	COMMUNITY #1	F	16	AC	
PATIENT3,BRITNEY	000003	COMMUNITY #1	F	16	AC	TOB: 10/26/11
CESSATION-SMOKER						
PATIENT4,LORETTA	000004	COMMUNITY #1	F	17	AC	ALC: 10/14/11 HF
CAGE 1/4						
PATIENT5,HALEY	000005	COMMUNITY #1	F	18	AC	BMI: 19.79; BP:
125/67						
PATIENT6,BRITTANY	000006	COMMUNITY #1	F	19	AC	ALC: 10/30/11 CPT
G0397; TOB: 08/11/11 CURRENT SMOKER, STATUS UNKNOWN; BMI: 21.01						

Figure 2-75: Sample Patient List, Comprehensive Health Screening

2.8.6 Cardiovascular Disease and Cholesterol Screening

Denominators

Active Clinical patients ages 23 and older. Broken down by gender.

User Population patients ages 23 and older. Broken down by gender.

Active IHD patients, defined as all Active Clinical patients diagnosed with ischemic heart disease (IHD) prior to the Report Period, *and* at least two visits during the Report Period, *and* two IHD-related visits ever. Broken down by gender.

Numerators

Patients with documented blood total cholesterol screening any time in the past five years.

- Patients with high total cholesterol levels, defined as equal to or greater than (\geq) 240

Patients with LDL completed in the past five years, regardless of result.

- Patients with LDL ≤ 100
- Patients with LDL 101–130
- Patients with LDL 131–160
- Patients with LDL > 160

Logic Description

Age is calculated at the beginning of the Report Period.

CRS uses the following codes to define the IHD denominator.

	ICD and Other Codes
Ischemic Heart Disease	V POV: 410.0-412.*, 414.0-414.9, 429.2

Total Cholesterol definition: Searches for most recent cholesterol test with a result during the Report Period. If more than one cholesterol test is found on the same day and/or the same visit and one test has a result and the other does not, the test with the result will be used. If a cholesterol test with a result is not found, CRS searches for the most recent cholesterol test without a result.

LDL Cholesterol definition: Searches for most recent LDL test with a result during the Report Period. If more than one LDL test is found on the same day and/or the same visit and one test has a result and the other does not, the test with the result will be used. If an LDL test with a result is not found, CRS searches for the most recent LDL test without a result.

CRS uses the following codes to define LDL and total cholesterol.

Test	CPT Codes	LOINC Codes	Taxonomy
LDL	80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F For numerator LDL =<100, CPT 3048F will count as meeting the measure.	Yes	DM AUDIT LDL CHOLESTEROL TAX
Total Cholesterol	82465	Yes	DM AUDIT CHOLESTEROL TAX

Key Logic Changes from CRS Version 11.0

1. Removed the CPT code range 428.* from Ischemic Heart Disease definition.
2. Added code 43396-1 to LOINC taxonomy for LDL.

Patient List Description

List of patients with cholesterol or LDL value, if any.

Measure Source

HP 2020 HDS-6, HDS-7

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2006 Performance (blood total cholesterol screening)	48.0%
IHS FY 2005 Performance (blood total cholesterol screening)	43.0%
HP 1998 baseline	67.0%
HP 2020 goal for adults who have had blood cholesterol checked (HDS-6)	82.1%
HP 2020 goal for adults with high cholesterol (HDS-7)	13.5%

DU	May 25, 2011							Page 185	
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Cardiovascular Disease and Cholesterol Screening									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical Pts => 23	986		676			569			
# w/ Total Cholesterol screen w/in 5 yrs	248	25.2	217	32.1	-6.9	201	35.3	-10.2	
A. # w/ High Chol =>240 w/ % of Total Chol Screen	21	8.5	23	10.6	-2.1	28	13.9	-5.5	
# w/LDL done in past 5 yrs	251	25.5	185	27.4	-1.9	114	20.0	+5.4	
A. # w/LDL =<100 w/ % of Total LDL Screen	108	43.0	95	51.4	-8.3	46	40.4	+2.7	
B. # w/LDL 101-130 w/ % of Total LDL Screen	73	29.1	44	23.8	+5.3	35	30.7	-1.6	
C. # w/LDL 131-160 w/ % of Total LDL Screen	25	10.0	25	13.5	-3.6	13	11.4	-1.4	
D. # w/LDL >160 w/ % of Total LDL Screen	15	6.0	9	4.9	+1.1	10	8.8	-2.8	
Male Active Clinical Pts =>23	408		272			220			
# w/ Total Cholesterol screen w/in 5 yrs	106	26.0	97	35.7	-9.7	85	38.6	-12.7	
A. # w/ High Chol =>240 w/ % of Total Chol Screen	11	10.4	14	14.4	-4.1	8	9.4	+1.0	
# w/LDL done in past 5 yrs	115	28.2	91	33.5	-5.3	59	26.8	+1.4	

A. # w/LDL ≤100 w/ % of Total LDL Screen	53	46.1	45	49.5	-3.4	23	39.0	+7.1
B. # w/LDL 101-130 w/ % of Total LDL Screen	25	21.7	18	19.8	+2.0	18	30.5	-8.8
C. # w/LDL 131-160 w/ % of Total LDL Screen	8	7.0	12	13.2	-6.2	5	8.5	-1.5
D. # w/LDL >160 w/ % of Total LDL Screen	10	8.7	8	8.8	-0.1	4	6.8	+1.9

Figure 2-76: Sample Report, CVD and Cholesterol Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Cardiovascular Disease and Cholesterol Screening: List of patients with cholesterol or LDL value, if any							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT100,JASON AARON	000100	COMMUNITY #1	M	46	UP;AC		
PATIENT101,JOHN THOMAS	000101	COMMUNITY #1	M	47	UP		
PATIENT102,DAKOTA CHEY	000102	COMMUNITY #1	M	47	UP		
PATIENT103,TRAVIS CLINT	000103	COMMUNITY #1	M	47	UP	CHOL	04/13/11 210
PATIENT104,TRACY MITCHE	000104	COMMUNITY #1	M	47	UP;AC;IHD	CHOL	03/15/11 167;
LDL 08/15/11 105							
PATIENT105,RUSSELL DALE	000105	COMMUNITY #1	M	48	UP	LDL	04/01/11 CPT:
3048F							
PATIENT106,CURTIS DWAYN	000106	COMMUNITY #1	M	49	UP;AC	CHOL	03/04/08 139;
LDL 06/04/09 68							
PATIENT107,RONALD	000107	COMMUNITY #1	M	49	UP;AC	CHOL	01/07/07 213;
LDL 08/01/07 122							

Figure 2-77: Sample Patient List, CVD and Cholesterol Screening

2.8.7 Cardiovascular Disease and Blood Pressure Control

Denominators

All *Active Clinical patients* ages 20 and over. Broken down by gender.

All *User Population patients* ages 20 and older. Broken down by gender.

Active IHD patients, defined as all Active Clinical patients diagnosed with IHD prior to the Report Period, *and* at least two visits during the Report Period, *and* two IHD-related visits ever. Broken down by gender.

Numerators

Patients with Blood Pressure value documented at least twice in prior two years.

- Patients with normal BP, defined as < 120/80, i.e., the mean systolic value is less than (<) 120 *and* the mean diastolic value is less than (<) 80.

- b. Patients with Pre Hypertension I BP, defined as \Rightarrow 120/80 and $<$ 130/80, i.e., the mean systolic value is equal to or greater than (\Rightarrow) 120 and less than ($<$) 130 *and* the mean diastolic value is equal to 80.
- c. Patients with Pre Hypertension II BP, defined as \Rightarrow 130/80 and $<$ 140/90, i.e., the mean systolic value is equal to or greater than (\Rightarrow) 130 and less than ($<$) 140 *and* the mean diastolic value is equal to or greater than (\Rightarrow) 80 and less than ($<$) 90.
- d. Patients with Stage 1 Hypertension BP, defined as \Rightarrow 140/90 and $<$ 160/100, i.e., the mean systolic value is equal to or greater than (\Rightarrow) 140 and less than ($<$) 160 *and* the mean diastolic value is equal to or greater than (\Rightarrow) 90 and less than ($<$) 100.
- e. Patients with Stage 2 Hypertension BP, defined as \Rightarrow 160/100, i.e., the mean systolic value is equal to or greater than (\Rightarrow) 160 *and* the mean diastolic value is equal to or greater than (\Rightarrow) 100.

Logic Description

Age of the patient is calculated at beginning of the Report Period.

Blood pressure definition: CRS uses mean of last three Blood Pressures documented on non-ER visits in the past two years. If three BPs are not available, uses mean of last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) blood pressures and dividing by three (or two). If the systolic and diastolic values do not *both* meet the current category, then the value that is least controlled determines the category.

For the BP documented numerator only, if CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F–3080F documented on a non-ER visit during the Report Period.

CRS uses the following codes to define the IHD numerator.

	ICD and Other Codes
Ischemic Heart Disease	V POV: 410.0-412.*, 414.0-414.9, 429.2

Key Logic Changes from CRS Version 11.0

1. Removed the CPT code range 428.* from Ischemic Heart Disease definition.
2. Added CPT codes 0001F and 2000F to BP Documented definition.

Patient List Description

List of Patients => 20 or who have IHD with BP value, if any.

Measure Source

HP 2020 HDS-5

Measure Past Performance and Long-Term Targets

Measure	Percent
HP 2020 goal for adults with high blood pressure (140/90)	26.9%

DU	May 25, 2011						Page 196	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Cardiovascular Disease and Blood Pressure Control								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Patients								
ages 20 and older	1,068		753			640		
# w/ BPs documented								
w/in 2 yrs	643	60.2	557	74.0	-13.8	478	74.7	-14.5
A. # w/Normal BP w/ %								
of Total Screened	128	19.9	134	24.1	-4.2	121	25.3	-5.4
B. # w/Pre HTN I BP w/ %								
of Total Screened	107	16.6	115	20.6	-4.0	83	17.4	-0.7
C. # w/Pre HTN II BP w/ %								
of Total Screened	148	23.0	114	20.5	+2.6	105	22.0	+1.1
D. # w/Stage 1 HTN BP w/ %								
of Total Screened	173	26.9	150	26.9	+0.0	130	27.2	-0.3
E. # w/Stage 2 HTN BP w/ %								
of Total Screened	37	5.8	39	7.0	-1.2	39	8.2	-2.4
Male Active Clinical Patients								
ages 20 and older	432		293			241		
# w/ BPs documented								
w/in 2 yrs	231	53.5	203	69.3	-15.8	177	73.4	-20.0
A. # w/Normal BP w/ %								
of Total Screened	8	3.5	23	11.3	-7.9	22	12.4	-9.0
B. # w/Pre HTN I BP w/ %								
of Total Screened	27	11.7	37	18.2	-6.5	22	12.4	-0.7
C. # w/Pre HTN II BP w/ %								
of Total Screened	64	27.7	46	22.7	+5.0	45	25.4	+2.3
D. # w/Stage 1 HTN BP w/ %								
of Total Screened	89	38.5	79	38.9	-0.4	63	35.6	+2.9
E. # w/Stage 2 HTN BP w/ %								
of Total Screened	16	6.9	17	8.4	-1.4	25	14.1	-7.2

Female Active Clinical Patients ages 20 and older								
	636		460			399		
# w/ BPs documented w/in 2 yrs	412	64.8	354	77.0	-12.2	301	75.4	-10.7
A. # w/Normal BP w/ % of Total Screened	120	29.1	111	31.4	-2.2	99	32.9	-3.8
B. # w/Pre HTN I BP w/ % of Total Screened	80	19.4	78	22.0	-2.6	61	20.3	-0.8
C. # w/Pre HTN II BP w/ % of Total Screened	84	20.4	68	19.2	+1.2	60	19.9	+0.5
D. # w/Stage 1 HTN BP w/ % of Total Screened	84	20.4	71	20.1	+0.3	67	22.3	-1.9
E. # w/Stage 2 HTN BP w/ % of Total Screened	21	5.1	22	6.2	-1.1	14	4.7	+0.4

Figure 2-78: Sample Report, CVD and Blood Pressure Control

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Cardiovascular Disease and Blood Pressure Control: List of Patients => 20 or who have IHD with BP value, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT1,SANDRA KAY	000001	COMMUNITY #1	F	21	UP;AC;		
PATIENT2,EVELYN	000002	COMMUNITY #1	F	21	UP;	/3080F	
PATIENT3,MICHELLE	000003	COMMUNITY #1	F	22	UP;AC;	125/67	PRE STG 1
PATIENT4,CAITLYN	000004	COMMUNITY #1	F	22	UP;AC;IHD	131/67	PRE STG II
PATIENT5,BRITNEY JANE	000005	COMMUNITY #1	F	22	UP;AC;	102/56	NORMAL
PATIENT6,KATHRYN ANNE	000006	COMMUNITY #1	F	22	UP;AC;	161/90	STG 2 HTN
PATIENT7,RHONDA	000007	COMMUNITY #1	F	22	UP;AC;	153/85	STG 1 HTN

Figure 2-79: Sample Patient List, CVD and Blood Pressure Control

2.8.8 Controlling High Blood Pressure

Denominator

Active Clinical patients ages 18 through 85 diagnosed with hypertension and no documented history of ESRD. Broken down by gender and age groups (18–45 and 46–85).

Numerators

Number of patients with Blood Pressure value documented during the Report Period.

- Patients with *normal blood pressure*, defined as < 120/80; that is, the mean systolic value is less than (<) 120 *and* the mean diastolic value is less than (<) 80.
- Patients with *Pre Hypertension I BP*, defined as => 120/80 and less than < 130/80, that is, the mean systolic value is equal to or greater than (=>) 120 and less than (<) 130 *and* the mean diastolic value is equal to 80.

- c. Patients with *Pre Hypertension II BP*, defined as \Rightarrow 130/80 and $<$ 140/90; that is, the mean systolic value is equal to or greater than (\Rightarrow) 130 and less than ($<$) 140 and the mean diastolic value is equal to or greater than (\Rightarrow) 80 and less than ($<$) 90.
- d. Patients with *Stage 1 Hypertension Blood Pressure (BP)*, defined as \Rightarrow 140/90 and $<$ 160/100; that is, the mean systolic value is equal to or greater than (\Rightarrow) 140 and less than ($<$) 160 and the mean diastolic value is equal to or greater than (\Rightarrow) 90 and less than ($<$) 100.
- e. Patients with *Stage 2 Hypertension BP*, defined as \Rightarrow 160/100; that is, the mean systolic value is equal to or greater than (\Rightarrow) 160 and the mean diastolic value is equal to or greater than (\Rightarrow) 100.

Logic Description

Age of the patient is calculated at beginning of the Report Period.

Blood pressure definition: CRS uses mean of last three BPs documented on non-ER visits during the Report Period. If three BPs are not available, uses mean of last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) blood pressures and dividing by three (or two). If the systolic and diastolic values do not *both* meet the current category, then the value that is least controlled determines the category.

For the BP documented numerator only, if CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F–3080F documented on non-ER visit during the Report Period.

CRS uses the following codes to define ESRD and hypertension.

	CPT Codes	ICD and Other Codes
ESRD	36145, 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831-36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951-90970 or old codes 90918-90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, G0257, G0308-G0327 (old codes), G0392 (old code), G0393 (old code), or S9339	V POV: 585.5, 585.6, V42.0, V45.1, (old code), V45.11, V45.12, or V56.* V Procedure: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93-39.95, 54.98, or 55.6*

	CPT Codes	ICD and Other Codes
Hypertension		V POV or Problem List Prior to the Report Period and at Least One Hypertension POV during Report Period: 401.*

Key Logic Changes from CRS Version 11.0

1. Added CPT code 36147 to ESRD definition.
2. Added CPT codes 0001F and 2000F to BP Documented definition.

Patient List Description

List of patients with hypertension and BP value, if any.

Measure Source

HP 2020 HDS-5, HDS-12

Measure Past Performance and Long-Term Targets

Measure	Percent
HP 2020 goal for adults with high blood pressure (140/90)	26.9%
HP 2020 goal for adults with high blood pressure and whose blood pressure is controlled	61.2%

DU	May 25, 2011						Page 200	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Controlling High Blood Pressure								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
18-85 w/HTN dx	108		101			91		
# w/ BPs								
documented	96	88.9	92	91.1	-2.2	85	93.4	-4.5
A. # w/Normal BP w/ %								
of Total Screened	5	5.2	6	6.5	-1.3	3	3.5	+1.7
B. # w/Pre HTN I BP w/ %								
of Total Screened	8	8.3	13	14.1	-5.8	8	9.4	-1.1
C. # w/Pre HTN II BP w/								
% of Total Screened	23	24.0	17	18.5	+5.5	19	22.4	+1.6
D. # w/Stage 1 HTN BP w/								
% of Total Screened	36	37.5	41	44.6	-7.1	42	49.4	-11.9
E. # w/Stage 2 HTN BP w/								

% of Total Screened	10	10.4	15	16.3	-5.9	13	15.3	-4.9
A. Active Clinical Pts 18-45 w/HTN dx	23		19			13		
# w/ BPs documented	17	73.9	16	84.2	-10.3	11	84.6	-10.7
A. # w/Normal BP w/ % of Total Screened	2	11.8	2	12.5	-0.7	0	0.0	+11.8
B. # w/Pre HTN I BP w/ % of Total Screened	1	5.9	1	6.3	-0.4	1	9.1	-3.2
C. # w/Pre HTN II BP w/ % of Total Screened	3	17.6	2	12.5	+5.1	2	18.2	-0.5
D. # w/Stage 1 HTN BP w/ % of Total Screened	7	41.2	6	37.5	+3.7	7	63.6	-22.5
E. # w/Stage 2 HTN BP w/ % of Total Screened	2	11.8	5	31.3	-19.5	1	9.1	+2.7
B. Active Clinical Pts 46-85 w/HTN dx	85		82			78		
# w/ BPs documented	79	92.9	76	92.7	+0.3	74	94.9	-1.9
A. # w/Normal BP w/ % of Total Screened	3	3.8	4	5.3	-1.5	3	4.1	-0.3
B. # w/Pre HTN I BP w/ % of Total Screened	7	8.9	12	15.8	-6.9	7	9.5	-0.6
C. # w/Pre HTN II BP w/ % of Total Screened	20	25.3	15	19.7	+5.6	17	23.0	+2.3
D. # w/Stage 1 HTN BP w/ % of Total Screened	29	36.7	35	46.1	-9.3	35	47.3	-10.6
E. # w/Stage 2 HTN BP w/ % of Total Screened	8	10.1	10	13.2	-3.0	12	16.2	-6.1

Figure 2-80: Sample Report, Controlling High Blood Pressure

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease										
Controlling High Blood Pressure: List of patients with hypertension and BP value, if any.										
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR				

PATIENT1,STELLA LYNN	000001	COMMUNITY #1	F	46	HTN PT	156/82	STG 1	HTN		
PATIENT2,TARA	000002	COMMUNITY #1	F	51	HTN PT	201/87	STG 2	HTN		
PATIENT3,BOBBIE	000003	COMMUNITY #1	F	52	HTN PT	3074F/				
PATIENT4,DARLENE	000004	COMMUNITY #1	F	54	HTN PT	139/73	PRE	STG II		
PATIENT5,NADINE	000005	COMMUNITY #1	F	61	HTN PT	159/86	STG 1	HTN		

Figure 2-81: Sample Patient List, Controlling High Blood Pressure

2.8.9 Comprehensive CVD-Related Assessment

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 33.0% for the proportion of at-risk patients who have a comprehensive assessment.

Denominators

Active IHD patients ages 22 and older, defined as all Active Clinical patients diagnosed with IHD prior to the Report Period, *and* at least two visits during the Report Period, *and* two IHD-related visits ever. (GPRA Denominator)

- a. Active IHD patients ages 22 and older who are not Active Diabetic
- b. Active IHD patients ages 22 and older who are Active Diabetic

Numerators

BP Assessed: Patients with Blood Pressure value documented at least twice in prior two years.

LDL Assessed: Patients with LDL completed in past five years, regardless of result.

Tobacco Use Assessed: Patients who have been screened for tobacco use during the Current Report period.

BMI Available: Patients for whom a BMI could be calculated.

Note: This numerator does *not* include refusals.

Lifestyle Counseling: Patients who have received any lifestyle adaptation counseling, including medical nutrition therapy, or nutrition, exercise or other lifestyle education during the Report Period.

Patients with *comprehensive CVD assessment*, defined as having BP, LDL, and tobacco use assessed, BMI calculated, and lifestyle counseling.

Note: This does *not* include depression screening and *does not include refusals of BMI*. (GPRA Numerator)

Refusal of BMI: Patients who refused a height or weight measurement and for whom a BMI could not be calculated.

Depression Screening: Patients screened for depression or diagnosed with a mood disorder at any time during the Report Period.

Note: This numerator does *not* include refusals.

Logic Description

Age of the patient is calculated at beginning of the Report Period.

Diabetes defined as: Diagnosed with diabetes (first POV in V POV with 250.00–250.93) prior to the Current Report Period, *and* at least two visits during the Current Report Period, *and* two DM-related visits ever. Patients not meeting these criteria are considered non-diabetics.

IHD diagnosis defined as: 410.0–412.*, 414.0–414.9, 428.* or 429.2 recorded in the V POV file.

Blood pressure definition: Having a minimum of two BPs documented on non-ER visits during the Report Period in past two years. If CRS does not find two BPs, it will search for CPT 0001F, 2000F, 3074F–3080F documented on non-ER visit during the past two years.

LDL definition: Finds the most recent test done in the last five years, regardless of the results of the measurement.

BMI definition: CRS calculates when the report is run, using NHANES II. For 19 through 50, height and weight must be recorded within last 5 years, not required to be on the same day. For over 50, height and weight within last 2 years; not required to be recorded on same day.

BMI Refusal definition: Refusals of a height and/or weight measurement include REF, NMI, and UAS and must be documented during the past year. For ages 19 and older, the height and the weight must be refused during the past year and are not required to be on the same visit.

CRS uses the following codes and taxonomies to define the numerators.

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
LDL - Finds the most recent test done in the last 5 years, regardless of the results of the measurement.	80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F		Yes	DM AUDIT LDL CHOLESTEROL TAX

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Tobacco Screening	D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455-G8457 (old codes), G8402 (old code) or G8453 (old code)	Any health factor for category Tobacco, TOBACCO (SMOKING), TOBACCO (SMOKELESS – CHEWING/DIP), or TOBACCO (EXPOSURE) (see table on next page) V POV or current Active Problem List: 305.1, 305.1* (old codes), 649.00-649.04, V15.82 Patient education codes: containing “TO-”, “-TO”, “-SHS”, 305.1, 305.1* (old codes), 649.00-649.04, V15.82, D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455- G8457 (old codes), G8402 (old code) or G8453 (old code) Dental code: 1320		
Medical Nutrition Therapy	97802-97804, G0270, G0271	Primary or secondary provider codes: 07, 29, 97, 99 Clinic codes: 67 (dietary) or 36 (WIC)		
Nutrition Education		V POV: V65.3 dietary surveillance and counseling Patient education codes: ending “-N” (nutrition) or “-MNT” (medical nutrition therapy) (or old code “-DT” (diet)) or containing V65.3.		
Exercise Education		V POV: V65.41 exercise counseling Patient education codes: ending “-EX” (exercise) or containing V65.41.		
Related Exercise and Nutrition Education		Patient education codes: ending “-LA” (lifestyle adaptation) or containing “OBS-“ (obesity) or 278.00 or 278.01.		
Depression Screening		V Exam: Exam Code 36 V POV: V79.0 CPT: 1220F BHS Problem Code: 14.1 (Screening for Depression) V Measurement in PCC or BH: PHQ2 or PHQ9		

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Mood Disorders		At least 2 visits in PCC or BHS for: Major Depressive Disorder, Dysthymic Disorder, Depressive Disorder NOS, Bipolar I or II Disorder, Cyclothymic Disorder, Bipolar Disorder NOS, Mood Disorder Due to a General Medical Condition, Substance-induced Mood Disorder, or Mood Disorder NOS (see codes below). V POV: 296.*, 291.89, 292.84, 293.83, 300.4, 301.13, or 311 BHS POV: 14, 15		

All existing national Tobacco Health Factors listed below are counted as tobacco screening.

Health Factor	
Ceremonial	Previous Smokeless
Cessation-Smokeless	Previous (Former) Smokeless
Cessation-Smoker	Previous Smoker
Current Smokeless	Previous (Former) Smoker
Current Smoker	Smoke Free Home
Current Smoker, status unknown	Smoker In Home
Current smoker, every day	Current Smoker & Smokeless
Current smoker, some day	Exposure To Environmental Tobacco Smoke
Non-Tobacco User	

Key Logic Changes from CRS Version 11.0

1. Added CPT code 1220F to Depression Screening definition.
2. Added CPT codes 0001F and 2000F to BP Documented definition.
3. For Tobacco Screening, noted that G8453, G8455, G8456, G8457, and G8402 are old codes.
4. Added code 43396-1 to LOINC taxonomy for LDL.

Patient List Description

List of patients with assessments received, if any.

Measure Source

Not Available

Measure Past Performance Long-Term Targets

Performance	Percent
IHS FY 2010 Performance (Comprehensive CVD Assessment)	35.0%
IHS FY 2009 Performance (Comprehensive CVD Assessment)	32.0%
IHS FY 2008 Performance (Comprehensive CVD Assessment)	30.0%
IHS FY 2007 Performance (Comprehensive CVD Assessment)	30.0%
IHS FY 2008 Performance (BP Assessed)	98.0%
IHS FY 2008 Performance (LDL Assessed)	90.0%
IHS FY 2008 Performance (Tobacco Assessed)	79.0%
IHS FY 2008 Performance (BMI Assessed or Refused)	85.0%
IHS FY 2008 Performance (Lifestyle Counseling)	38.0%

DU	May 25, 2011						Page 207	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Comprehensive CVD-Related Assessment								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active IHD Pts 22+ (GPRA)	77		45			36		
# w/ BPs documented w/in 2 yrs	72	93.5	45	100.0	-6.5	36	100.0	-6.5
# w/ LDL done in past 5 yrs	60	77.9	38	84.4	-6.5	30	83.3	-5.4
# w/Tobacco Screening w/in 1 yr	66	85.7	38	84.4	+1.3	27	75.0	+10.7
# w/BMI calculated -No Refusals	72	93.5	44	97.8	-4.3	35	97.2	-3.7
# w/ lifestyle educ w/in 1 yr	43	55.8	22	48.9	+7.0	22	61.1	-5.3
# w/ BP, LDL, tobacco, BMI and life counseling -No Refusals (GPRA)	35	45.5	19	42.2	+3.2	14	38.9	+6.6
# w/BMI refusal (No BMI)	2	2.6	0	0.0	+2.6	0	0.0	+2.6
# w/ Depression screening or Mood Disorder DX -No Refusals	20	26.0	4	8.9	+17.1	2	5.6	+20.4
A. Active IHD Pts 22+ and are NOT Active								

Diabetic	41		20		17		
# w/ BPs documented							
w/in 2 yrs	36	87.8	20	100.0	-12.2	17	100.0 -12.2
# w/LDL done							
in past 5 yrs	28	68.3	17	85.0	-16.7	13	76.5 -8.2
# w/Tobacco Screening							
w/in 1 yr	33	80.5	15	75.0	+5.5	13	76.5 +4.0
# w/BMI calculated							
-No Refusals	38	92.7	20	100.0	-7.3	16	94.1 -1.4
# w/ lifestyle							
educ w/in 1 yr	24	58.5	7	35.0	+23.5	7	41.2 +17.4
# w/ BP, LDL, tobacco,							
BMI and life counseling							
-No Refusals	19	46.3	6	30.0	+16.3	4	23.5 +22.8
# w/BMI refusal							
(No BMI)	0	0.0	0	0.0	+0.0	0	0.0 +0.0
# w/ Depression screening,							
or Mood Disorder DX							
-No Refusals	13	31.7	1	5.0	+26.7	1	5.9 +25.8

Figure 2-82: Sample Report, Comprehensive CVD-Related Assessment

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Comprehensive CVD-Related Assessment: List of patients with assessments received, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,CAITLYN	000001	COMMUNITY #1	F	22	IHD	ALL: BP: 131/67 PRE
STG; LDL: 06/08/11; TOB: 07/25/11		NEVER SMOKED; BMI: 25.4; LIFE: 08/15/01				UTI-N
SN; DEP: Meas PHQ9						
PATIENT2,CARLA	000002	COMMUNITY #1	F	40	IHD	BP: 112/66 NORMAL;
TOB: 08/26/11 NEVER SMOKED; BMI: 33.7						
PATIENT3,JENNY	000003	COMMUNITY #1	F	47	IHD	BP: 3074F/; TOB:
11/01/11 D1320; BMI: 39.5; LIFE: 03/03/11						Prv: 97; DEP: CPT 1220F
PATIENT4,SHERRY	000004	COMMUNITY #1	F	68	IHD;AD	ALL: BP: 150/82 STG
1; LDL: 09/13/10; TOB: 12/30/11 NEVER SMOKED; BMI: 26.8; LIFE: 10/19/11						PM-LA
PATIENT5,PAULINE	000005	COMMUNITY #1	F	70	IHD;AD	BP: 149/72 STG 1;
LDL: 10/24/11; TOB: 10/24/11 CURRENT SMOKER, STATUS UNKNOWN; BMI: 37.5						
PATIENT6,TINA MARIE	000006	COMMUNITY #1	F	78	IHD;AD	BP: 3077F/; TOB:
01/27/11 G8402; BMI: 43.4; DEP: Meas PHQ2						

Figure 2-83: Sample Patient List: Comprehensive CVD-Related Assessment

2.8.10 Appropriate Medication Therapy after a Heart Attack

Denominator

Active Clinical patients aged 35 and older discharged for an AMI during the first 51 weeks of the Report Period and were not readmitted for any diagnosis within 7 days of discharge. Broken down by gender.

Numerators

Patients with active prescription for or who have a contraindication/previous adverse reaction to *beta-blockers*.

Note: This numerator does *not* include refusals.

Patients with active prescription for or who have a contraindication/previous adverse reaction to *ASA (aspirin) or other anti-platelet agent*.

Note: This numerator does *not* include refusals.

Patients with active prescription for or who have a contraindication/ previous adverse reaction to *ACEIs/ARBs*.

Note: This numerator does *not* include refusals.

Patients with active prescription for or who have a contraindication/ previous adverse reaction to *statins*.

Note: This numerator does *not* include refusals.

Also included for the numerators above are subnumerators:

- a. Patients with active prescription for the specified medication
- b. Patients with contraindication/previous adverse reaction to the specified medication

Patients with documented refusal of the specified medication

Patients with active prescriptions for *all post-AMI medications* (i.e., beta-blocker, ASA/anti-platelet, ACEI/ARB, and statin), and/or who have a contraindication/previous adverse reaction.

Note: This numerator does *not* include refusals.

Logic Description

Age is calculated at the beginning of the Report Period.

Acute Myocardial Infarction (AMI) definition: POV 410.*1 (i.e., first eligible episode of an AMI) with Service Category H. If patient has more than one episode of AMI during the first 51 weeks of the Report Period, CRS will include only the first discharge.

Denominator Exclusions

Patients meeting any of the following conditions will be excluded from the denominator.

1. Patients with Discharge Type of Irregular (AMA), Transferred, or contains "Death."
2. Patients readmitted for any diagnosis within seven days of discharge.
3. Patients with a Diagnosis Modifier of C (Consider), D (Doubtful), M (Maybe, Possible, Perhaps), O (Rule Out), P (Probable), R (Resolved), S (Suspect, Suspicious), or T (Status Post).
4. Patients with a Provider Narrative beginning with "Consider"; "Doubtful"; "Maybe"; "Possible"; "Perhaps"; "Rule Out"; "R/O"; "Probable"; "Resolved"; "Suspect"; "Suspicious"; or "Status Post."

Numerator Logic

In the logic below, "ever" is defined as anytime through the end of the Report Period.

To be included in the numerators, a patient must meet one of the three conditions below:

1. An active prescription (not discontinued as of [discharge date + 7 days] and does not have a comment of RETURNED TO STOCK) that was prescribed prior to admission, during the inpatient stay, or within seven days after discharge. "Active" prescription defined as: Days Prescribed > ((Discharge Date + 7 days) - Order Date); *or*
2. A refusal of the medication at least once during hospital stay through seven days after discharge date; *or...*
3. Have a contraindication/previous adverse reaction to the indicated medication.

Refusals and contraindications/previous adverse drug reactions (ADR)/allergies are only counted if a patient did not have a prescription for the indicated medication. Patients without a prescription who have a refusal and/or contraindication/ADR/allergy will be counted in subnumerators B–C. Because a patient may have both a refusal and a contraindication/ADR/allergy, the subnumerator totals of A–C may not add up to the numerator total.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Beta-Blocker Numerator Logic

Beta-blocker medication codes defined with medication taxonomy BGP HEDIS BETA BLOCKER MEDS. (Medications are: Noncardioselective Beta Blockers: Carteolol, Carvedilol, Labetalol, Nadolol, Penbutolol, Pindolol, Propranolol, Timolol, Sotalol; Cardioselective Beta Blockers: Acebutolol, Atenolol, Betaxolol, Bisoprolol, Metoprolol, Nebivolol; and Antihypertensive Combinations: Atenolol-chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Hydrochlorothiazide-metoprolol, and Hydrochlorothiazide-propranolol.)

Refusal of beta-blocker: REF refusal of any beta-blocker medication in site-populated medication taxonomy BGP HEDIS BETA BLOCKER MEDS at least once during hospital stay through seven days after discharge date.

CRS uses the following codes to define contraindications to beta-blockers.

Contraindication to Beta-Blockers (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Asthma		POV: 2 diagnoses of 493* on different visit dates
Hypotension		POV: 1 diagnosis of 458*
Heart Block >1 Degree		POV: 1 diagnosis of 426.0, 426.12, 426.13, 426.2, 426.3, 426.4, 426.51, 426.52, 426.53, 426.54, or 426.7
Sinus Bradycardia		POV: 1 diagnosis of 427.81
COPD		POV: 2 diagnoses on different visit dates of 491.2*, 496, or 506.4, or a combination of any of these codes, such as 1 visit with 491.20 and 1 with 496
NMI Refusal	G8011 (old code) at least once during hospital stay through 7 days after discharge date	Refusal: NMI (not medically indicated) refusal for any beta-blocker at least once during hospital stay through 7 days after discharge date

CRS uses the following codes to define adverse drug reactions/documentated allergies to beta-blockers.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to Beta-Blockers (any of the codes occurring ever)	POV: 995.0-995.3 AND E942.0
	Entry in ART (Patient Allergies File): “beta block*”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “beta block*”, “bblock*” or “b block*”

ASA (aspirin)/Other Anti-Platelet Numerator Logic

ASA medication codes defined with medication taxonomy DM AUDIT ASPIRIN DRUGS.

Other anti-platelet medication codes defined with medication taxonomy site-populated DM AUDIT ANTI-PLATELET DRUGS taxonomy.

Refusal of ASA/other anti-platelet: REF refusal of any ASA or anti-platelet medication in site-populated medication taxonomies DM AUDIT ASPIRIN DRUGS or BGP ANTI-PLATELET DRUGS at least once during hospital stay through seven days after discharge date.

CRS uses the following codes to define contraindications to ASA/other anti-platelets.

Contraindication to ASA/Other Anti-Platelets (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Active prescription for Warfarin/Coumadin at time of arrival or prescribed at discharge		Site-Populated Drug Taxonomy: BGP CMS WARFARIN MEDS
Hemorrhage		POV: 459.0
NMI Refusal	G8008 (old code) at least once during hospital stay through 7 days after discharge date	Refusal: NMI (not medically indicated) refusal for any aspirin at least once during hospital stay through 7 days after discharge date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ASA/other anti-platelets.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to ASA/Other Anti-Platelets (any of the codes occurring ever)	POV: 995.0-995.3 AND E935.3
	Entry in ART (Patient Allergies File): “aspirin”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “ASA” or “aspirin”

ACEI/ARB Numerator Logic

Ace Inhibitor (ACEI) medication codes defined with medication taxonomy BGP HEDIS ACEI MEDS. ACEI medications are: Angiotensin Converting Enzyme Inhibitors (Benazepril, Captopril, Enalapril, Fosinopril, Lisinopril, Moexipril, Perindopril, Quinapril, Ramipril, Trandolopril).

Antihypertensive Combinations (Amlodipine-benazepril, Benazepril-hydrochlorothiazide, Captopril-hydrochlorothiazide, Enalapril-hydrochlorothiazide, Fosinopril-hydrochlorothiazide, Hydrochlorothiazide-lisinopril, Hydrochlorothiazide-moexipril, Hydrochlorothiazide-quinapril, Trandolapril-verapamil).

Refusal of ACEI: REF refusal of any ACE inhibitor medication in site-populated medication taxonomy BGP HEDIS ACEI MEDS at least once during hospital stay through seven days after discharge date.

CRS uses the following codes to define contraindications to ACE inhibitors.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0-V23.9, V72.42, 640.*-649.*, or 651.*-676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI (not medically indicated) refusal for any ACE inhibitor at least once during hospital stay through 7 days after discharge date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ACE inhibitors.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ACE Inhibitors (any of the codes occurring ever)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): "ace inhibitor" or "ACEI"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "ace i*" or "ACEI"

ARB (Angiotensin Receptor Blocker) medication codes defined with medication taxonomy BGP HEDIS ARB MEDS. ARB medications: Angiotensin II Inhibitors (Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan.)

Antihypertensive Combinations (Aliskiren-valsartan, Amlodipine-hydrochlorothiazide-valsartan, Amlodipine-olmesartan, Candesartan-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Hydrochlorothiazide-Irbesartan, Hydrochlorothiazide-Losartan, Hydrochlorothiazide-olmesartan, Hydrochlorothiazide-Telmisartan, Hydrochlorothiazide-Valsartan).

Refusal of ARB: REF refusal of any ARB medication in site-populated medication taxonomy BGP HEDIS ARB MEDS at least once during hospital stay through seven days after discharge date.

CRS uses the following codes to define contraindications to ARBs.

Contraindication to ARBs (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0-V23.9, V72.42, 640.*-649.*, or 651.*-676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI refusal for any ARB at least once during hospital stay through 7 days after discharge date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ARBs.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to ARBs (any of the codes occurring ever)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): "Angiotensin Receptor Blocker" or "ARB"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "Angiotensin Receptor Blocker" or "ARB"

Statins Numerator Logic

Statin medication codes defined with medication taxonomy BGP HEDIS STATIN MEDS. Statin medications: Atorvastatin (Lipitor), Fluvastatin (Lescol), Lovastatin (Altacor), Mevacor, Pravastatin (Pravachol), Simvastatin (Zocor), Rosuvastatin (Crestor).

Statin Combination Products: Advicor, Caduet, PraviGard Pac, Vytorin.

Refusal of Statin: REF refusal of any statin medication in site-populated medication taxonomy BGP HEDIS STATIN MEDS at least once during hospital stay through seven days after discharge date.

CRS uses the following codes to define contraindications to statins.

Contraindication to Statins (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0–V23.9, V72.42, 640.*–649.*, or 651.*–676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period
Acute Alcoholic Hepatitis		POV: 571.1 during the Report Period
NMI Refusal		Refusal: NMI refusal for any statin at least once during hospital stay through 7 days after discharge date

CRS uses the following codes to define adverse drug reactions/documentated allergies to statins.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to Statins (any of the codes occurring ever unless otherwise noted)	Site-Populated Lab Taxonomy or LOINC Taxonomy: DM AUDIT ALT TAX, with ALT and/or AST > 3x the Upper Limit of Normal (ULN) (i.e., Reference High) on 2 or more consecutive visits during the Report Period
	Site-Populated Lab Taxonomy or LOINC Taxonomy: BGP CREATINE KINASE TAX, with Creatine Kinase (CK) levels > 10x ULN or CK > 10,000 IU/L during the Report Period
	POV: Myopathy/Myalgia, defined as any of the following during the Report Period: 359.0-359.9, 729.1, 710.5, or 074.1
	POV: 995.0-995.3 AND E942.9
	Entry in ART (Patient Allergies File): "statin" or "statins"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "Statin" or "Statins"

All Medications Numerator Logic

To be included in this numerator, a patient must have a prescription, refusal, or a contraindication for *all* of the four medication classes (i.e., beta-blocker, ASA/other anti-platelet, ACEI/ARB, *and* statin).

Key Logic Changes from CRS Version 11.0

1. Removed refusals from main numerators. Refusal numerator is a separate numerator, and no longer a subnumerator.
2. Added codes 16325-3, 48136-6, and 54500-4 to LOINC taxonomy for AST.
3. For contraindications to beta-blockers, noted that G8011 is an old code.
4. For contraindications to ASA, noted that G8008 is an old code.

Patient List Description

List of patients with AMI, with appropriate medication therapy, if any.

Measure Source

American Heart Association/American College of Cardiology Guidelines for the Treatment of AMI.

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011	Page 216
----	--------------	----------

*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Appropriate Medication Therapy after a Heart Attack									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR	% PERIOD		BASE %	
Active Clinical Pts 35+ hospitalized for AMI	71		0			0			
# w/beta-blocker Rx/contr/ADR									
-No Refusals	27	38.0	0	0.0	+38.0	0	0.0	+38.0	
A. # w/beta-blocker Rx w/ % of Total	5	18.5	0	0.0	+18.5	0	0.0	+18.5	
B. # w/contr/ADR w/ % of Total	22	81.5	0	0.0	+81.5	0	0.0	+81.5	
# w/ beta-blocker refusal	2	2.8	0	0.0	+2.8	0	0.0	+2.8	
# w/ASA Rx/contr/ADR									
-No Refusals	9	12.7	0	0.0	+12.7	0	0.0	+12.7	
A. # w/ASA Rx w/% of Total	3	33.3	0	0.0	+33.3	0	0.0	+33.3	
B. # w/contr/ADR w/ % of Total	6	66.7	0	0.0	+66.7	0	0.0	+66.7	
# w/ ASA refusal	3	4.2	0	0.0	+4.2	0	0.0	+4.2	
# w/ACEI/ARB Rx/contr/ADR									
-No Refusals	19	26.8	0	0.0	+26.8	0	0.0	+26.8	
A. # w/ACEI/ARB Rx w/% of Total	4	21.1	0	0.0	+21.1	0	0.0	+21.1	
B. # w/contr/ADR w/ % of Total	15	78.9	0	0.0	+78.9	0	0.0	+78.9	
# w/ ACEI/ARB refusal	2	2.8	0	0.0	+2.8	0	0.0	+2.8	
# w/statin Rx/contr/ADR									
-No Refusals	17	23.9	0	0.0	+23.9	0	0.0	+23.9	
A. # w/statin Rx w/% of Total	5	29.4	0	0.0	+29.4	0	0.0	+29.4	
B. # w/contr/ADR w/ % of Total	12	70.6	0	0.0	+70.6	0	0.0	+70.6	
# w/ Statin Refusal	2	2.8	0	0.0	+2.8	0	0.0	+2.8	
# w/Rx/contr/ADR of ALL meds-No Refusals	4	5.6	0	0.0	+5.6	0	0.0	+5.6	

Figure 2-84: Sample Report, Appropriate Medication Therapy after a Heart Attack

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Appropriate Medication Therapy after a Heart Attack: List of patients with AMI, with appropriate medication therapy, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,CECELIA	000001	COMMUNITY #1	F	37	AC	ACEI/ARB: Contra
ACEI - pregnant; STATIN: contra statin - pregnant						
PATIENT2,KATHLEEN	000002	COMMUNITY #1	F	38	AC	ASA: ADR 06/01/11
POV 995.3; ACEI/ARB: Contra POV [396.0]; STATIN: Statin contra - BF-HC						
PATIENT3,KIMBERLY A	000003	COMMUNITY #1	F	49	AC	BETA: 03/19/11
Refused PROPRANOLOL 10MG TAB; ASA: ADR 02/03/03 POV 995.3 w/E935.3						
PATIENT4,TIMOTHY JOHN	000004	COMMUNITY #1	M	57	AC	BETA: heart blk
contra;						
PATIENT5,FELIPE	000005	COMMUNITY #1	M	57	AC	
PATIENT6,JAMES DALTON	000006	COMMUNITY #1	M	77	AC	ALL MEDS: BETA:
04/23/11 Contra CPT code G8011; ASA: 04/23/11 Contra CPT code G8008; ACEI/ARB:						
04/27/11 Contra POV [396.0]; STATIN: 05/05/11 SIMVASTATIN 40MG TAB						

Figure 2-85: Sample Patient List: Appropriate Medication Therapy after a Heart Attack

2.8.11 Persistence of Appropriate Medication Therapy after a Heart Attack

Denominator

Active Clinical patients 35 and older diagnosed with an AMI six months prior to the Report Period through the first six months of the Report Period. Broken down by gender.

Numerators

Patients with a 135-day course of treatment with beta-blockers, or who have a contraindication/previous adverse reaction to beta-blocker therapy.

Note: This numerator does *not* include refusals.

Patients with a 135-day course of treatment with ASA (aspirin) or other anti-platelet agent or who have a contraindication/previous adverse reaction to ASA/anti-platelet therapy.

Note: This numerator does *not* include refusals.

Patients with a 135-day course of treatment with ACEIs/ARBs or who have a contraindication/previous adverse reaction to ACEI/ARB therapy.

Note: This numerator does *not* include refusals.

Patients with a *135-day course of treatment with statins* or who have a contraindication/previous adverse reaction to statin therapy.

Note: This numerator does *not* include refusals.

Also included for the numerators above are subnumerators:

- a. Patients with active prescription for the specified medication
- b. Patients with contraindication/previous adverse reaction to the specified medication

Patients with documented refusal of the specified medication

Patients with a *135-day course of treatment for all post-AMI medications* (i.e., beta-blocker, ASA/anti-platelet, ACEI/ARB, *and* statin) following first discharge date or visit date, including previous active prescriptions; and/or who have a contraindication/previous adverse reaction.

Note: This numerator does *not* include refusals.

Logic Description

Age is calculated at the beginning of the Report Period.

Acute Myocardial Infarction (AMI) definition: POV 410.0*–410.9* or 412. AMI diagnosis may be made at an inpatient or outpatient visit but must occur between six months prior to beginning of Report Period through first six months of the Report Period. Inpatient visit defined as Service Category of H (Hospitalization). If patient has more than one episode of AMI during the timeframe, CRS will include only the first hospital discharge or ambulatory visit.

Denominator Exclusions

Patients meeting any of the following conditions will be excluded from the denominator.

- If inpatient visit, patients with Discharge Type of Irregular (AMA), Transferred, or contains “Death.”
- Patients with a Diagnosis Modifier of C (Consider), D (Doubtful), M (Maybe, Possible, Perhaps), O (Rule Out), P (Probable), R (Resolved), S (Suspect, Suspicious), or T (Status Post).
- Patients with a Provider Narrative beginning with “Consider,” “Doubtful,” “Maybe,” “Possible,” “Perhaps,” “Rule Out,” “R/O,” “Probable,” “Resolved,” “Suspect,” “Suspicious,” or “Status Post.”

Numerator Logic

In the logic below, “ever” is defined as anytime through the end of the Report Period.

To be included in the numerators, a patient must meet one of the three conditions below.

1. A total days' supply ≥ 135 days in the 180 days following discharge date for inpatient visits or visit date for ambulatory visits. Medications must not have a comment of RETURNED TO STOCK. Prior active prescriptions can be included if the treatment days fall within the 180 days following discharge/visit date. Prior active prescription defined as most recent prescription (see codes below) prior to admission/visit date with the number of days supply equal to or greater than the discharge/visit date minus the prescription date; *or*
2. A refusal of the medication at least once at time of diagnosis through the 180 days after AMI; *or*
3. Have a contraindication/previous adverse reaction to the indicated medication.
4. Refusals and contraindications/previous ADR/allergies are only counted if a patient did not have a prescription for the indicated medication. Patients without a prescription who have a refusal and/or contraindication/ADR/allergy will be counted in subnumerators B–C. Because a patient may have both a refusal and a contraindication/ADR/allergy, the sub-numerator totals of A–C may not add up to the numerator total.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Example of patient included in the beta-blocker numerator who has prior active prescription:

- Admission Date: 2/1/2011, Discharge Date: 2/15/2011
- Must have 135 days prescribed by 8/13/2011 (Discharge Date+180)
- Prior Beta-Blocker Rx Date: 1/15/2011
- # Days Prescribed: 60 (treats patient through 3/15/2011)
- Discharge Date minus Rx Date: 2/15/2011-1/15/2011 = 31, 60 is ≥ 31 , prescription is considered Prior Active Rx
- 3/15/2011 is between 2/15 and 8/13/2011, thus remainder of Prior Active Rx can be counted toward 180-day treatment period

- # Remaining Days Prescribed from Prior Active Rx:
- $(60 - (\text{Discharge Date} - \text{Prior Rx Date})) = 60 - (2/15/2011 - 1/15/2011) = 60 - 31 = 29$
- Rx #2: 4/1/2011, # Days Prescribed: 90
- Rx #3: 7/10/2011, #Days Prescribed: 90
- Total Days Supply Prescribed between 2/15 and 8/13/2011: $29 + 90 + 90 = 209$

Beta-Blocker Numerator Logic

Beta-blocker medication codes defined with medication taxonomy BGP HEDIS BETA BLOCKER MEDS. (Medications are: Noncardioselective Beta Blockers: Carteolol, Carvedilol, Labetalol, Nadolol, Penbutolol, Pindolol, Propranolol, Timolol, Sotalol; Cardioselective Beta Blockers: Acebutolol, Atenolol, Betaxolol, Bisoprolol, Metoprolol, Nebivolol; and Antihypertensive Combinations: Atenolol-chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Hydrochlorothiazide-metoprolol and Hydrochlorothiazide-propranolol.)

Refusal of beta-blocker: REF refusal of any beta-blocker medication in site-populated medication taxonomy BGP HEDIS BETA BLOCKER MEDS at least once during the period admission/visit date through the 180 days after discharge/visit date.

CRS uses the following codes to define contraindications to beta-blockers.

Contraindication to Beta-Blockers (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
Asthma		POV: 2 diagnoses of 493* on different visit dates
Hypotension		POV: 1 diagnosis of 458*
Heart Block >1 Degree		POV: 1 diagnosis of 426.0, 426.12, 426.13, 426.2, 426.3, 426.4, 426.51, 426.52, 426.53, 426.54, or 426.7
Sinus Bradycardia		POV: 1 diagnosis of 427.81
COPD		POV: 2 diagnoses on different visit dates of 491.2*, 496, or 506.4, or a combination of any of these codes, such as 1 visit with 491.20 and 1 with 496
NMI Refusal	G8011 (old code) at least once during the period admission/visit date through the 180 days after discharge/visit date	Refusal: NMI refusal for any beta-blocker at least once during the period admission/visit date through the 180 days after discharge/visit date

CRS uses the following codes to define adverse drug reactions/documentated allergies to beta-blockers.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to Beta-Blockers (any of the codes occurring anytime up to the 180 days after discharge/visit date)	POV: 995.0-995.3 AND E942.0
	Entry in ART (Patient Allergies File): "beta block*"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "beta block*", "bblock*" or "b block*"

ASA (Aspirin)/Other Anti-Platelet Numerator Logic

ASA medication codes defined with medication taxonomy DM AUDIT ASPIRIN DRUGS.

Other anti-platelet medication codes defined with medication taxonomy site-populated BGP ANTI-PLATELET DRUGS taxonomy.

Refusal of ASA/other anti-platelet: REF refusal of any ASA or anti-platelet medication in site-populated medication taxonomies DM AUDIT ASPIRIN DRUGS or DM AUDIT ANTI-PLATELET DRUGS at least once during the period admission/visit date through the 180 days after discharge/visit date.

CRS uses the following codes to define contraindications to ASA/other anti-platelets.

Contraindication to ASA/Other Anti-Platelets (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
Active prescription for Warfarin/Coumadin during the period admission/visit date through the 180 days after discharge/visit date		Site-Populated Drug Taxonomy: BGP CMS WARFARIN MEDS
Hemorrhage		POV: 459.0
NMI Refusal	G8008 (old code) at least once during the period admission/visit date through the 180 days after discharge/visit date	Refusal: NMI (not medically indicated) refusal for any aspirin/other anti-platelet at least once during the period admission/visit date through the 180 days after discharge/visit date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ASA/other anti-platelets.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to ASA/Other Anti-Platelets (any of the codes occurring anytime up to the 180 days after discharge/visit date)	POV: 995.0-995.3 AND E935.3
	Entry in ART (Patient Allergies File): “aspirin”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “ASA” or “aspirin”

ACEI/ARB Numerator Logic

Ace Inhibitor (ACEI) medication codes defined with medication taxonomy BGP HEDIS ACEI MEDS. ACEI medications are: Angiotensin Converting Enzyme Inhibitors (Benazepril, Captopril, Enalapril, Fosinopril, Lisinopril, Moexipril, Perindopril, Quinapril, Ramipril, Trandolopril).

Antihypertensive Combinations (Amlodipine-benazepril, Benazepril-hydrochlorothiazide, Captopril-hydrochlorothiazide, Enalapril-hydrochlorothiazide, Fosinopril-hydrochlorothiazide, Hydrochlorothiazide-lisinopril, Hydrochlorothiazide-moexipril, Hydrochlorothiazide-quinapril, Trandolapril-verapamil).

Refusal of ACEI: REF refusal of any ACE Inhibitor medication in site-populated medication taxonomy BGP HEDIS ACEI MEDS at least once during the period admission/visit date through the 180 days after discharge/visit date.

CRS uses the following codes to define contraindications to ACE inhibitors.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the period admission/visit date through the 180 days after discharge/visit date with V22.0-V23.9, V72.42, 640.*-649.*, or 651.*-676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the period, CRS will use the first two visits in the period.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the period admission/visit date through the 180 days after discharge/visit date Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the period admission/visit date through the 180 days after discharge/visit date
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI refusal for any ACE inhibitor at least once during the period admission/visit date through the 180 days after discharge/visit date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ACE inhibitors.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ACE Inhibitors (any of the codes occurring anytime up to the 180 days after discharge/visit date)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): "ace inhibitor" or "ACEI"
	Contained within or Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "ace i*" or "ACEI"

ARB (Angiotensin Receptor Blocker) medication codes defined with medication taxonomy BGP HEDIS ARB MEDS. ARB medications are: Angiotensin II Inhibitors (Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan.).

Antihypertensive Combinations (Aliskiren-valsartan, Amlodipine-hydrochlorothiazide-valsartan, Amlodipine-olmesartan, Candesartan-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Hydrochlorothiazide-Irbesartan, Hydrochlorothiazide-Losartan, Hydrochlorothiazide-olmesartan, Hydrochlorothiazide-Telmisartan, Hydrochlorothiazide-Valsartan).

Refusal of ARB: REF refusal of any ARB medication in site-populated medication taxonomy BGP HEDIS ARB MEDS at least once during the period admission/visit date through the 180 days after discharge/visit date.

CRS uses the following codes to define contraindications to ARBs.

Contraindication to ARBs (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the period admission/visit date through the 180 days after discharge/visit date with V22.0-V23.9, V72.42, 640.* –649.*, or 651.*–676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the period, CRS will use the first two visits in the period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the period admission/visit date through the 180 days after discharge/visit date Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the period admission/visit date through the 180 days after discharge/visit date
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22

Contraindication to ARBs (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
NMI Refusal		Refusal: NMI refusal for any ARB at least once during the period admission/visit date through the 180 days after discharge/visit date

CRS uses the following codes to define adverse drug reactions/documentated allergies to ARBs.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ARBs (any of the codes occurring anytime up to the 180 days after discharge/visit date)	POV: 995.0-995.3 AND E942.6 Entry in ART (Patient Allergies File): "Angiotensin Receptor Blocker" or "ARB" Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "Angiotensin Receptor Blocker" or "ARB"

Statins Numerator Logic

Statin medication codes defined with medication taxonomy BGP HEDIS STATIN MEDS. Statin medications: Atorvastatin (Lipitor), Fluvastatin (Lescol), Lovastatin (Altacor), Mevacor, Pravastatin (Pravachol), Simvastatin (Zocor), Rosuvastatin (Crestor).

Statin Combination Products: Advicor, Caduet, PraviGard Pac, Vytarin.

Refusal of Statin: REF refusal of any statin medication in site-populated medication taxonomy BGP HEDIS STATIN MEDS at least once during admission/visit date through the 180 days after discharge/visit date.

CRS uses the following codes to define contraindications to statins.

Contraindication to Statins (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the period admission/visit date through the 180 days after discharge/visit date with V22.0–V23.9, V72.42, 640.*-649.*, or 651.* –676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the period, CRS will use the first two visits in the period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedure: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the period admission/visit date through the 180 days after discharge/visit date Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the period admission/visit date through the 180 days after discharge/visit date
Acute Alcoholic Hepatitis		POV: 571.1 during the period admission/visit date through the 180 days after discharge/visit date
NMI Refusal		Refusal: NMI refusal for any statin at least once during the period admission/visit date through the 180 days after discharge/visit date

CRS uses the following codes to define adverse drug reactions/documentated allergies to statins.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to Statins (any of the codes occurring anytime up to the 180 days after discharge/visit date unless otherwise noted)	Site-Populated Lab Taxonomy or LOINC Taxonomy: DM AUDIT ALT TAX, with ALT and/or AST > 3x the Upper Limit of Normal (ULN) (i.e., Reference High) on 2 or more consecutive visits during the period admission/visit date through the 180 days after discharge/visit date
	Site-Populated Lab Taxonomy or LOINC Taxonomy: BGP CREATINE KINASE TAX, with Creatine Kinase (CK) levels > 10x ULN or CK > 10,000 IU/L during the period admission/visit date through the 180 days after discharge/visit date
	POV: Myopathy/Myalgia, defined as any of the following during the period admission/visit date through the 180 days after discharge/visit date: 359.0–359.9, 729.1, 710.5, or 074.1
	POV: 995.0–995.3 AND E942.9
	Entry in ART (Patient Allergies File): “statin” or “statins”
	Entry in Problem List or in Provider Narrative for any POV 995.0–995.3 or V14.8: “Statin” or “Statins”

All Medications Numerator Logic

To be included in this numerator, a patient must have a prescription, refusal, or a contraindication for *all* of the four medication classes (i.e., beta-blocker, ASA/other anti-platelet, ACEI/ARB, *and* statin).

Key Logic Changes from CRS Version 11.0

1. Removed refusals from main numerators. Refusal numerator is a separate numerator, and no longer a subnumerator.
2. Added codes 16325-3, 48136-6, and 54500-4 to LOINC taxonomy for AST.
3. For contraindications to beta-blockers, noted that G8011 is an old code.
4. For contraindications to ASA, noted that G8008 is an old code.

Patient List Description

List of patients with AMI, with persistent medication therapy, if any.

Measure Source

American Heart Association/American College of Cardiology Guidelines for the Treatment of AMI.

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 228

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Persistence of Appropriate Medication Therapy after a Heart Attack

	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR	BASE % PERIOD	%	CHG from BASE %
Active Clinical Pts 35+ w/ AMI DX	61		4			4		
# w/135-day beta-blocker Rx/contra/ADR								
-No Refusals	23	37.7	2	50.0	-12.3	3	75.0	-37.3
A. # w/135-day beta blocker Rx w/ % of Total	3	13.0	2	100.0	-87.0	2	66.7	-53.6
B. # w/contra/ADR w/ % of Total	20	87.0	0	0.0	+87.0	1	33.3	+53.6
# w/ beta-blocker refusal	1	1.6	0	0.0	+1.6	0	0.0	+1.6
# w/135-day ASA Rx/contra/ADR								
-No Refusals	5	8.2	0	0.0	+8.2	2	50.0	-41.8
A. # w/135-day ASA Rx w/% of Total	1	20.0	0	0.0	+20.0	2	100.0	-80.0
B. # w/contra/ADR w/ % of Total	4	80.0	0	0.0	+80.0	0	0.0	+80.0
# w/ ASA refusal	2	3.3	0	0.0	+3.3	0	0.0	+3.3
# w/135-day ACEI/ARB Rx/contra/ADR								
-No Refusals	13	21.3	1	25.0	-3.7	1	25.0	-3.7
A. # w/135-day ACEI/ARB Rx w/% of Total	1	7.7	1	100.0	-92.3	1	100.0	-92.3
B. # w/contra/ADR w/ % of Total	12	92.3	0	0.0	+92.3	0	0.0	+92.3
# w/ ACEI/ARB refusal	1	1.6	0	0.0	+1.6	0	0.0	+1.6
# w/135-day statin Rx/contra/ADR								
-No Refusals	6	9.8	2	50.0	-40.2	2	50.0	-40.2
A. # w/135-day statin Rx w/% of Total	2	33.3	2	100.0	-66.7	2	100.0	-66.7
B. # w/contra/ADR w/ % of Total	4	66.7	0	0.0	+66.7	0	0.0	+66.7
# w/ Statin refusal	2	3.3	0	0.0	+3.3	0	0.0	+3.3
# w/Rx/contra/ADR of ALL								

meds-No Refusals	2	3.3	0	0.0	+3.3	1	25.0	-21.7
------------------	---	-----	---	-----	------	---	------	-------

Figure 2-86: Sample Report, Persistence of Appropriate Medication Therapy after a Heart Attack

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Persistence of Appropriate Medication Therapy after a Heart Attack: List of patients with AMI, with persistent medication therapy, if any							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT1,RHONDA	000001	COMMUNITY #1	F	35	AC	ALL MEDS: BETA:	
03/07/09 03/13/09 Contra 2 dx asthma; ASA: 03/23/11 Contra NMI ASPIRIN 325MG CAP; ACEI/ARB: 04/01/11 ADR Problem List 995.0 ACEI; STATIN: 04/01/11 ADR creat kinase of 5000							
PATIENT2,KATHLEEN	000002	COMMUNITY #1	F	38	AC	BETA: 04/30/11	
Contra 2/3 heart block dx 426.3							
PATIENT3,KIMBERLY A	000003	COMMUNITY #1	F	49	AC	BETA: 05/01/11	
Refused PROPRANOLOL 10MG TAB; ASA: 05/01/11 Refused CLOPIDOGREL 37.5MG TAB U/D; ACEI/ARB: 05/02/11 Refused IRBESARTAN 150MG TAB U/D							
PATIENT4,TIMOTHY	000004	COMMUNITY #1	M	57	AC	BETA: Beta Blocker	
contra NMI med 05/09/11; STATIN: adr Statin - AST/ALT							
PATIENT5,JOSHUA	000005	COMMUNITY #1	M	63	AC	BETA: 01/15/11	
Contra sinus brady dx 427.81							

Figure 2-87: Sample Patient List: Persistence of Appropriate Medication Therapy after a Heart Attack

2.8.12 Appropriate Medication Therapy in High Risk Patients

Denominators

Active IHD patients ages 22 and older; defined as all Active Clinical patients diagnosed with IHD prior to the Report Period, *and* at least two visits during the Report Period, *and* two IHD-related visits ever.

- Active IHD patients ages 22 and older who are not Active Diabetic.
- Active IHD patients ages 22 and older who are Active Diabetic

Numerators

Patients with a 180-day course of treatment with *beta-blockers* during the Report Period, or who have a contraindication/previous adverse reaction to beta-blocker therapy.

Note: This numerator does *not* include refusals.

Patients with a 180-day course of treatment with *ASA (aspirin) or other anti-platelet agent* during the Report Period, or who have a contraindication/previous adverse reaction to ASA/anti-platelet therapy.

Note: This numerator does *not* include refusals.

Patients with a 180-day course of treatment with *ACEIs/ARBs* during the Report Period, or who have a contraindication/previous adverse reaction to ACEI/ARB therapy.

Note: This numerator does *not* include refusals.

Patients with a 180-day course of treatment with *statins* during the Report Period, or who have a contraindication/previous adverse reaction to statin therapy.

Note: This numerator does *not* include refusals.

Also included for the numerators above are subnumerators:

- a. Patients with active prescription for the specified medication
- b. Patients with contraindication/previous adverse reaction to the specified medication

Patients with documented refusal of the specified medication

Patients with a 180-day course of treatment for *all medications* (i.e., beta-blocker, aspirin/anti-platelet, ACEI/ARB, *and* statin) during the Report Period, and/or who have a contraindication/previous adverse reaction.

Note: This numerator does *not* include refusals.

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

IHD diagnosis defined as: 410.0–412.*, 414.0–414.9, or 429.2 recorded in the V POV file.

Diabetes defined as: Diagnosed with diabetes (first POV in V POV with 250.00–250.93) prior to the Current Report Period, *and* at least two visits during the Current Report period, *and* two DM-related visits ever. Patients not meeting these criteria are considered non-diabetics.

Numerator Logic

In the logic below, “ever” is defined as anytime through the end of the Report Period.

To be included in the numerators, a patient must meet one of the three conditions below:

1. Prescription(s) for the indicated medication with a total days supply of 180 days or more during the Report Period. Medications must not have a comment of RETURNED TO STOCK; *or*
2. A refusal of the medication during the Report Period; *or*
3. Have a contraindication/previous adverse reaction to the indicated medication

Refusals and contraindications/previous ADR/allergies are only counted if a patient did not have a prescription for the indicated medication. Patients without a prescription who have a refusal and/or contraindication/ADR/ allergy will be counted in sub-numerators B–C. Because a patient may have both a refusal and a contraindication/ADR/allergy, the sub-numerator totals of A–C may not add up to the numerator total.

For prescriptions, the days supply requirement may be met with a single prescription or from a combination of prescriptions for the indicated medication that were filled during the Report Period and prescriptions filled prior to the Report Period but which are still active (i.e., prior active prescription). Prior active prescriptions can be included if the treatment days fall within the Report Period. Prior active prescription defined as most recent prescription for the indicated medication (see codes below) prior to Report Period Start Date with the number of days supply equal to or greater than the Report Period Start Date minus the prescription date.

Note: If a prescription for a medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Example of patient included in the beta-blocker numerator with prior active prescription:

- Report Period: 07/01/2010–06/30/2011
- Must have 180 days supply of indicated medication 6/30/2011 (end of Report Period)
- Prior Beta-Blocker Rx Date: 06/01/2010
- # Days Prescribed: 60 (treats patient through 07/31/2010)
- Report Period Start Date minus Rx Date: 07/01/2010-06/01/2010 = 30; 60 (#Days Prescribed) is ≥ 30 , prescription is considered Prior Active Rx
- 07/31/2010 is between the Report Period of 07/01/2010 and 06/30/2011, thus remainder of Prior Active Rx can be counted toward 180-days supply

- # Remaining Days Prescribed from Prior Active Rx:
- (# Days Prescribed-(Report Period Start Date-Prior Rx Date) =
60-(07/01/2010-06/01/2010) = 60-30 = 30
- Rx #2: 08/05/2010, # Days Prescribed: 90
- Rx #3: 11/10/2010, #Days Prescribed: 90
- Total Days Supply Prescribed between 07/01/2010 and 06/30/2011, including prior active prescription: 30+90+90=210

Beta-Blocker Numerator Logic

Beta-blocker medication codes defined with medication taxonomy BGP HEDIS BETA BLOCKER MEDS. Medications are: Noncardioselective Beta Blockers: Carteolol, Carvedilol, Labetalol, Nadolol, Penbutolol, Pindolol, Propranolol, Timolol, Sotalol; Cardioselective Beta Blockers: Acebutolol, Atenolol, Betaxolol, Bisoprolol, Metoprolol, Nebivolol; and Antihypertensive Combinations: Atenolol-chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Hydrochlorothiazide-metoprolol, and Hydrochlorothiazide-propranolol.

Refusal of beta-blocker: REF refusal of any beta-blocker medication in site-populated medication taxonomy BGP HEDIS BETA BLOCKER MEDS at least once during the Report Period.

CRS uses the following codes to define contraindications to beta-blockers.

Contraindication to Beta-Blockers (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
Asthma		POV: 2 diagnoses of 493* on different visit dates
Hypotension		POV: 1 diagnosis of 458*
Heart Block >1 Degree		POV: 1 diagnosis of 426.0, 426.12, 426.13, 426.2, 426.3, 426.4, 426.51, 426.52, 426.53, 426.54, or 426.7
Sinus Bradycardia		POV: 1 diagnosis of 427.81
COPD		POV: 2 diagnoses on different visit dates of 491.2*, 496, or 506.4, or a combination of any of these codes, such as 1 visit with 491.20 and 1 with 496
NMI Refusal	G8011 (old code) at least once during the Report Period	Refusal: NMI refusal for any beta-blocker at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to beta-blockers.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to Beta-Blockers (any of the codes occurring ever)	POV: 995.0-995.3 AND E942.0
	Entry in ART (Patient Allergies File): "beta block**"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "beta block**", "bblock*" or "b block**"

ASA (aspirin)/Other Anti-Platelet Numerator Logic

ASA medication codes defined with medication taxonomy DM AUDIT ASPIRIN DRUGS.

Other anti-platelet medication codes defined with medication taxonomy site-populated BGP ANTI-PLATELET DRUGS taxonomy.

Refusal of ASA/other anti-platelet: REF refusal of any ASA or anti-platelet medication in site-populated medication taxonomies DM AUDIT ASPIRIN DRUGS or BGP ANTI-PLATELET DRUGS at least once during the Report Period.

CRS uses the following codes to define contraindications to ASA/other anti-platelets.

Contraindication to ASA/Other Anti-Platelets (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
180-day course of treatment for Warfarin/Coumadin during the Report Period		Site-Populated Drug Taxonomy: BGP CMS WARFARIN MEDS
Hemorrhage		POV: 459.0
NMI Refusal	G8008 (old code) at least once during the Report Period	Refusal: NMI (not medically indicated) refusal for any aspirin at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to ASA/other anti-platelets.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to ASA/Other Anti-Platelets (any of the codes occurring ever)	POV: 995.0-995.3 AND E935.3
	Entry in ART (Patient Allergies File): "aspirin"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "ASA" or "aspirin"

ACEI/ARB Numerator Logic

Ace Inhibitor (ACEI) medication codes defined with medication taxonomy BGP HEDIS ACEI MEDS. ACEI medications are: Angiotensin Converting Enzyme Inhibitors (Benazepril, Captopril, Enalapril, Fosinopril, Lisinopril, Moexipril, Perindopril, Quinapril, Ramipril, Trandolopril.)

Antihypertensive Combinations: (Amlodipine-benazepril, Benazepril-hydrochlorothiazide, Captopril-hydrochlorothiazide, Enalapril-hydrochlorothiazide, Fosinopril-hydrochlorothiazide, Hydrochlorothiazide-lisinopril, Hydrochlorothiazide-moexipril, Hydrochlorothiazide-quinapril, Trandolapril-verapamil.)

Refusal of ACEI: REF refusal of any ACE Inhibitor medication in site-populated medication taxonomy BGP HEDIS ACEI MEDS at least during the Report Period.

CRS uses the following codes to define contraindications to ACE inhibitors.

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0–V23.9, V72.42, 640.* –649.*, or 651.* –676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period

Contraindication to ACE Inhibitors (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI refusal for any ACE inhibitor at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to ACE inhibitors.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ACE Inhibitors (any of the codes occurring anytime through the end of the Report Period)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): "ace inhibitor" or "ACEI"
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: "ace i*" or "ACEI"

ARB (Angiotensin Receptor Blocker) medication codes defined with medication taxonomy BGP HEDIS ARB MEDS. ARB medications are: Angiotensin II Inhibitors (Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan.)

Antihypertensive Combinations (Aliskiren-valsartan, Amlodipine-hydrochlorothiazide-valsartan, Amlodipine-olmesartan, Candesartan-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Hydrochlorothiazide-Irbesartan, Hydrochlorothiazide-Losartan, Hydrochlorothiazide-olmesartan, Hydrochlorothiazide-Telmisartan, Hydrochlorothiazide-Valsartan.)

Refusal of ARB: REF refusal of any ARB medication in site-populated medication taxonomy BGP HEDIS ARB MEDS at least once during the Report Period.

CRS uses the following codes to define contraindications to ARBs.

Contraindication to ARBs (any of the codes occurring ever unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0–V23.9, V72.42, 640.*–649.*, or 651.*–676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedures: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period
Moderate or Severe Aortic Stenosis		POV: 395.0, 395.2, 396.0, 396.2, 396.8, 424.1, 425.1, or 747.22
NMI Refusal		Refusal: NMI (not medically indicated) refusal for any ARB at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to ARBs.

	ICD and Other Codes
Adverse Drug Reaction/ Allergy to ARBs (any of the codes occurring anytime through the end of the Report Period)	POV: 995.0-995.3 AND E942.6
	Entry in ART (Patient Allergies File): “Angiotensin Receptor Blocker” or “ARB”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “Angiotensin Receptor Blocker” or “ARB”

Statins Numerator Logic

Statin medication codes defined with medication taxonomy BGP HEDIS STATIN MEDS. Statin medications are: Atorvastatin (Lipitor), Fluvastatin (Lescol), Lovastatin (Altacor), Mevacor, Pravastatin (Pravachol), Simvastatin (Zocor), Rosuvastatin (Crestor).

Statin Combination Products: Advicor, Caduet, PraviGard Pac, Vytori.

Refusal of Statin: REF refusal of any statin medication in site-populated medication taxonomy BGP HEDIS STATIN MEDS at least once during the Report Period.

CRS uses the following codes to define contraindications to statins.

Contraindication to ACE Inhibitors (any of the codes occurring ever, unless otherwise noted)	CPT Codes	ICD and Other Codes
Pregnancy		POV or Problem List: At least two visits during the Report Period with V22.0–V23.9, V72.42, 640.*–649.*, 651.*–676.* and with no documented miscarriage or abortion (defined below) occurring after the second pregnancy POV. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the Report Period, CRS will use the first two visits in the Report Period.
	Abortion: Any of the following: 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260-S2267	Abortion: Any of the following POVs: 635*, 636*, or 637* Procedure: 69.01, 69.51, 74.91, 96.49
	Miscarriage: Any of the following: 59812, 59820, 59821, 59830	Miscarriage: Any of the following POVs: 630, 631, 632, 633*, or 634*
Breastfeeding		POV: V24.1 during the Report Period Patient Education: BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, or BF-N during the Report Period
Acute Alcoholic Hepatitis		POV: 571.1 during the Report Period
NMI Refusal		Refusal: NMI refusal for any statin at least once during the Report Period

CRS uses the following codes to define adverse drug reactions/documentated allergies to statins.

	ICD and Other Codes
Adverse Drug Reaction/Allergy to Statins (any of the codes occurring anytime through the end of the Report Period, unless otherwise noted)	Site-Populated Lab Taxonomy or LOINC Taxonomy: DM AUDIT ALT TAX, with ALT and/or AST > 3x the Upper Limit of Normal (ULN) (i.e., Reference High) on 2 or more consecutive visits during the Report Period
	Site-Populated Lab Taxonomy or LOINC Taxonomy: BGP CREATINE KINASE TAX, with Creatine Kinase (CK) levels > 10x ULN or CK > 10,000 IU/L during the Report Period
	POV: Myopathy/Myalgia, defined as any of the following during the Report Period: 359.0-359.9, 729.1, 710.5, or 074.1
	POV: 995.0-995.3 AND E942.9
	Entry in ART (Patient Allergies File): “statin” or “statins”
	Entry in Problem List or in Provider Narrative for any POV 995.0-995.3 or V14.8: “Statin” or “Statins”

All Medications Numerator Logic

To be included in this numerator, a patient must have a prescription, refusal, or a contraindication for *all* of the four medication classes (i.e., beta-blocker, ASA/other anti-platelet, ACEI/ARB, *and* statin).

Key Logic Changes from CRS Version 11.0

1. Removed refusals from main numerators. Refusal numerator is a separate numerator, and no longer a subnumerator.
2. Removed CPT code range 428.* from Ischemic Heart Disease definition.
3. Added codes 16325-3, 48136-6, and 54500-4 to LOINC taxonomy for AST.
4. For contraindications to beta-blockers, noted that G8011 is an old code.
5. For contraindications to ASA, noted that G8008 is an old code.

Patient List Description

List of IHD patients 22+ with 180-day medication therapy during the Report Period, if any.

Measure Source

American Heart Association/American College of Cardiology Guidelines

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011					Page 241		
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Appropriate Medication Therapy in High Risk Patients								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active IHD pts 22+	59		38			30		
# w/180 day beta-blocker								
Rx/contra/ADR								
-No Refusals	32	54.2	22	57.9	-3.7	14	46.7	+7.6
A. # w/180 day beta-blocker								
Rx w/% of Total	18	56.3	14	63.6	-7.4	11	78.6	-22.3
B: # w/contra/ADR								
w/ % of Total	14	43.8	8	36.4	+7.4	3	21.4	+22.3
# w/ beta-blocker refusal								
	1	1.7	0	0.0	+1.7	0	0.0	+1.7
# w/180 day ASA								
Rx/contra/ADR								
-No Refusals	22	37.3	21	55.3	-18.0	23	76.7	-39.4
A. # w/180 day ASA								
Rx w/% of Total	17	77.3	18	85.7	-8.4	18	78.3	-1.0
B. # w/contra/ADR								
w/ % of Total	5	22.7	3	14.3	+8.4	5	21.7	+1.0
# w/ ASA refusal								
	1	1.7	0	0.0	+1.7	0	0.0	+1.7
# w/180 day ACEI/ARB								
Rx/contra/ADR								
-No Refusals	29	49.2	15	39.5	+9.7	18	60.0	-10.8
A. # w/180 day ACEI/ARB								
Rx w/% of Total	26	89.7	14	93.3	-3.7	17	94.4	-4.8
B. # w/contra/ADR								
w/ % of Total	3	10.3	1	6.7	+3.7	1	5.6	+4.8
# w/ ACEI/ARB refusal								
	1	1.7	0	0.0	+1.7	0	0.0	+1.7
# w/180 day statin								
Rx/contra/ADR								
-No Refusals	30	50.8	22	57.9	-7.0	14	46.7	+4.2
A. # w/180 day statin								
Rx w/% of Total	26	86.7	20	90.9	-4.2	14	100.0	-13.3
B. # w/contra/ADR								
w/ % of Total	4	13.3	2	9.1	+4.2	0	0.0	+13.3
# w/ Statin refusal								
	1	1.7	0	0.0	+1.7	0	0.0	+1.7
# w/180 day Rx/contra/ADR/								
of ALL meds								
-No Refusals	16	27.1	8	21.1	+6.1	6	20.0	+7.1

Figure 2-88: Sample Report, Appropriate Medication Therapy in High Risk Patients

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Appropriate Medication Therapy in High Risk Patients: List of IHD patients 22+
 with 180-day medication therapy during the Report Period, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,CAITLYN	000001	COMMUNITY #1	F	22	IHD	BETA: 05/11/11
Refused METOPROLOL TARTRATE 50MG TAB; ASA: 05/11/11 Refused ASPIRIN 325MG E.C. TAB; ACEI/ARB: 05/11/11 Refused CAPTOPRIL 25MG TABS; STATIN: 05/11/11 Refused						
PATIENT2,CARLA	000002	COMMUNITY #1	F	40	IHD	
PATIENT3,GENEVA	000003	COMMUNITY #1	F	47	IHD,AD	BETA: 03/27/11
05/12/11 Contra 2 dx asthma						
PATIENT4,SHERRY LISA	000004	COMMUNITY #1	F	68	IHD	ASA: aspirin
contra total days WARFARIN: 676; ACEI/ARB: 01/06/11(99);04/15/11(42);05/27/11(41); 07/07/11(42);08/18/11(38);09/25/11(40);11/04/11(35);12/09/11(22) (359 TOTAL DAYS); STATIN: 01/06/11(99);04/15/11(42);05/27/11(41);07/07/11(42);08/18/11(38); 09/25/11(40);11/04/11(35);12/09/11(22) (359 TOTAL DAYS)						
PATIENT5,PAULINE	000005	COMMUNITY #1	F	70	IHD,AD	ALL MEDS: BETA:
07/23/11 Contra CPT G8011; ASA: 07/23/11 Contra CPT G8008; ACEI/ARB: 07/27/11 Contra POV 396.0; STATIN: 06/05/11						

Figure 2-89: Sample Patient List: Appropriate Medication Therapy in High Risk Patients

2.8.13 Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge

Denominator

Number of visits for *User Population patients* ages 18 and older who were hospitalized during the Report Period with ischemic stroke or transient ischemic attack (TIA) with documented permanent, persistent, or paroxysmal atrial fibrillation.

Numerators

Number of visits where patients received a prescription for anticoagulant at discharge

Number of visits where patients refused anticoagulant therapy

Number of visits where patients did not receive anticoagulation therapy

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

CRS uses the following codes to define ischemic stroke or transient ischemic attack with atrial fibrillation.

	ICD and Other Codes
Ischemic Stroke or TIA with Atrial Fibrillation (Non-CHS inpatient visit - Type not equal to C and Service Category=H) The patient must be admitted to the hospital during the report period with a condition described here but the discharge may occur after the report period.	V POV: 433.01, 433.11, 433.21, 433.31, 433.81, 433.91, 434.01, 434.11, 434.91, 435.0, 435.1, 435.2, 435.3, 435.8, or 435.9 AND 427.31 (atrial fibrillation)

Anticoagulant Therapy: Patient must meet one of the conditions below to be counted as receiving anticoagulant therapy. For all prescriptions, medications must not have a comment of RETURNED TO STOCK.

1. Active prescription for Warfarin, aspirin, or other anti-platelet as of discharge date. "Active" prescription defined as:

Rx Days Supply \geq (Discharge Date - Prescription Date), where the prescription has not been discontinued as of the discharge date.

2. Prescription for Warfarin, aspirin, or other anti-platelet on discharge date.

Warfarin Medication: Any medication in site-populated BGP CMS WARFARIN MEDS taxonomy.

Aspirin Medication: Any medication in site-populated DM AUDIT ASPIRIN DRUGS taxonomy.

Other Anti-Platelet/Anticoagulant Medication: Any medication in the site-populated BGP ANTI-PLATELET DRUGS taxonomy, any medication with VA Drug Class BL700.

Refusal of Anticoagulant Therapy: Refusal of any of the following documented on discharge date:

1. Any medication in site-populated taxonomies BGP CMS WARFARIN MEDS, DM AUDIT ASPIRIN DRUGS, or BGP ANTI-PLATELET DRUGS; or
2. Any medication with VA Drug Class BL700.

No Anticoagulant Therapy: Patients who did not have an active prescription for anticoagulant therapy at time of discharge and did not receive or refuse anticoagulant therapy at discharge.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with stroke/TIA and atrial fibrillation with anticoagulant therapy, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 248

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge

	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
# Stroke/TIA w/ Atrial Fib Visits for User								
Pop Pts 18+	17		0			0		
# Visits w/ Anticoagulant Rx	5	29.4	0	0.0	+29.4	0	0.0	+29.4
# Visits w/ Refusal	1	5.9	0	0.0	+5.9	0	0.0	+5.9
# Visits w/ No Anticoagulant Therapy	11	64.7	0	0.0	+64.7	0	0.0	+64.7

Figure 2-90: Sample Report, Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic						
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge: List of patients with stroke/TIA and atrial fibrillation with anticoagulant therapy, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,SHERRY	000001	COMMUNITY #1	F	38	UP; Visit: 1)	04/01/11
433.81+427.31	THERAPY: 04/01/11					
PATIENT2,CODY JACK	000002	COMMUNITY #1	M	31	UP; Visit: 1)	05/01/11
433.81+427.31	THERAPY: 05/01/11; 2) 09/01/11 433.21+427.31 THERAPY: 09/20/11; 3)					
10/15/11 434.01+427.31	THERAPY: 10/15/11					
PATIENT3,TIMOTHY ALLEN	000003	COMMUNITY #1	M	33	UP; Visit: 1)	11/20/11
434.11+427.31	THERAPY: 11/25/11					
PATIENT4,TRACE	000004	COMMUNITY #1	M	37	UP; Visit: 1)	05/01/11
433.21+427.31	THERAPY: 03/01/11					

Figure 2-91: Sample Patient List: Stroke and Stroke Rehabilitation: Anticoagulant Therapy Prescribed for Atrial Fibrillation at Discharge

2.8.14 Cholesterol Management for Patients with Cardiovascular Conditions

Denominators

Active Clinical patients ages 18 to 75 who, during the first ten months of the year prior to the beginning of the Report Period, were diagnosed with AMI, coronary artery bypass graft (CABG), or percutaneous coronary interventions (PCI) *or* who were diagnosed with IVD during the Report Period and the year prior to the Report Period. Broken down by gender.

User Population patients ages 18 to 75 who, during the first 10 months of the year prior to the beginning of the Report period, were diagnosed with AMI, coronary artery bypass graft (CABG), or percutaneous coronary interventions (PCI) *or* who were diagnosed with IVD during the Report Period and the year prior to the Report Period. Broken down by gender.

Numerators

Patients with LDL completed during the Report Period, regardless of result.

- Patients with LDL ≤ 100 , completed during the Report Period.
- Patients with LDL 101-130, completed during the Report Period.
- Patients with LDL > 130 , completed during the Report Period.

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Searches for most recent LDL test with a result during the Report Period. If more than one LDL test is found on the same day and/or the same visit and one test has a result and the other does not, the test with the result will be used. If an LDL test with a result is not found, CRS searches for the most recent LDL test without a result. For numerator LDL ≤ 100 , CPT 3048F will count as meeting the measure.

CRS uses the following codes to define the denominator and numerators.

Diagnosis or Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
AMI		V POV: 410.*0, 410.*1		
PCI	33140, 92980, 92982, 92995, G0290	V Procedure: 00.66, 36.01 (old code), 36.02 (old code), 36.05, (old code), 36.06-36.07, or 36.09		
CABG	33510-33514, 33516-33519, 33521-33523, 33533-33536, S2205-S2209	V Procedure: 36.1*, 36.2		
IVD		V POV: 411.*, 413.*, 414.0*, 414.2, 414.8, 414.9, 429.2, 433.*-434.*, 440.1, 440.2*, 440.4, 444.*, or 445.*		
LDL	80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F		Yes	DM AUDIT LDL CHOLESTEROL TAX

Key Logic Changes from CRS Version 11.0

1. Replaced the term “PTCA” with “PCI”.
2. Added HCPCS code G0290 to PCI definition.
3. Added code 43396-1 to LOINC taxonomy for LDL.

Patient List Description

List of patients with AMI, CABG, PCI, or IVD w/LDL value, if any.

Measure Source

HEDIS

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011	Page 250
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		

Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Cholesterol Management for Patients with Cardiovascular Conditions								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical pts 18-75								
with DX of AMI, CABG,								
PCI, or IVD								
	42		28			18		
# w/LDL done	38	90.5	21	75.0	+15.5	9	50.0	+40.5
A. # w/LDL <=100								
w/% of Total								
Screened	14	36.8	12	57.1	-20.3	3	33.3	+3.5
B. # w/LDL 101-130								
w/% of Total								
Screened	5	13.2	3	14.3	-1.1	2	22.2	-9.1
C. # w/LDL >130								
w/% of Total								
Screened	5	13.2	4	19.0	-5.9	4	44.4	-31.3
Male Active Clinical pts								
18-75 with DX of AMI, CABG								
PCI, or IVD								
	22		16			10		
# w/LDL done	20	90.9	11	68.8	+22.2	4	40.0	+50.9
A. # w/LDL <=100								
w/% of Total								
Screened	5	25.0	4	36.4	-11.4	1	25.0	+0.0
B. # w/LDL 101-130								
w/% of Total								
Screened	2	10.0	3	27.3	-17.3	1	25.0	-15.0
C. # w/LDL >130								
w/% of Total								
Screened	4	20.0	2	18.2	+1.8	2	50.0	-30.0
Female Active Clinical pts								
18-75 with DX of AMI, CABG								
PCI, or IVD								
	20		12			8		
# w/LDL done	18	90.0	10	83.3	+6.7	5	62.5	+27.5
A. # w/LDL <=100								
w/% of Total								
Screened	9	50.0	8	80.0	-30.0	2	40.0	+10.0
B. # w/LDL 101-130								
w/% of Total								
Screened	3	16.7	0	0.0	+16.7	1	20.0	-3.3
C. # w/LDL >130								
w/% of Total								
Screened	1	5.6	2	20.0	-14.4	2	40.0	-34.4

Figure 2-92: Sample Report, Cholesterol Management for Patients with Cardiovascular Conditions

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Cholesterol Management for Patients with Cardiovascular Conditions: List of patients with AMI, CABG, PCI, or IVD w/LDL value, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,SHERRY 200.02	000001	COMMUNITY #1	F	68	UP;AC;IVD	DXS02/02/11 LDL:
PATIENT2,CODY JACK	000002	COMMUNITY #1	M	41	UP;AC;IVD	DXS01/18/11 LDL: ***
PATIENT3,TIMOTHY ALLEN	000003	COMMUNITY #1	M	43	UP;AC;IVD	DXSLDL 05/16/11 128
PATIENT4,TRACE	000004	COMMUNITY #1	M	47	UP;AC;IVD	DXS
PATIENT5,KENNETH	000005	COMMUNITY #1	M	60	UP;AC;CABG	CPT11/24/03 LDL: 6
PATIENT6,ROSS WAYNE 83700	000006	COMMUNITY #1	M	60	UP;AC;AMI	DX03/03/DL: CPT
PATIENT7,WILLIAM 3048F	000007	COMMUNITY #1	M	62	UP;AC;PCI	CPT07/30/11 LDL: CPT
PATIENT8,JASON LEE	000008	COMMUNITY #1	M	63	UP;AC;AMI	DX
PATIENT30,ALLISON	000030	COMMUNITY #2	F	52	UP;AC;IVD	DXS11/12/11 LDL: 97
PATIENT31,ALLEN JAMES	000031	COMMUNITY #2	M	44	UP;AC;PCI	PROC11/18/11 LDL: 64

Figure 2-93: Sample Patient List: Cholesterol Management for Patients with Cardiovascular Conditions

2.8.15 Heart Failure and Evaluation of LVS Function

Denominator

Active Clinical patients ages 18 or older discharged with heart failure during the Report Period.

Numerators

Patients whose LVS function was evaluated before arrival, during hospitalization, or is planned for after discharge.

Logic Description

Age of the patient is calculated as of the hospital admission date.

Denominator exclusions are defined as any of the following:

1. Patients receiving comfort measures only (i.e., patients who received palliative care and usual interventions were not received because a medical decision was made to limit care).
2. Patients with a Discharge Type of Transferred or Irregular or containing "Death."
3. Patients who had a left ventricular assistive device (LVAD) or heart transplant procedure during hospitalization.

CRS uses the following codes to define the denominator and numerators.

	CPT Codes	ICD and Other Codes
Denominator Exclusions		
Comfort Measures		V POV: V66.7 (Encounter for palliative care) documented during hospital stay
LVAD/Heart Transplant		V Procedure: 33.6, 37.41, 37.51–37.54, 37.61–37.66, 37.68 documented during hospital stay
Denominator Definition		
Heart Failure		V POV (Primary Diagnosis only): 398.91, 402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 404.93, 428.0, 428.1, 428.20, 428.21, 428.22, 428.23, 428.30, 428.31, 428.32, 428.33, 428.40, 428.41, 428.42, 428.43, 428.9, 429.1, or 997.1 <i>and</i> with Service Category H (hospitalization). NOTE: If a patient has multiple admissions matching these criteria during the Report Period, the earliest admission will be used.
Numerator Definition (Evaluation of LVS Function): Any of the codes listed below		
Ejection Fraction (ordered or documented anytime one year prior to discharge date)	78414, 78468, 78472, 78473, 78480, 78481, 78483, 78494, 93303, 93304, 93307, 93308, 93312, 93314–93318, 93350, 93543, 93555	V Measurement: "CEF" V Procedure: 88.53, 88.54
RCIS Order for Cardiovascular Disorders Referral (ordered during the hospital stay but no later than the hospital discharge date)		ICD Diagnostic Category: "Cardiovascular Disorders" AND one of the following: CPT Categories: "Evaluation and/or Management," "Non-surgical Procedures" or "Diagnostic Imaging"
Other Procedures (documented anytime one year prior to discharge date)		Echocardiogram: V Procedure 88.72, 37.28, 00.24; Nuclear Medicine Test: V Procedure 92.2*; Cardiac Catheterization with a Left Ventriculogram: V Procedure 37.22, 37.23, 88.53, 88.54

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of Active Clinical heart failure patients 18+ who received evaluation of LVS function, if any.

Measure Source

CMS HF-2

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 255	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Heart Failure and Evaluation of LVS Function								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR	% PERIOD		BASE %
AC 18+ w/Heart								
Failure Dx	44		2			1		
Patients w/Eval								
of LVS Function	14	31.8	0	0.0	+31.8	0	0.0	+31.8

Figure 2-94: Sample Report, Heart Failure and Evaluation of LVS Function

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Heart Failure and Evaluation of LVS Function: List of Active Clinical heart failure patients 18+ who received evaluation of LVS function, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	

PATIENT1,JOAN	000164	COMMUNITY	#1	F	36	AC	Admission: 06/01/11
LVS: 06/03/11 Procedure: 88.72							
PATIENT2,SARAH	000127	COMMUNITY	#1	F	35	AC	Admission: 05/01/11
LVS: NOT DOCUMENTED							
PATIENT3,JOHN	000151	COMMUNITY	#1	M	36	AC	Admission: 06/01/11
LVS: 06/01/11 CPT: 78468							
PATIENT4,ROGER	000125	COMMUNITY	#1	M	47	AC	Admission: 05/20/11
LVS: 04/30/11 CEF Measurement 40							
PATIENT5,DANIEL	000129	COMMUNITY	#1	M	57	AC	Admission: 03/14/11
LVS: NOT DOCUMENTED							

Figure 2-95: Sample Patient List: Heart Failure and Evaluation of LVS Function

2.9 STD-Related Measure Topics

2.9.1 HIV Screening

GPRA Measure Description

During FY 2011, achieve the tentative target rate of 73.6% for the proportion of pregnant patients who are screened for HIV.

Denominators

All pregnant Active Clinical female User Population patients with no documented miscarriage or abortion and with no recorded HIV diagnosis ever (GPRA Denominator).

User Population patients ages 13–64 with no recorded HIV diagnosis prior to the Report Period. (GPRA Developmental Denominator).

Numerators

Patients who were screened for HIV during the past 20 months.

Note: This numerator does *not* include refusals. (GPRA Numerator).

Patients with documented HIV screening refusal during the past 20 months.

Patients who were screened for HIV during the Report Period.

Note: This numerator does *not* include refusals. (GPRA Developmental Numerator)

Patients with documented HIV screening refusal during the Report Period.

No denominator. This measure is a total count only, not a percentage. Number of HIV screens provided to User Population patients during the report period, where the patient was not diagnosed with HIV anytime prior to the screen.

Note: This numerator does not include refusals. (GPRA Developmental Numerator)

Logic Description

Age of the patient is calculated at the beginning of the Report period.

Pregnancy definition: At least two visits during the past 20 months from the end of the Report Period. Pharmacy-only visits (Clinic Code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the past 20 months, CRS will use the first two visits in the 20-month period. The patient must not have a documented miscarriage or abortion occurring after the second pregnancy-related visit. In addition, the patient must have at least one pregnancy-related visit occurring during the reporting period. The time period is extended to include patients who were pregnant during the Report Period but whose initial diagnosis (and HIV test) were documented prior to Report Period.

HIV Screening definition: For the number of HIV screens provided to User Population patients numerator (count only), a maximum of one HIV screen per patient per day will be counted.

Note 1: The time frame for both screening and refusals for the pregnant patients denominator is anytime during the past 20 months and for User Population patients 13–64 is anytime during the Report Period.

Note 2: Refusals are allowed during the past 20 months for pregnant patients (vs. only during the Report Period) in the event the patient is at the end of her pregnancy at the beginning of the Report Period and refused the HIV test earlier in her pregnancy during the previous year

CRS uses the following codes and taxonomies to define the denominator and numerators.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Pregnancy (at least 2 visits in past 20 months with 1 during the Report Period)		V POV: V22.0–V23.9, V72.42, 640.* –649.*, 651.* –676.*		
Miscarriage (after second pregnancy POV in past 20 months)	59812, 59820, 59821, 59830	V POV: 630, 631, 632, 633*, 634*		
Abortion (after second pregnancy POV in past 20 months)	59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260–S2267	V POV: 635*, 636* 637* V Procedure: 69.01, 69.51, 74.91, 96.49.		

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
HIV Diagnosis (documented anytime prior to the end of the Report Period)		V POV or Problem List: 042, 042.0–044.9 (old codes), 079.53, V08, or 795.71		
HIV Screening	86689, 86701–86703, 87390, 87391, 87534–87539		Yes	BGP HIV TEST TAX
Refusal of HIV lab test in past 20 months				BGP HIV TEST TAX

Key Logic Changes from CRS Version 11.0

1. Updated patient list.

Patient List Description

List of pregnant patients or User Population patients with documented HIV test or refusal, if any.

Measure Source

HP 2020 HIV-14.3

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	78.0%
IHS FY 2009 Performance	76.0%
IHS FY 2008 Performance	75.0%
IHS FY 2007 Performance	74.0%
IHS FY 2006 Performance	65.0%
IHS FY 2005 Performance	54.0%
<i>HP 2020 Goal</i>	<i>74.1%</i>

DU	May 25, 2011	Page 256
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

HIV Screening		
REPORT	%	PREV YR % CHG from BASE % CHG from

	PERIOD		PERIOD		PREV YR % PERIOD		BASE %	
Pregnant AC Pts w/ no HIV ever (GPRA)	39		38		32			
# w/HIV screening -No Refusals (GPRA)	15	38.5	6	15.8	+22.7	0	0.0	+38.5
# w/HIV screening refusal	1	2.6	0	0.0	+2.6	0	0.0	+2.6
User Pop Pts 13-64 w/ no HIV (GPRA Dev.)	2,024		1,663		1,519			
# w/HIV screening -No Refusals (GPRA Dev.)	46	2.3	21	1.3	+1.0	0	0.0	+2.3
# w/HIV screening refusal	4	0.2	0	0.0	+0.2	0	0.0	+0.2
# HIV screens for User Pop Pts w/ no prior HIV-No Refusals (GPRA Dev.)	51		21		+30		0	
							+51	

Figure 2-96: Sample Report, HIV Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
HIV Screening: List of pregnant patients or User Population patients with documented HIV test or refusal, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,HELEN MARY	000001	COMMUNITY #1	F	12	UP	
PATIENT2,CECELIA	000002	COMMUNITY #1	F	19	UP, UP13-64	03/31/11 LAB
result: NEGA ; 03/31/11 LAB result: NEGATIVE ;Screen Count: 2						
PATIENT15,BRENDA G	000015	COMMUNITY #2	F	30	UP,UP13-64,AC PREG (NO	
DX)03/03/11 lab refusal						
PATIENT16,ALYSHA	000016	COMMUNITY #2	F	33	UP,UP13-64	

Figure 2-97: Sample Patient List, HIV Screening

2.9.2 HIV Quality of Care

Denominator

All User Population patients ages 13 and older with at least two direct care visits (i.e., not Contract/CHS) with HIV diagnosis during the Report Period, including one HIV diagnosis in last six months.

Numerators

Patients who received CD4 test only (without HIV viral load) during the Report Period

Patients who received HIV viral load only (without CD4) during the Report Period

Patients who received both CD4 and HIV viral load during the Report Period

Total patients receiving any test

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

CRS uses the following codes and taxonomies to define the denominator and numerators.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
HIV		042, 042.0-044.9 (old codes), 079.53, V08, 795.71		
CD4	86359, 86360 86361		Yes	BGP CD4 TAX
HIV Viral Load	87536, 87539		Yes	BGP HIV VIRAL LOAD TAX

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients 13 and older diagnosed with HIV, with CD4 test, if any.

Measure Source

HP 2010 developmental measure 13–13a Viral Load Testing

Measure Past Performance and Long-Term Targets

Performance	Percent
<i>IHS 2020 goal for viral load testing</i>	<i>Nearly 100%</i>
<i>IHS 2020 baseline for CD4 testing</i>	<i>Nearly 100%</i>

DU	May 25, 2011	Page 258
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

HIV Quality of Care		
REPORT	%	PREV YR % CHG from BASE % CHG from

	PERIOD		PERIOD		PREV YR % PERIOD		BASE %	
User Pop Pts >13 w/ HIV Dx	6		1		2			
# w/CD4 only	1	16.7	0	0.0	+16.7	0	0.0	+16.7
# w/viral load only	1	16.7	0	0.0	+16.7	0	0.0	+16.7
# w/both	1	16.7	1	100.0	-83.3	2	100.0	-83.3
TOTAL # w/ any tests	3	50.0	1	100.0	-50.0	2	100.0	-50.0

Figure 2-98: Sample Report HIV Quality of Care

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
HIV Quality of Care: List of patients 13 and older diagnosed with HIV, with CD4 test, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,MARY 86359;	000001	COMMUNITY #1	F	19	UP	CD4: 02/01/11
PATIENT2,TANYA	000002	COMMUNITY #1	F	37	UP	
PATIENT15,JOHN 05/01/11 87539	000015	COMMUNITY #2	M	18	UP	Viral Load:
PATIENT16,HAROLD 86360; Viral Load: 03/01/11 87536	000016	COMMUNITY #2	M	20	UP	CD4: 03/01/11

Figure 2-99: Sample Patient List, HIV Quality of Care

2.9.3 Chlamydia Screening

Denominators

Female Active Clinical patients ages 16 through 25.

- Female Active Clinical 16–20.
- Female Active Clinical 21–25.

Female User Population patients ages 16 through 25.

- Female User Population 16–20.
- Female User Population 21–25.

Numerator

Patients tested for Chlamydia during the Report Period.

Logic Description

Age is calculated at beginning of the Report Period. The following codes are used to determine a test for Chlamydia.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Chlamydia Test	86631, 86632, 87110, 87270, 87320, 87490-92, 87810, 3511F	V POV: V73.88, V73.98	Yes	BGP CHLAMYDIA TESTS TAX

Key Logic Changes from CRS Version 11.0

1. Added CPT code 3511F to Chlamydia Test definition.
2. Added codes 57680-1, 57679-3, 57288-3, 57287-5, 56911-1, 56910-3, 56909-5, and 53926-2 to LOINC taxonomy for Chlamydia Test.

Patient List Description

List of patients with documented Chlamydia screening, if any.

Measure Source

HP 2020 STD-4, annual screening for genital Chlamydia–females enrolled in commercial MCOs (aged 25 years and under); STD-3, annual screening for genital Chlamydia–females enrolled in Medicaid MCOs (aged 25 years and under).

Measure Past Performance and Long-Term Targets

Performance	Percent
HP 2020 goal for Females 16-20 with Medicaid (STD-3.1)	57.9%
HP 2020 goal for Females 21-24 with Medicaid (STD-3.2)	65.3%
HP 2020 goal for Females 16-20 with Commercial Health Insurance (STD-4.1)	44.1%
HP 2020 goal for Females 21-24 with Commercial Health Insurance (STD-4.2)	47.9%

DU	May 25, 2011	Page 260
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Chlamydia Testing		

	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
Female Active Clinical 16-25	166		143			128		
# w/Chlamydia Screen	53	31.9	49	34.3	-2.3	43	33.6	-1.7
A. Female Active Clinical 16-20	70		55			57		
# w/Chlamydia Screen	24	34.3	16	29.1	+5.2	23	40.4	-6.1
B. Female Active Clinical 21-25	96		88			71		
# w/Chlamydia Screen	29	30.2	33	37.5	-7.3	20	28.2	+2.0
Female User Population 16-25	287		255			239		
# w/Chlamydia Screen	69	24.0	58	22.7	+1.3	51	21.3	+2.7
A. Female User Population 16-20	139		118			119		
# w/Chlamydia Screen	32	23.0	19	16.1	+6.9	25	21.0	+2.0
B. Female User Population 21-25	148		137			120		
# w/Chlamydia Screen	37	25.0	39	28.5	-3.5	26	21.7	+3.3

Figure 2-100: Sample Report Chlamydia Testing

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Chlamydia Testing: List of patients with documented Chlamydia screening, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,MELISSA ANNE	000001	COMMUNITY #1	F	16	UP;AC	NUMERATOR:
PATIENT2,LISA MARIE	000002	COMMUNITY #1	F	16	UP;AC	NUMERATOR:
PATIENT3,CRYSTAL LEE	000003	COMMUNITY #1	F	17	UP;	NUMERATOR:
PATIENT4,DANIELLE	000004	COMMUNITY #1	F	18	UP;AC	NUMERATOR: lab test
03/17/11						
PATIENT5,KELLYE	000005	COMMUNITY #1	F	19	UP;	NUMERATOR:
PATIENT6,RUBY	000006	COMMUNITY #1	F	19	UP;AC	NUMERATOR: CPT
87490						
06/01/11						
PATIENT7,SANDRA KAY	000007	COMMUNITY #1	F	21	UP;AC	NUMERATOR: lab test
01/27/11						

Figure 2-101: Sample Patient List, Chlamydia Testing

2.9.4 Sexually Transmitted Infection Screening

Denominators

Screenings needed for incidents of key sexually transmitted infections (STIs) for Active Clinical patients that occurred during the period 60 days prior to the beginning of the Report Period through the first 300 days of the Report Period. Key STIs defined as Chlamydia, Gonorrhea, HIV/AIDS, and Syphilis. Broken down by gender.

Chlamydia screenings needed for key STI incidents for Active Clinical patients that occurred during the defined period. Broken down by gender.

Gonorrhea screenings needed for key STI incidents for Active Clinical patients that occurred during the defined period. Broken down by gender.

HIV/AIDS screenings needed for key STI incidents for Active Clinical patients that occurred during the defined period. Broken down by gender.

Syphilis screenings needed for key STI incidents for Active Clinical patients that occurred during the defined period. Broken down by gender.

Screenings needed for incidents of key sexually transmitted infections (STIs) for User Population patients that occurred during the period 60 days prior to the beginning of the Report Period through the first 300 days of the Report Period. Key STIs defined as Chlamydia, gonorrhea, HIV/AIDS, and syphilis. Broken down by gender.

Chlamydia screenings needed for key STI incidents for User Population patients that occurred during the defined period. Broken down by gender.

Gonorrhea screenings needed for key STI incidents for User Population patients that occurred during the defined period. Broken down by gender.

HIV/AIDS screenings needed for key STI incidents for User Population patients that occurred during the defined period. Broken down by gender.

Syphilis screenings needed for key STI incidents for User Population patients that occurred during the defined period. Broken down by gender.

Numerators

No denominator; count only. The total count of *Active Clinical* patients who were diagnosed with one or more key STIs during the period 60 days prior to the Report Period through the first 300 days of the Report Period. Broken down by gender.

No denominator; count only. The total count of separate *key STI incidents for Active Clinical patients* during the defined period. Broken down by gender.

Total number of needed screenings performed or refused from one month prior to the date of relevant STI incident through two months after.

- Number of documented screening refusals.

Number of needed Chlamydia screenings performed or refused from one month prior to the date of first STI diagnosis of each incident through two months after.

- Number of documented screening refusals.

Number of needed Gonorrhea screenings performed or refused from one month prior to the date of first STI diagnosis of each incident through two months after.

- Number of documented screening refusals.

Number of needed HIV/AIDS screenings performed or refused from one month prior to the date of first STI diagnosis of each incident through two months after.

- Number of documented screening refusals.

Number of needed Syphilis screenings performed or refused from one month prior to the date of first STI diagnosis of each incident through two months after.

- Number of documented screening refusals.

No denominator; count only. Total count of *User Population patients* who were diagnosed with one or more key STIs during the period 60 days prior to the Report Period through the first 300 days of the Report Period. Broken down by gender.

No denominator; count only. Total count only of *separate key STI incidents for User Population patients* during the defined period. Broken down by gender.

Logic Description

Key STIs are Chlamydia, Gonorrhea, HIV/AIDS, and Syphilis. Key STI diagnoses are defined with the following codes.

	ICD and Other Codes
Chlamydia	V POV: 078.8*, 079.88, 079.98, 099.41, 099.50-099.59
Gonorrhea	V POV: 098.0-098.89
HIV/AIDS	V POV: 042, 042.0-044.9, 079.53, 795.71, V08
Syphilis	V POV: 090.0-093.9, 094.1-097.9

Logic for Identifying Patients Diagnosed with Key STI:

Any patient with one or more diagnoses of any of the key STIs defined above during the period 60 days prior to the beginning of the Report Period through the first 300 days of the Report Period.

Logic for Identifying Separate Incidents of Key STIs:

One patient may have one or multiple occurrences of one or multiple STIs during the year. Incidents of an STI are identified beginning with the date of the first key STI diagnosis (see definition above) occurring between 60 days prior to the beginning of the Report Period through the first 300 days of the Report Period. A second incident of the same STI is counted if another diagnosis with the same STI occurs two months or more after the initial diagnosis. A different STI diagnosis that occurs during the same 60-day time period as the first STI counts as a separate incident.

Example of Patient with Multiple Incidents of Single STI:

Visit	Total Incidents
08/01/10: Patient screened for Chlamydia	0
08/08/10: Patient diagnosed with Chlamydia	1
10/15/10: Patient diagnosed with Chlamydia	2
10/25/10: Follow-up for Chlamydia	2
11/15/0: Patient diagnosed with Chlamydia	2
03/01/11: Patient diagnosed with Chlamydia	3

Denominator Logic for Needed Screenings:

One patient may need multiple screening tests based on one or more STI incidents occurring during the time period.

To be included in the needed screening tests denominator, the count will be derived from the number of separate STI incidents and the type(s) of screenings recommended for each incident. The recommended screenings for each key STI are listed below.

STI	Screenings Needed
Chlamydia	Gonorrhea, HIV/AIDS, Syphilis
Gonorrhea	Chlamydia, HIV/AIDS, Syphilis
HIV/AIDS	Chlamydia, Gonorrhea, Syphilis
Syphilis	Chlamydia, Gonorrhea, HIV/AIDS

“Needed” screenings are recommended screenings that are further evaluated for contraindications. The following are reasons that a recommended screening is identified as not needed (i.e., contraindicated).

1. The patient has a documented STI diagnosis corresponding to the screening type in the same time period. For example, a patient with both a Chlamydia and a gonorrhea diagnosis on the same visit does not need the recommended Chlamydia screening based on the gonorrhea diagnosis.

2. Only one screening for each type of STI is needed during the relevant time period, regardless of the number of different STI incidents identified. For example, if a patient is diagnosed with Chlamydia and Gonorrhea on the same visit, only one screening each is needed for HIV/AIDS and Syphilis.
3. A patient with HIV/AIDS diagnosis prior to any STI diagnosis that triggers a recommended HIV/AIDS screening does not need the screening ever.

Numerator Logic:

To be counted in the numerator, each needed screening in the denominator must have a corresponding laboratory test or test refusal documented in the period from one month prior to the relevant STI diagnosis date through two months after the STI incident.

Key STI screenings are defined with the following codes.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Chlamydia	86631–86632, 87110, 87270, 87320, 87490–87492, 87810	POV: V73.88, V73.98	Yes	BGP CHLAMYDIA TESTS TAX
Gonorrhea	87590–87592, 87850		Yes	BKM GONORRHEA TEST TAX
HIV/AIDS	86689, 86701–86703, 87390–87391, 87534–87539		Yes	BGP HIV TEST TAX
Syphilis	86592–86593, 86781, 87285		Yes	BKM FTA-ABS TESTS TAX or BKM RPR TESTS TAX
Refusal of any screening	Any refusal type (REF, NMI, etc.) for any of the four screening tests as defined above during the specified time period.			

Logic Examples

Example of Patient with Single Diagnosis of Single STI:

08/01/10:	Patient screened for Chlamydia
08/08/10:	Patient diagnosed with Chlamydia–3 screens needed: Gonorrhea, HIV/AIDS, Syphilis
08/13/10:	Patient screened for Gonorrhea, HIV/AIDS, Syphilis
Result:	Denominator: 3 screens needed, Numerator: 3 screens performed.

Example of Patient with Multiple Diagnoses of Single STI:

08/01/10:	Patient screened for Chlamydia
08/08/10:	Patient diagnosed with Chlamydia (Incident #1)–3 screens needed: Gonorrhea, HIV/AIDS, Syphilis
12/01/10:	Patient screened for Chlamydia
12/08/10:	Patient diagnosed with Chlamydia (Incident #2) –3 screens needed: Gonorrhea, HIV/AIDS, Syphilis
Result:	Denominator: 6 screens needed (2 each of 3 types), Numerator: 3 screens performed (1 each of 3 types)

Example of Patient with Single Diagnosis of Multiple STIs:

10/15/10:	Patient screened for Chlamydia, Gonorrhea, HIV/AIDS, Syphilis
10/18/10:	Patient diagnosed with Chlamydia–3 screens needed: Gonorrhea, HIV/AIDS, Syphilis
10/20/10:	Patient diagnosed with Syphilis–removes needed screen for Syphilis (see above)
Result:	Denominator: 2 screens needed, Numerator: 2 screens performed prior to triggering diagnoses but within timeframe

Example of Patient with Multiple Diagnoses of Multiple STIs:

06/15/05:	Patient diagnosed with HIV/AIDS
08/01/10:	Patient screened for Chlamydia and Gonorrhea
08/08/10:	Patient diagnosed with Chlamydia and Gonorrhea (Incident #1)–1 screen needed: Syphilis (HIV/AIDS not needed since prior diagnosis)
08/08/10:	Patient screened for HIV/AIDS and Syphilis–since only the Syphilis screen is needed, the HIV/AIDS screen is not counted at all
12/01/10:	Patient screened for Chlamydia
12/08/10:	Patient diagnosed with Chlamydia (Incident #2) –2 screens needed: Gonorrhea and Syphilis
12/10/10:	Patient screened for Syphilis
Result:	Denominator: 3 screens needed (2 Syphilis and 1 Gonorrhea), Numerator: 2 screens performed (2 Syphilis)

Key Logic Changes from CRS Version 11.0

1. Added codes 57680-1, 57679-3, 57288-3, 57287-5, 56911-1, 56910-3, 56909-5, and 53926-2 to LOINC taxonomy for Chlamydia Screening.

Patient List Description

List of patients diagnosed with one or more STIs during the defined time period with related screenings.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU

May 25, 2011

Page 266

*** IHS 2011 Selected Measures with Community Specified Report ***

DEMO INDIAN HOSPITAL

Report Period: Jan 01, 2011 to Dec 31, 2011

Previous Year Period: Jan 01, 2010 to Dec 31, 2010

Baseline Period: Jan 01, 2000 to Dec 31, 2000

Sexually Transmitted Infection (STI) Screening (con't)

	REPORT PERIOD		PREV YR PERIOD		CHG from PREV YR		BASE PERIOD		CHG from BASE
Active Clinical Pts w/ Key STI Dx	40		13		+27		10		+30
Male Active Clinical Pts w/ Key STI Dx	10		7		+3		8		+2
Female Active Clinical Pts w/ Key STI Dx	30		6		+24		2		+28
# Key STI Incidents for Active Clinical Pts	47		16		+31		15		+32
# Male AC Key STI Incidents	12		9		+3		13		-1
# Female AC Key STI Incidents	35		7		+28		2		+33
# Key STI Screens Needed for AC Pts	131		48				45		
# Needed Screens Performed/Refused	31	23.7	6	12.5	+11.2		8	17.8	+5.9
A. # Documented Refusals	2	1.5	0	0.0	+1.5		0	0.0	+1.5
# Key STI Screens Needed for Male AC Pts	35		27				39		
# Needed Screens Performed/Refused	5	14.3	4	14.8	-0.5		7	17.9	-3.7
A. # Documented Refusals	0	0.0	0	0.0	+0.0		0	0.0	+0.0
# Key STI Screens Needed for Female AC Pts	96		21				6		

# Needed Screens								
Performed/Refused	26	27.1	2	9.5	+17.6	1	16.7	+10.4
A. # Documented								
Refusals	2	2.1	0	0.0	+2.1	0	0.0	+2.1

Figure 2-102: Sample Report Sexually Transmitted Infection (STI) Screening

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease								
Sexually Transmitted Infection (STI) Screening: List of patients diagnosed with one or more STIs during the defined time period with related screenings.								
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR		

PATIENT1,DIANE	000001	COMMUNITY #1	F	15	UP 1) 08/28/11	POV: CHL 079.98;		
1) GC-Y 08/20/11 Lab [GonoDNA]; HIV-N ; SYP-N ; ;								
PATIENT2,LEIGHANN	000002	COMMUNITY #1	F	18	UP;AC 1 1) 04/30/11	POV: HIV 042.; 1) CHL-Y 04/04/03 Lab [ChlaDNA]; GC-Y 05/15/11 Lab [HGB]; SYP-N ; ;		
PATIENT3,WHITNEY	000003	COMMUNITY #1	F	25	UP;AC 1) 11/02/10	POV: SYP 094.1; 1) CHL-N ; GC-N ; HIV-Y 10/05/10 CPT [86703]; ;		
PATIENT4,NANCY	000004	COMMUNITY #1	F	29	UP;AC 1) 10/26/11	POV: GC 098.89; 1) CHL-N ; HIV-N ; SYP-N ; ;		
PATIENT5,JOHN	000005	COMMUNITY #1	M	40	UP;AC 1) 03/13/11	POV: HIV 042.; 1) CHL-N ; GC-Y 03/13/11 Lab [HGB]; SYP-N ; ;		
PATIENT6,NORMAN	000006	COMMUNITY #1	M	42	UP;AC 1) 04/14/11	POV: HIV 079.53; 2) 06/14/11 POV: HIV V08.; 1) CHL-N ; GC-N ; SYP-N ; ; 2) CHL-N ; GC-N ; SYP-N ; ;		

Figure 2-103: Sample Patient List, Sexually Transmitted Infection (STI) Screening

2.10 Other Clinical Measures Topics

2.10.1 Osteoporosis Management

Denominators

Female Active Clinical patients ages 67 and older who had a new fracture occurring 6 months (180 days) prior to the Report Period through the first 6 months of the Report Period with no osteoporosis screening or treatment in year prior to the fracture.

Female User Population patients ages 67 and older who had a new fracture occurring 6 months (180 days) prior to the Report Period through the first 6 months of the Report Period with no osteoporosis screening or treatment in year prior to the fracture.

Numerator

Patients treated or tested for osteoporosis after the fracture.

Logic Description

Age is calculated at the beginning of the Report Period. Fractures do not include fractures of finger, toe, face, or skull. CRS will search for the first (i.e., earliest) fracture during the period 6 months (180) days prior to the beginning of the Report Period and the first 6 months of the Report Period. If multiple fractures are present, only the first fracture will be used.

Index Episode Start Date definition: The Index Episode Start Date is the date the fracture was diagnosed. If the fracture was diagnosed at an outpatient visit (Service Category A, S, or O), the Index Episode Start Date is equal to the Visit Date. If diagnosed at an inpatient visit (Service Category H), the Index Episode Start Date is equal to the Discharge Date.

Denominator Exclusions

1. Patients receiving osteoporosis screening or treatment in the year (365 days) prior to the Index Episode Start Date. Osteoporosis screening or treatment is defined as a Bone Mineral Density (BMD) test (see below for codes) or receiving any osteoporosis therapy medication (see below for codes).
2. Patients with a fracture diagnosed at an outpatient visit, which also had a fracture within 60 days prior to the Index Episode Start Date.
3. Patients with a fracture diagnosed at an inpatient visit, which also had a fracture within 60 days prior to the ADMISSION DATE.

Osteoporosis Treatment and Testing definition: (1) For fractures diagnosed at an outpatient visit: a nondiscontinued prescription within 6 months (180 days) of the Index Episode Start Date (i.e., visit date) or B) a BMD test within 6 months of the Index Episode Start Date. (2) For fractures diagnosed at an inpatient visit, a BMD test performed during the inpatient stay.

CRS uses the following codes to define fracture and BMD test.

	CPT Codes	ICD and Other Codes
Fracture Codes	21800–21825, 22305–22314, 22316–22324, 22520, 22521, 22523, 22524, 23500–23515, 23570–23630, 23665–23680, 24500–24585, 24620, 24635, 24650–24685, 25500–25609, 25611 (old code), 25620 (old code), 25622–25652, 25680, 25685, 27193–27248, 27254, 27500–27514, 27520–27540, 27750–27828, S2360, S2362	V POV: 733.1*, 805*–806*, 807.0*–807.4, 808*–815*, 818*–825*, 827*, 828* V Procedure: 79.01–79.03, 79.05–79.07, 79.11–79.13, 79.15–79.17, 79.21–79.23, 79.25–79.27, 79.31–79.33, 79.35–79.37, 79.61–79.63, 79.65–79.67, 81.65–81.66.

BMD Test Codes	77078, 76070 (old code), 77079, 76071 (old code), 77080, 76075 (old code), 77081, 76076 (old code), 77083, 76078 (old code), 76977, 78350, 78351, G0130	V POV: V82.81 V Procedure: 88.98
-----------------------	---	---

Treatment medication codes are defined with medication taxonomy BGP HEDIS OSTEOPOROSIS MEDS. (Medications are: Alendronate, Alendronate-Cholecalciferol (Fosomax Plus D), Ibandronate (Boniva), Risedronate, Calcitonin, Raloxifene, Estrogen, Injectable Estrogens, and Teriparatide.) Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of female patients with new fracture who have had osteoporosis treatment or testing, if any.

Measure Source

HEDIS

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 277	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Osteoporosis Management								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Female Active Clinical Pts								
67 and older								
w/fracture	8		0			0		
# w/osteoporosis treatment								
or testing	3	37.5	0	0.0	+37.5	0	0.0	+37.5
Female User Pop Pts								
67 and older								
w/fracture	9		0			0		

# w/osteoporosis treatment or testing	4	44.4	0	0.0	+44.4	0	0.0	+44.4
---------------------------------------	---	------	---	-----	-------	---	-----	-------

Figure 2-104: Sample Report Osteoporosis Management

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Osteoporosis Management: List of female patients with new fracture who have had osteoporosis treatment or testing, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,ALWENA	000001	COMMUNITY #1	F	68	UP,AC FX: 01/01/11 CPT 22524	
PATIENT2,SYBIL	000002	COMMUNITY #1	F	69	UP,AC FX: 02/10/11 CPT 22524TX:	
02/10/11 CPT G0130						
PATIENT3,ELIZABETH	000003	COMMUNITY #1	F	78	UP,AC FX: 02/15/11 CPT S2362	
PATIENT4,KATIE	000004	COMMUNITY #1	F	80	UP,AC FX: 02/05/11 PROC 81.66	
PATIENT5,LINDSAY	000005	COMMUNITY #1	F	81	UP FX: 02/01/11 DX 733.13TX:	
02/15/11 CPT 77081						
PATIENT6,ELIZABETH	000006	COMMUNITY #1	F	86	UP,AC FX: 01/15/11 DX 733.13TX:	
01/31/11 CPT 77081						

Figure 2-105: Sample Patient List, Osteoporosis Management

2.10.2 Osteoporosis Screening in Women

Denominators

Female Active Clinical patients ages 65 and older without a documented history of osteoporosis.

Female User Population patients ages 65 and older without a documented history of osteoporosis.

Numerators

Patients who had osteoporosis screening documented in the past two years.

Note: This numerator does *not* include refusals.

Patients with documented refusal in past year.

Logic Description

Age is calculated at the beginning of the Report Period.

Osteoporosis definition: No osteoporosis diagnosis ever (POV 733.*).

CRS uses the following codes to define osteoporosis screening.

	CPT Codes	ICD and Other Codes
Osteoporosis Screening (any test documented in the past two years)	Central DEXA: 77080, 76075 (old code) Peripheral DEXA: 77081, 76076 (old code) SEXA: G0130 Central CT: 77078, 76070 (old code) Peripheral CT: 77079, 76071 (old code) US Bone Density: 76977	V Procedure: 88.98 (Quantitative CT) V POV: V82.81 Special screening for other conditions, Osteoporosis
Osteoporosis Screening Refusal	Refusal (in past year): CPT or V Radiology: Central DEXA: 77080 or 76075 (old code); Peripheral DEXA: 77081 or 76076 (old code); SEXA: G0130; Central CT: 77078 or 76070 (old code); Peripheral CT: 77079 or 76071 (old code); US Bone Density: 76977	

Key Logic Changes from CRS Version 11.0

1. Removed refusals from main numerators. Refusal numerator is a separate numerator, and no longer a subnumerator.

Patient List Description

List of female patients ages 65 and older with osteoporosis screening, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011	Page 279
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Osteoporosis Screening in Women		
REPORT PERIOD	%	PREV YR PERIOD % CHG from BASE PREV YR % PERIOD % CHG from BASE %
Female Active Clinical		

Pts =>65	54	29	29					
# w/osteoporosis screening in past 2 years								
-No Refusals	6	11.1	0	0.0	+11.1	0	0.0	+11.1
# w/ Osteoporosis Screening Refusal	2	3.7	0	0.0	+3.7	0	0.0	+3.7
Female User Pop								
Pts =>65	111	79	80					
# w/osteoporosis screening in past 2 years								
-No Refusals	6	5.4	0	0.0	+5.4	0	0.0	+5.4
# w/ Osteoporosis Screening Refusals	2	1.8	0	0.0	+1.8	0	0.0	+1.8

Figure 2-106: Sample Report, Osteoporosis Screening in Women

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Osteoporosis Screening in Women: Osteoporosis Screening in Women: List of female patients ages 65 and older with osteoporosis screening or refusal, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,SHERRY	000001	COMMUNITY #1	F	68	UP;AC	05/15/11	Refusal
88.98							
PATIENT2,APRIL	000002	COMMUNITY #1	F	68	UP;AC	04/01/11	CPT G0130
PATIENT3,JACKIE	000003	COMMUNITY #1	F	69	UP;AC	08/21/11	CPT G0130
PATIENT4,PAULINE	000004	COMMUNITY #1	F	70	UP;AC		
PATIENT5,SHANNON	000005	COMMUNITY #1	F	72	UP;AC		
PATIENT6,TINA MARIE	000006	COMMUNITY #1	F	78	UP;AC	04/15/11	CPT 77081

Figure 2-107: Sample Patient List, Osteoporosis Screening in Women

2.10.3 Rheumatoid Arthritis Medication Monitoring

Denominator

Active Clinical patients ages 16 and older diagnosed with *rheumatoid arthritis (RA)* prior to the Report Period and with at least two RA-related visits any time during the Report Period who were prescribed maintenance therapy medication chronically during the Report Period.

Numerator

Patients who received appropriate monitoring of chronic medication during the Report Period.

Logic Description

Age is calculated at the beginning of the Report Period.

RA defined as diagnosis (POV or Problem List) 714.* prior to the Report Period, and at least two RA POVs during the Report Period.

For all maintenance therapy medications *except* intramuscular gold, each medication must be prescribed within the past 465 days of the end of the Report Period (i.e., the Medication Period) and the sum of the days supply ≥ 348 . This means the patient must have been on the medication at least 75% of the Medication Period. Two examples are shown below to illustrate this logic. *All* medications must not have a comment of RETURNED TO STOCK.

Example of Patient Not on Chronic Medication (not included in Denominator):

Report Period: Jan 1–Dec 31, 2011

Medication Period: 465 days from end of Report Period (Dec 31, 2011): Sep 22, 2010–Dec 31, 2011

Medication Prescribed:

Diclofenac: 1st Rx: Oct 15, 2010, Days Supply=90; 2nd Rx: Jan 01, 2011: Days Supply=90; 3rd Rx: Mar 15, 2011: Days Supply=90.

Total Days Supply=270. 270 is not >348 . Patient is not considered on chronic medication and is not included in the denominator.

Example of Patient on Chronic Medication (included in Denominator):

Report Period: Jan 1–Dec 31, 2011

Medication Period: 465 days from end of Report Period (Dec 31, 2011): Sep 22, 2010–Dec 31, 2011

Medications Prescribed:

Sulfasalazine: 1st Rx: Sep 30, 2010, Days Supply=90; 2nd Rx: Dec 30, 2010, Days Supply=90; 3rd Rx: Mar 15, 2011 Days Supply=180.

Total Days Supply=360. 360 is >348 . Patient is considered on chronic medication and is included in denominator.

The days supply requirement may be met with a single prescription or from a combination of prescriptions for the same medication that were filled during the Medication Period. However, for all medications, there must be at least one prescription filled during the Report period.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

For intramuscular gold, the patient must have 12 or more prescriptions during the Report Period.

Appropriate monitoring of rheumatoid arthritis medications is defined with lab tests and varies by medication, as shown in the table below. If patient is prescribed two or more types of medications, patient must meet criteria for all of the medications.

Maintenance Therapy Medications Definition

- **Medications shown in table below.** *Except* for Gold, Intramuscular, all medications requiring more than one of each type of test during the Report Period, there must be a minimum of ten days between tests. For example, if a Sulfasalazine test was performed on March 1, March 7, and March 21, 2011, the March 7 test will not be counted since it was performed only six days after the March 1 test.

Medication	Required Monitoring Tests
Gold, Intramuscular	CBC and urine Protein on same day as each injection during Report Period
Azathioprine or Sulfasalazine	4 CBCs during the Report Period
Leflunomide or Methotrexate	6 each of CBC, Serum Creatinine, and Liver Function Tests during the Report Period
Cyclosporin	CBC, Liver Function Tests, and Potassium within past 180 days from Report Period end date 12 Serum Creatinine tests during the Report Period
Gold, Oral or Penicillamine	4 each of CBC and Urine Protein during the Report Period
Mycophenolate	CBC within past 180 days from Report Period end date

The medications in the above table are defined with medication taxonomies: BGP RA IM GOLD MEDS, BGP RA AZATHIOPRINE MEDS, BGP RA LEFLUNOMIDE MEDS, BGP RA METHOTREXATE MEDS, BGP RA CYCLOSPORINE MEDS, BGP RA ORAL GOLD MEDS, BGP RA MYCOPHENOLATE MEDS, BGP RA PENICILLAMINE MEDS, BGP RA SULFASALAZINE MEDS.

1. **NSAID Medications:** All of the following medications must have Creatinine, Liver Function Tests, and CBC during the Report Period: Diclofenac, Etodolac, Indomethacin, Ketorolac, Sulindac, Tolmetin, Meclofenamate, Mefenamic Acid, Nabumetone, Meloxicam, Piroxicam, Fenoprofen, Flurbiprofen, Ibuprofen, Ketoprofen, Naproxen, Oxaprozin, Aspirin, Choline Magnesium Trisalicylate, Diflunisil, Magnesium Salicylate, Celcoxib. All of these medications *except* aspirin are defined with medication taxonomy BGP RA OA NSAID MEDS. Aspirin defined with medication taxonomy DM AUDIT ASPIRIN DRUGS.

2. **Glucocorticoid Medications:** Dexamethasone, Methylprednisolone, Prednisone, Hydrocortisone, Betamethasone, Prednisolone, Triamcinolone. These medications defined with medication taxonomy BGP RA GLUCOCORTICIDS MEDS. Glucocorticoids must have a glucose test, which must be performed during the Report Period.

Example of Patient Not Included in Numerator:

Medications Prescribed and Required Monitoring:

- Gold, Oral, last Rx Jun 15, 2011. Requires CBC and Urine Protein within past 90 days of Report Period end date.
- CBC performed on Dec 1, 2011, which is within past 90 days of Report Period end date of Dec 31, 2011. No Urine Protein performed during that period. Patient is not in numerator.

Example of Patient Included in Numerator:

Medications Prescribed and Required Monitoring:

- Diclofenac, last Rx Sep 1, 2011. Requires LFT and CBC during Report Period.
- Mycophenolate, last Rx Mar 10, 2011. Requires CBC within past 180 days from Report Period end date.
- LFT and CBC performed during Report Period. CBC performed Nov 1, 2011, which is within past 180 days of Report Period end date of Dec 31, 2011. Patient is in numerator.

Monitoring Test	CPT Codes	LOINC Codes	Taxonomy
CBC	85025, 85027	Yes	BGP CBC TESTS
Urine Protein		Yes	DM AUDIT URINE PROTEIN TAX
Serum Creatinine	82540, 82565-75	Yes	DM AUDIT CREATININE TAX
Liver Function Tests-ALT	84460	Yes	DM AUDIT ALT TAX
Liver Function Tests-AST	84450	Yes	DM AUDIT AST TAX
Liver Function	80076	Yes	BGP LIVER FUNCTION TESTS
Glucose	82947, 82948, 82950, 82951, 82952, 82962	Yes	DM AUDIT GLUCOSE TESTS TAX
Potassium	84132	Yes	BGP POTASSIUM TESTS

Key Logic Changes from CRS Version 11.0

1. Added codes 16325-3, 48136-6, and 54500-4 to LOINC taxonomy for AST.
2. Added codes 47288-6, 58410-2, and 55429-5 to LOINC taxonomy for CBC.

- Added codes 2885-2, 1751-7, 1975-2, 1968-7, 6768-6, 1920-8, and 1742-6 to LOINC taxonomy for Liver Function.

Patient List Description

List of RA patients age 16 and older prescribed maintenance therapy medication with monitoring laboratory tests, if any. The numerator values for patients who meet the measure are prefixed with YES and patients who did not meet the measure are prefixed with NO. The chronic medications and all laboratory tests the patient *did* have are displayed.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 284	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Rheumatoid Arthritis Medication Monitoring								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts =>16								
w/RA DX and maintenance								
therapy RX	3		0			0		
# w/RA chronic med								
monitoring	2	66.7	0	0.0	+66.7	0	0.0	+66.7

Figure 2-108: Sample Report, Rheumatoid Arthritis Medication Monitoring

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Rheumatoid Arthritis Medication Monitoring: List of RA patients 16 and older prescribed maintenance therapy medication with monitoring lab tests, if any. The numerator values for patients who meet the measure are prefixed with YES: and patients who did not meet the measure are prefixed with NO:. The chronic medications and all lab tests the patient DID have are displayed.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,RUTH	000001	COMMUNITY #1	F	64	;AC	YES: NSAID:
10/21/11 CREAT, 09/22/11 CBC, 05/21/11 LFT10/21/11 CREAT, 09/22/11 CBC, 05/21/11 LFT						

PATIENT2,SHANNON	000002	COMMUNITY #1	F	72	;AC	YES:
Glucocorticoids:	has	Glucose				
PATIENT34,CATHERINE	000034	COMMUNITY #3	F	50	;AC	NO:
Glucocorticoids:	does not have	Glucose				

Figure 2-109: Sample Patient List, Rheumatoid Arthritis Medication Monitoring

2.10.4 Osteoarthritis Medication Monitoring

Denominator

Active Clinical patients ages 40 and older diagnosed with *osteoarthritis (OA)* prior to the Report Period and with at least two OA-related visits any time during the Report Period and prescribed maintenance therapy medication chronically during the Report Period.

Numerator

Patients who received appropriate monitoring of chronic medication during the Report Period.

Logic Description

Age is calculated at the beginning of the Report Period.

OA defined as diagnosis (POV or Problem List) 715.* prior to the Report period, and at least two OA POVs during the Report Period.

For all maintenance therapy medications, each medication must be prescribed within the past 465 days of the end of the Report Period (i.e., the Medication Period) and the sum of the days supply =>348. This means the patient must have been on the medication at least 75% of the Medication Period. Two examples are shown below to illustrate this logic. Medications must not have a comment of RETURNED TO STOCK.

Example of Patient Not on Chronic Medication (not included in Denominator):

- Report Period: Jan 1–Dec 31, 2011
- Medication Period: 465 days from end of Report Period (Dec 31, 2011): Sep 22, 2010–Dec 31, 2011
- Medication Prescribed:
 - Diclofenac: 1st Rx: Oct 15, 2010, Days Supply=90; 2nd Rx: Jan 1, 2011: Days Supply=90; 3rd Rx: Mar 15, 2011: Days Supply=90.
 - Total Days Supply=270. 270 is not >348. Patient is not considered on chronic medication and is not included in the denominator.

Example of Patient on Chronic Medication (included in Denominator):

- Report Period: Jan 1–Dec 31, 2011
- Medication Period: 465 days from end of Report Period (Dec 31, 2011): Sep 22, 2010–Dec 31, 2011
- Medication Prescribed:
 - Etodolac: 1st Rx: Sep 30, 2010, Days Supply=90; 2nd Rx: Dec 30, 2010, Days Supply=90; 3rd Rx: Mar 15, 2011: Days Supply =180.
 - Total Days Supply=360. 360 is >348. Patient is considered on chronic medication and is included in denominator.

The days supply requirement may be met with a single prescription or from a combination of prescriptions for the same medication that were filled during the Medication Period. However, for all medications, there must be at least one prescription filled during the Report Period.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Appropriate monitoring of OA medications is defined with laboratory tests and varies by medication, as shown below.

Maintenance Therapy Medications Defined with the Following NSAID Medications:

Diclofenac, Etodolac, Indomethacin, Ketorolac, Sulindac, Tolmetin, Meclofenamate, Mefenamic Acid, Nabumetone, Meloxicam, Piroxicam, Fenoprofen, Flurbiprofen, Ibuprofen, Ketoprofen, Naproxen, Oxaprozin, Aspirin, Choline Magnesium Trisalicylate, Diflunisil, Magnesium Salicylate, Celcoxib. All of these medications, *except* aspirin, are defined with medication taxonomy BGP RA OA NSAID MEDS. Aspirin defined with medication taxonomy DM AUDIT ASPIRIN DRUGS

All NSAID medications must have Creatinine, Liver Function Tests and CBC during the Report Period.

Example of Patient Not Included in Numerator:

- Medication Prescribed and Required Monitoring:
 - Diclofenac, last Rx Jun 15, 2011. Requires Creatinine, LFT and CBC during Report Period. Only the LFT was performed during Report Period. Patient is not in numerator.

Example of Patient Included in Numerator:

- Medications Prescribed and Required Monitoring:

- Diclofenac, last Rx Sep 1, 2011. Requires Creatinine, LFT and CBC during Report Period. Creatinine, LFT, and CBC performed during Report Period. Patient is in numerator.

CRS uses the following codes to define the monitoring tests.

Monitoring Test	CPT Codes	LOINC Codes	Taxonomy
Serum Creatinine	82540, 82565-82575	Yes	DM AUDIT CREATININE TAX
CBC	85025, 85027	Yes	BGP CBC TESTS
Liver Function Tests-ALT	84460	Yes	DM AUDIT ALT TAX
Liver Function Tests-AST	84450	Yes	DM AUDIT AST TAX
Liver Function	80076	Yes	BGP LIVER FUNCTION TESTS

Key Logic Changes from CRS Version 11.0

1. Added codes 16325-3, 48136-6, and 54500-4 to LOINC taxonomy for AST.
2. Added codes 47288-6, 58410-2, and 55429-5 to LOINC taxonomy for CBC.
3. Added codes 2885-2, 1751-7, 1975-2, 1968-7, 6768-6, 1920-8, and 1742-6 to LOINC taxonomy for Liver Function.

Patient List Description

List of OA patients 40 and older prescribed maintenance therapy medication with monitoring laboratory tests, if any. The numerator values for patients who meet the measure are prefixed with YES and patients who did not meet the measure are prefixed with NO. All laboratory tests the patient did have are displayed.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011					Page 287	
*** IHS 2011 Selected Measures with Community Specified Report ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Previous Year Period: Jan 01, 2010 to Dec 31, 2010							
Baseline Period: Jan 01, 2000 to Dec 31, 2000							

Osteoarthritis Medication Monitoring							
REPORT	%	PREV YR	%	CHG from	BASE	%	
PERIOD		PERIOD		PREV YR	% PERIOD	BASE %	

Active Clinical Pts =>40								
w/OA DX and maintenance								
therapy RX	3		6			4		
# w/OA chronic								
med monitoring	2	66.7	3	50.0	+16.7	2	50.0	+16.7

Figure 2-110: Sample Report, Osteoarthritis Medication Monitoring

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Osteoarthritis Medication Monitoring: List of OA patients 40 and older prescribed maintenance therapy medication with monitoring lab tests, if any. The numerator values for patients who meet the measure are prefixed with "YES:" and patients who did not meet the measure are prefixed with "NO:". All lab tests the patient DID have are displayed.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,RUTH 10/21/11 CREAT, 09/22/11 CBC, 05/21/11 LFT	000001	COMMUNITY #1	F	64	;AC	451 days of nsaid YES:
PATIENT2,JACKIE CREAT: 10/30/11 CBC: 08/06/11 LFT: 10/30/11	000002	COMMUNITY #1	F	69	;AC	472 days of nsaid YES:
PATIENT15,RAYMOND CREAT: 09/12/11	000015	COMMUNITY #2	M	84	;AC	804 days of nsaid NO:
PATIENT33,ROBERT LEE 12/02/11 LFT: 08/06/11	000033	COMMUNITY #3	M	62	;AC	397 days of nsaid NO: CBC:
Total # of Patients on list: 4						

Figure 2-111: Sample Patient List, Osteoarthritis Medication Monitoring

2.10.5 Asthma

Denominators

All *Active Clinical patients*. Broken down by age groups (under 5, 5 to 64, and 65 and older).

Patients who have had two asthma-related visits during the Report Period or with persistent asthma. Broken down by age groups (under 5, 5 to 64, and 65 and older).

Numerators

Patients who have had two asthma-related visits during the Report Period or with persistent asthma.

Patients from Numerator 1 who have been hospitalized at any hospital for asthma during the Report Period.

Logic Description

Age is calculated at beginning of Report Period.

Asthma visits definition: Diagnosis (POV) 493.*.

Persistent asthma definition: Any of the following:

- Active entry in PCC Problem List for 493.* with Severity of 2, 3 or 4 at *any* time before the end of the Report Period, *or*
- Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 2, 3, or 4 documented *any* time before the end of the Report Period.

Hospitalizations definition: Service Category H with primary POV 493.*.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients diagnosed with asthma and any asthma-related hospitalizations.

Measure Source

HP 2020 RD-2

Measure Past Performance and Long-Term Targets

Measure	Target
HP1998 baseline for hospitalizations for asthma:	
Under 5	45.6 per 10,000
5-64	12.5 per 10,000
65 and older	17.7 per 10,000
HP 2020 goal for hospitalizations for asthma:	
<i>Under 5</i>	<i>18.1 per 10,000</i>
<i>5-64</i>	<i>8.6 per 10,000</i>
<i>65 and older</i>	<i>20.3 per 10,000</i>

DU	May 25, 2011							Page 289	
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Asthma (con't)	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Total Active Clinical Patients	1,554		1,209			1,101			
# w/asthma	51	3.3	38	3.1	+0.1	25	2.38	+1.0	
A. Under 5	9	17.6	13	34.2	-16.6	12	48.0	-30.4	

B. 5-64	39	76.5	24	63.2	+13.3	11	44.0	+32.5
C. 65 and older	3	5.9	1	2.6	+3.3	2	8.0	-2.1
# w/asthma	51		38			25		
# w/asthma hospitalization	0	0.0	1	2.6	-2.6	2	8.0	-8.0
A. Under 5	0	0.0	0	0.0	+0.0	1	50.0	-50.0
B. 5-64	0	0.0	1	100.0	-100.0	1	50.0	-50.0
C. 65 and older	0	0.0	0	0.0	+0.0	0	0.0	+0.0

Figure 2-112: Sample Report, Asthma

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Asthma: List of patients diagnosed with asthma and any asthma-related hospitalizations.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,GENEVA 02/02/10	000001	COMMUNITY #1	F	47	AC	Severity 4 on visit	
PATIENT2,JACKIE 06/24/11 08/27/11	000002	COMMUNITY #1	F	69	AC	2 Dxs PCC:	
PATIENT3,PAULINE 02/12/11 02/20/11	000003	COMMUNITY #1	F	70	AC	2 Dxs PCC:	
PATIENT4,WILLIAM R	000004	COMMUNITY #1	M	7	AC	Severity 2 on PL	
PATIENT5,ZACHARY 03/03/11	000005	COMMUNITY #1	M	11	AC	Severity 2 on visit	
PATIENT42,JOSEPHINE	000042	COMMUNITY #2	F	4	AC	Severity 4 on PL	

Figure 2-113: Sample Patient List, Asthma

2.10.6 Asthma Quality of Care

Denominators

Active Clinical patients ages 5–56 with persistent asthma within the year prior to the beginning of the Report Period and during the Report Period, without a documented history of emphysema or chronic obstructive pulmonary disease (COPD). Broken down by age groups (5–9, 10–17, and 18–56).

User Population patients ages 5–56 with persistent asthma within the year prior to the beginning of the Report Period and during the Report Period, without a documented history of emphysema and chronic COPD. Broken down by age groups (5–9, 10–17, and 18–56).

Numerator

Patients who had at least one dispensed prescription for preferred asthma therapy medication during the Report Period.

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Emphysema definition: Any visit at any time on or before the end of the Report Period with POV codes: 492.*, 506.4, 518.1, 518.2.

COPD definition: Any visit at any time on or before the end of the Report Period with POV codes: 491.20, 491.21, 491.22, 493.2*, 496, 506.4.

Persistent asthma definition:

1. Meeting any of the following four criteria below within the year prior to the beginning of the Report Period *and* during the Report Period:
 - a. At least one visit to Clinic Code 30 (Emergency Medicine) with primary diagnosis 493* (asthma)
 - b. At least one acute inpatient discharge with Primary Diagnosis 493.*. Acute inpatient discharge defined as Service Category of H
 - c. At least four outpatient visits, defined as Service Categories A, S, or O, with primary or secondary diagnosis of 493.* *and* at least two asthma medication dispensing events (see definition below)
 - d. At least four asthma medication dispensing events (see definition below). If the sole medication was leukotriene modifiers, then *must* also have at least one visit with POV 493.* in the same year as the leukotriene modifier (i.e., during the Report Period or within the year prior to the beginning of the Report Period.)
2. Meeting any of the following criteria below:
 - a. Active entry in PCC Problem List for 493.* with Severity of 2, 3 or 4 at any time before the end of the Report Period, or
 - b. Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 2, 3, or 4 documented any time before the end of the Report Period.

Dispensing event definition: One prescription of an amount lasting 30 days or less. For RXs longer than 30 days, divide the days' supply by 30 and round down to convert. For example, a 100-day RX is equal to three dispensing events ($100/30 = 3.33$, rounded down to 3). Also, two different RXs dispensed on the same day are counted as two different dispensing events. Inhalers should also be counted as one dispensing event.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Asthma medication codes for denominator defined with medication taxonomies: BGP HEDIS ASTHMA MEDS, BGP HEDIS ASTHMA LEUK MEDS, BGP HEDIS ASTHMA INHALED MEDS. Medications are: Antiasthmatic Combinations (Dyphylline-Guaifenesin, Guaifenesin-Theophylline, Potassium Iodide-Theophylline), Antibody Inhibitor (Omalizumab), Inhaled Steroid Combinations (Budesonide-Formoterol, Fluticasone-Salmeterol), Inhaled Corticosteroids (Belclomethasone, Budesonide, Flunisolide, Fluticasone CFC Free, Mometasone, Triamcinolone), Lekotriene Modifiers (Montelukast, Zafirlukast, Zileuton), Long-Acting, Inhaled Beta-2 Agonists (Aformoterol, Formoterol, Salmeterol), Mast Cell Stabilizers (Cromolyn, Nedocromil), Methylxanthines (Aminophylline, Dyphylline, Oxtriphylline, Theophylline), Short-Acting, Inhaled Beta-2 Agonists (Albuterol, Levalbuterol, Metaproterenol, Pirbuterol). Medications must not have a comment of RETURNED TO STOCK.

To be included in the numerator, patient must have a non-discontinued prescription for preferred asthma therapy (see list of medications below) during the Report Period.

Preferred asthma therapy medication codes for numerator defined with medication taxonomy: BGP HEDIS PRIMARY ASTHMA MEDS Medications are: Antiasthmatic Combinations (Dyphylline-Guaifenesin, Guaifenesin-Theophylline, Potassium Iodide-Theophylline), Antibody Inhibitor (Omalizumab), Inhaled Steroid Combinations (Budesonide-Formoterol, Fluticasone-Salmeterol), Inhaled Corticosteroids (Belclomethasone, Budesonide, Flunisolide, Fluticasone CFC Free, Mometasone, Triamcinolone), Lekotriene Modifiers (Montelukast, Zafirlukast, Zileuton), Mast Cell Stabilizers (Cromolyn, Nedocromil), Methylxanthines (Aminophylline, Dyphylline, Oxtriphylline, Theophylline). Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of asthmatic patients with preferred asthma therapy medications, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011					Page 292			
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Asthma Quality of Care (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical Pts 5-56									
w/persistent									
asthma	21		10			5			
# w/ preferred asthma									
control med	7	33.3	5	50.0	-16.7	3	60.0	-26.7	
A. Active Clinical									
ages 5-9	11		3			1			
# w/ preferred asthma									
control med	3	27.3	1	33.3	-6.1	1	100.0	-72.7	
B. Active Clinical									
ages 10-17	2		2			1			
# w/ preferred asthma									
control med	2	100.0	2	100.0	+0.0	0	0.0	+100.0	
C. Active Clinical									
ages 18-56	8		5			3			
# w/ preferred asthma									
control med	2	25.0	2	40.0	-15.0	2	66.7	-41.7	
User Pop Pts 5-56									
w/persistent									
asthma	21		11			6			
# w/ preferred asthma									
control med	7	33.3	5	45.5	-12.1	3	50.0	-16.7	
A. User Pop									
ages 5-9	11		4			1			
# w/ preferred asthma									
control med	3	27.3	1	25.0	+2.3	1	100.0	-72.7	
B. User Pop									
ages 10-17	2		2			1			
# w/ preferred asthma									

control med	2 100.0	2 100.0	+0.0	0 0.0	+100.0
C. User Pop ages 18-56	8	5		4	
# w/ preferred asthma control med	2 25.0	2 40.0	-15.0	2 50.0	-25.0

Figure 2-114: Sample Report, Asthma Quality of Care

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Asthma Quality of Care: List of asthmatic patients with preferred asthma therapy medications, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,ZACHARY Asthma 02/02/11	000011	COMMUNITY	F	5	UP; Severity	4 in V
PATIENT12,TINA DANIELLE for 493.00	000012	COMMUNITY	F	6	UP;AC Severity	>1 on PL
PATIENT13,THERESA LYNN Asthma 03/03/11	000013	COMMUNITY	M	47	UP;AC Severity	2 in V
PATIENT36,NATHAN BRADLEY 03/07/11, MONTELUKAST NA 10MG TAB	000014	COMMUNITY	M	16	UP;AC 4 meds	NUM:
PATIENT37,JANELLE MARIE 09/26/08 4 POVS AND 2 MEDS	000015	COMMUNITY	F	50	UP;AC DX ON HOSP/OR ER ON	NUM: 09/19/11, FLUTICASONE PROPIONATE 110MCG INHALER
PATIENT38,THOMAS ELLIS NUM: 02/03/11, MONTELUKAST NA 10MG TAB	000016	COMMUNITY	M	6	UP;AC 4 POVS AND 2 MEDS	

Figure 2-115: Sample Patient List, Asthma Quality of Care

2.10.7 Asthma and Inhaled Steroid Use

Denominators

Active Clinical patients ages 1 or older with persistent asthma or who have had two asthma-related visits during the Report Period. Broken down by age groups (1–4, 5–19, 20–44, 45–64, and >65).

User Population patients ages 1 or older with persistent asthma or who have had two asthma-related visits during the Report Period. Broken down by age groups (1–4, 5–19, 20–44, 45–64, and >65).

Numerator

Patients prescribed an inhaled corticosteroid during the Report Period.

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Denominator Exclusion: Patients with intermittent asthma, defined as any of the following:

- An Active entry in PCC Problem List for 493.* with a Severity of 1 at *any* time before the end of the Report Period, *or*
- Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 1 documented *any* time before the end of the Report period.

Asthma definition:

1. CRS will first search to see if the patient has persistent asthma, which is defined as any of the following:
 - a. An Active entry in PCC Problem List for 493.* with a Severity of 2, 3, or 4 at *any* time before the end of the Report Period, *or*
 - b. Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 2, 3, or 4 documented *any* time before the end of the Report Period.
2. If the patient does not meet any of the above criteria, then CRS will search for two asthma-related visits during the Report Period. Asthma-related visit defined as any primary or secondary POV of *asthma 493.**.

Note: For facilities not using asthma staging (severity assessment) in the PCC Problem List, CRS will rely on visit criteria for this assessment. This will result in patients with intermittent asthma being included in the denominator. The Expert Guideline driven method for managing patients with asthma is by staging them in the PCC Problem List. Doing so will improve the accuracy of the information reported by CRS.

To be included in the numerator, patient must have a non-discontinued prescription for an inhaled corticosteroid during the Report period. Inhaled corticosteroid medications defined with medication taxonomy BGP ASTHMA INHALED STEROIDS. (Medications are: Mometasone (Asmanex), Beclovent, Qvar, Vancenase, Vanceril, Vanceril DS, Bitolerol (Tornalate), Pulmicort, Pulmicort Respules, Pulmicort Turbohaler, Salmeterol/fluticasone (Advair), Triamcinolone (Azmacort), Fluticasone (Flovent), Budesonide-Formoterol (Symbicort).) Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with asthma with inhaled corticosteroid prescription, if any.

Measure Source

HP 2010, 24–7 measure (developmental), National Health Interview Survey (NHIS), CDC, NCHS

Measure Past Performance and Long-term Targets

None

DU	May 25, 2011							Page 295	
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Asthma and Inhaled Steroid Use									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
Active Clinical Ages 1 and older with asthma									
	51		34			21			
# w/ Inhaled Steroid Rx	13	25.5	7	20.6	+4.9	2	9.5	+16.0	
Active Clinical ages 1-4 with asthma									
	9		9			7			
# w/ Inhaled Steroid Rx	2	22.2	1	11.1	+11.1	1	14.3	+7.9	
Active Clinical ages 5-19 with asthma									
	21		11			7			
# w/Inhaled Steroid Rx	5	23.8	3	27.3	-3.5	0	0.0	+23.8	
Active Clinical ages 20-44 with asthma									
	8		8			5			
# w/ Inhaled Steroid Rx	3	37.5	2	25.0	+12.5	0	0.0	+37.5	
Active Clinical ages 45-64 with asthma									
	10		5			0			
# w/ Inhaled Steroid Rx	3	30.0	1	20.0	+10.0	0	0.0	+30.0	
Active Clinical ages 65 and older with asthma									
	3		1			2			
# w/ Inhaled Steroid Rx	0	0.0	0	0.0	+0.0	1	50.0	-50.0	

User Pop Ages 1 and older with asthma	52	37	24
# w/ Inhaled Steroid Rx	13 25.0	7 18.9 +6.1	2 8.3 +16.7

Figure 2-116: Sample Report, Asthma and Inhaled Steroid Use

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Asthma and Inhaled Steroid Use: List of patients with asthma with inhaled corticosteroid prescription, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1, GENEVA FLUTICASONE PROPIONATE 110MCG INHALER	000001	COMMUNITY #1	F	47	UP;AC	2 DXs NUM: 11/06/11
PATIENT2, JACKIE	000002	COMMUNITY #1	F	69	UP;AC	PL 493.00=Pers: 3
PATIENT3, PAULINE	000003	COMMUNITY #1	F	70	UP;AC	V Asthma=Pers
PATIENT4, WILLIAM	000004	COMMUNITY #1	M	7	UP;AC	
PATIENT5, ZACHARY LEE	000005	COMMUNITY #1	M	11	UP;AC	
PATIENT25, JOSEPHINE FLUTICASONE PROPIONATE 110MCG INHALER	000025	COMMUNITY #2	F	4	UP;AC	2 DXs NUM: 06/20/11

Figure 2-117: Sample Patient List, Asthma and Inhaled Steroid Use

2.10.8 Medication Therapy for Persons with Asthma

Denominators

Active Clinical patients ages 5-50 with persistent asthma or who have had two asthma-related visits during the Report Period.

Numerators

Suboptimal Control: Patients who were dispensed more than 3 canisters of a short-acting beta2 agonist inhaler during the same 90-day period during the Report Period.

Absence of Controller Therapy: Patients who were dispensed more than 3 canisters of short acting beta2 agonist inhalers over a 90-day period and who did not receive controller therapy during the same 90-day period.

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Denominator Exclusion: Patients with intermittent asthma, defined as any of the following:

- An Active entry in PCC Problem List for 493.* with a Severity of 1 at *any* time before the end of the Report Period, *or*

- Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 1 documented *any* time before the end of the Report period.

Asthma definition:

1. CRS will first search to see if the patient has persistent asthma, which is defined as any of the following:
 - a. An Active entry in PCC Problem List for 493.* with a Severity of 2, 3, or 4 at *any* time before the end of the Report Period, *or*
 - b. Most recent visit-related asthma entry (i.e., V Asthma) with Severity of 2, 3, or 4 documented *any* time before the end of the Report Period.
2. If the patient does not meet any of the above criteria, then CRS will search for two asthma-related visits during the Report Period. Asthma-related visit defined as any primary or secondary POV of *asthma 493.**.

Note: For facilities not using asthma staging (severity assessment) in the PCC Problem List, CRS will rely on visit criteria for this assessment. This will result in patients with intermittent asthma being included in the denominator. The Expert Guideline driven method for managing patients with asthma is by staging them in the PCC Problem List. Doing so will improve the accuracy of the information reported by CRS.

To be included in the numerator, patient must have one or more non-discontinued prescriptions for short acting Beta2 Agonist inhalers totaling at least four canisters in one 90 day period. Short acting Beta2 Agonist inhaler medications defined with medication taxonomy BGP PQA SABA MEDS. (Medications are: Albuterol, Levalbuterol, Pirbuterol). Medications must not have a comment of RETURNED TO STOCK.

Controller Therapy definition:

At least one non-discontinued prescription of controller therapy medications during the same 90 day period. Controller therapy medications defined with medication taxonomy BGP PQA CONTROLLER MEDS. (Medications are: Beclomethasone, Budesonide, Budesonide-Formoterol, Ciclesonide, Cromolyn, Flunisolide, Fluticasone, Fluticasone-Salmeterol, Formoterol, Mometasone, Mometasone-Formoterol, Montelukast, Nedocromil, Salmeterol, Theophylline, Triamcinolone, Zafirlukast, Zileuton). Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

New topic.

Patient List Description

List of patients with asthma with suboptimal control and controller therapy, if any.

Measure Source

PQA (Pharmacy Quality Alliance)

Measure Past Performance and Long-term Targets

None

DU	May 25, 2011						Page 299	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Medication Therapy for Persons with Asthma								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical ages								
5-50 w/ asthma	35		22			12		
# w/ Suboptimal								
Control	3	8.6	0	0.0	+8.6	0	0.0	+8.6
# w/ Absence of Controller								
Therapy	2	66.7	0	0.0	+66.7	0	0.0	+66.7

Figure 2-118: Sample Report, Medication Therapy for Persons with Asthma

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic						
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Medication Therapy for Persons with Asthma: List of patients with asthma						
with suboptimal control and controller therapy, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,GWEN	000001	COMMUNITY #1	F	5	AC,V Asthma=Pers	
PATIENT2,ALICE	000002	COMMUNITY #1	F	6	AC,PL 493.00=Pers: 2	SABA:
06/15/11 ALBUTEROL 90MCG/INHALATION MDI(4)						
PATIENT3,GENEVA	000003	COMMUNITY #1	F	47	AC,2 DXs	
PATIENT22,MELANIE	000022	COMMUNITY #1	F	47	AC,PL 493.00=Pers: 2	
PATIENT27,RANDALL	000027	COMMUNITY #1	M	6	AC,V Asthma=Pers	
PATIENT28,JASON	000028	COMMUNITY #1	M	6	AC,PL 493.00=Pers: 4	
PATIENT36,WILLIAM	000036	COMMUNITY #1	M	7	AC,2 DXs	
PATIENT42,JEFFERY	000042	COMMUNITY #1	M	7	AC,V Asthma=PersSABA: 03/01/11	
ALBUTEROL 90MCG/INHALATION MDI(4)						
PATIENT45,ALBERT	000045	COMMUNITY #1	M	39	AC,2 DXs	SABA: 05/25/11
ALBUTEROL 90MCG/INHALATION MDI(2) 07/01/11 ALBUTEROL 90MCG/INHALATION MDI (2); CONT:						
06/01/11 SALMETEROL XINAFOATE 50MCG INH POWDER						

Figure 2-119: Sample Patient List, Medication Therapy for Persons with Asthma

2.10.9 Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation

Denominator

Number of visits for User Population patients ages 18 and older diagnosed with community-acquired bacterial pneumonia at an outpatient visit during the Report Period.

Numerators

Number of visits where patients had *oxygen saturation documented* and reviewed.

Number of visits where patients *refused oxygen saturation assessment*.

Number of visits where patients *did not have their oxygen saturation documented* and reviewed.

Logic Description

Age of the patient is calculated at the beginning of the report period.

If a patient has more than one visit for community-acquired bacterial pneumonia during the report period, each visit will be counted as long as it has been more than 45 days since the date of the prior visit. For example, a patient was diagnosed on January 1, 2011 and 35 days later he was diagnosed again with pneumonia. That second diagnosis does not count as a separate visit. However, if the patient were diagnosed again on February 16, 2011 (46 days after onset), that diagnosis counts as a separate visit. Because RPMS does not store the date of onset, visit date will be used as a surrogate for onset date.

CRS uses the following codes and taxonomies to define the denominator and numerators.

	ICD and Other Codes
Community-Acquired Bacterial Pneumonia (Non-CHS outpatient visit, defined as visit type not equal to "C" and service category of "A" for Ambulatory, "S" for Day Surgery, or "O" for Observation)	V POV: 481, 482.0, 482.1, 482.2, 482.30, 482.31, 482.32, 482.39, 482.40, 482.41, 482.42, 482.49, 482.81, 482.82, 482.83, 482.84, 482.89, 482.9, 483.0, 483.1, 483.8, 485, 486, 487.0

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Oxygen Saturation Assessment (any of the arterial blood gas (ABG) or pulse oximetry tests performed at the visit)	94760-94762, 82803, 82805, 82810, or 3028F or 3028F, where 3028F has no modifier of 1P, 2P, 3P, or 8P	V Measurement: O2 Saturation	Yes	BGP CMS ABG TESTS

Refusal of Oxygen Saturation Assessment definition: Patients whose oxygen saturation was not assessed due to a patient refusal of assessment on visit date. Refusal is defined as refusal of any of the tests listed above.

No Assessment definition: Patients whose oxygen saturation was not assessed or refused.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with community-acquired bacterial pneumonia, with oxygen saturation assessment or documented reason for no assessment, if any.

Measure Source

CMS PQRI Measure #57

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 301		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation (con't)									
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from	
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %	
# Pneumonia Visits for									
User Pop Pts 18+	53		12			8			
# Visits w/ O2 Sat									
Assmt	23	43.4	1	8.3	+35.1	2	25.0	+18.4	
# Visits w/ Refusal	4	7.5	0	0.0	+7.5	0	0.0	+7.5	
# Visits w/ No O2									

Sat Assmt	26	49.1	11	91.7	-42.6	6	75.0	-25.9
-----------	----	------	----	------	-------	---	------	-------

Figure 2-120: Sample Report, Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation: List of patients with community-acquired bacterial pneumonia, with oxygen saturation assessment or documented reason for no assessment, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1, GENEVA	000001	COMMUNITY #1	F	27	UP 1) 11/25/11	482.81 NUM: 1)
11/25/11 CPT [3028F]						
PATIENT2, JACKIE	000002	COMMUNITY #1	F	29	UP 1) 05/31/11	482.42 NUM: 1)
05/31/11 None						
PATIENT3, PAULINE	000003	COMMUNITY #1	F	38	UP 1) 05/31/11	482.0; 2)
08/01/11 482.49; 3) 09/17/11						
487.0 NUM: 1) 05/31/11 None; 2) 08/01/11						
CPT [82805]; 3) 09/17/11 CPT [82810]						
PATIENT4, WILLIAM	000004	COMMUNITY #1	M	38	UP 1) 06/15/11	482.89 NUM: 1)
06/15/11 MET BLOOD GASES						
PATIENT5, ZACHARY LEE	000005	COMMUNITY #1	M	36	UP 1) 05/27/11	486. NUM: 1)
05/27/11 O2 MEAS						

Figure 2-121: Sample Patient List, Community-Acquired Pneumonia (CAP): Assessment of Oxygen Saturation

2.10.10 Chronic Kidney Disease Assessment

Denominators

Active Clinical patients ages 18 and older with serum creatinine test during the Report Period.

User Population patients ages 18 and older with serum creatinine test during the Report Period.

Numerators

Patients with Estimated GFR.

- Patients with GFR less than (<) 60.
- Patients with normal GFR (i.e., >=60).

Logic Description

Age is calculated at beginning of the Report Period.

For the GFR <60 numerator, CRS will include GFR results containing a numeric value less than 60 or with a text value of "<60". For the normal GFR (>=60) numerator, CRS will include GFR results containing a numeric value equal to or greater than 60 or with a text value of "60"

CRS uses the following codes and taxonomies to define the denominator and numerators.

	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Creatinine test	82540, 82565-75		Yes	DM AUDIT CREATININE TAX
Estimated GFR test			Yes	BGP GPRA ESTIMATED GFR TAX

Key Logic Changes from CRS Version 11.0

1. Added codes 62238-1 and 50110-4 to LOINC taxonomy for Estimated GFR.

Patient List Description

List of patients with Creatinine test, with GFR and value, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 303	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Chronic Kidney Disease Assessment (con't)								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR	% PERIOD		BASE %
Active Clinical Pts								
=> 18 with Serum								
Creatinine test	271		258			221		
w/Est GFR	189	69.7	33	12.8	+57.0	16	7.2	+62.5
A. # w/ GFR <60	52	19.2	30	11.6	+7.6	15	6.8	+12.4
B. # w/Normal GFR								

(>=60)	137	50.6	2	0.8	+49.8	0	0.0	+50.6
User Pop Pts =>18 with Serum Creatinine	331		311			262		
# w/ Est GFR	221	66.8	33	10.6	+56.2	16	6.1	+60.7
A. # w/GFR <60	55	16.6	30	9.6	+7.0	15	5.7	+10.9
B. # w/Normal GFR (>=60)	165	49.8	2	0.6	+49.2	0	0.0	+49.8

Figure 2-122: Sample Report, Chronic Kidney Disease Assessment

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Chronic Kidney Disease Assessment: List of patients with Creatinine test, with GFR and value, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,SHERISA	000001	COMMUNITY #1	F	18	UP;AC	08/16/11, GFR: 78	
PATIENT2,CAITLYN	000002	COMMUNITY #1	F	22	UP;AC	07/09/11, GFR: >60	
PATIENT3,HALEY DEBRA	000003	COMMUNITY #1	F	25	UP;AC		
PATIENT4,HELENE MARIE	000004	COMMUNITY #1	F	29	UP;AC	10/24/11, GFR: 0.2	
PATIENT5,MARTHA	000005	COMMUNITY #1	F	30	UP;AC	02/28/11, GFR: <60	
PATIENT6,PAULA KAYE	000006	COMMUNITY #1	F	34	UP;AC		
PATIENT7,KATHLEEN	000007	COMMUNITY #1	F	38	UP;AC	02/09/11, GFR: 85	

Figure 2-123: Sample Patient List, Chronic Kidney Disease Assessment

2.10.11 Prediabetes/Metabolic Syndrome

Denominators

Active Clinical patients ages 18 and older diagnosed with prediabetes/metabolic syndrome without a documented history of diabetes.

User Population patients ages 18 and older diagnosed with prediabetes/metabolic syndrome without a documented history of diabetes.

Numerators

All Screenings: Patients with all screenings (BP, LDL, fasting glucose, nephropathy screening, tobacco screening, BMI, lifestyle counseling, and depression screening).

BP Assessed: Patients with Blood Pressure documented at least twice during the Report Period.

LDL Assessed: Patients with LDL completed, regardless of result, during the Report Period.

Fasting Glucose Assessed: Patients with fasting glucose test, regardless of result, during the Report Period.

Nephropathy Assessed: Patients with nephropathy assessment, defined as an estimated GFR with result *and* a quantitative urinary protein assessment during the Report period *or* with evidence of diagnosis and/or treatment of ESRD at any time before the end of the Report period.

Tobacco Use Assessed: Patients who have been screened for tobacco use during the Report Period.

BMI Available: Patients for whom a BMI could be calculated.

Note: This numerator does *not* include refusals.

Lifestyle Counseling: Patients who have received any lifestyle adaptation counseling, including medical nutrition therapy, or nutrition, exercise or other lifestyle education during the Report Period.

Depression Screening: Patients screened for depression or diagnosed with a mood disorder at any time during the Report period, including documented refusals in past year.

Logic Description

Age is calculated at beginning of the Report Period.

Prediabetes/Metabolic Syndrome defined as:

1. Diagnosis of prediabetes/metabolic syndrome, defined as: Two visits during the Report Period with POV 277.7, *or*
2. One each of at least three different conditions listed below, occurring during the Report Period except as otherwise noted:
 - BMI \Rightarrow 30 *or* Waist Circumference >40 inches for men or >35 inches for women
 - Triglyceride value ≥ 150
 - HDL value <40 for men or <50 for women
 - Patient diagnosed with hypertension *or* mean BP value \Rightarrow 130/85 where systolic is \Rightarrow 130 *or* diastolic is \Rightarrow 85
 - Fasting Glucose value \Rightarrow 100 *and* <126

Note: Waist circumference and fasting glucose values will be checked last.

Definition for patients without diabetes: No diabetes diagnosis ever (POV 250.00–250.93).

BMI definition: CRS calculates BMI at the time the report is run, using NHANES II. For 18 and under, a height and weight must be taken on the same day any time during the Report Period. For 19 through 50, height and weight must be recorded within last 5 years, not required to be on the same day. For over 50, height and weight within last 2 years, not required to be recorded on same day.

BP definition: CRS uses mean of last three BPs documented on non-ER visits during the Report Period. If three BPs are not available, uses mean of last two non-ER BPs. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean Systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean Diastolic value is calculated by adding the diastolic values from the last three (or two) blood pressures and dividing by three (or two).

For the BP documented numerator, if CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F–3080F documented on non-ER visit during the Report Period.

Hypertension: Diagnosis of (POV or problem list) 401.* occurring prior to the Report Period, and at least one hypertension POV during the Report Period.

Nephropathy Assessment Definition:

- Estimated GFR with result during the Report Period *and* Quantitative Urinary Protein Assessment during the Report Period

Note: Be sure and check with your laboratory supervisor that the names you add to your taxonomy reflect quantitative test values

- ESRD diagnosis/treatment (see table for codes)

CRS uses the following codes and taxonomies to define the denominator and numerators.

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Triglyceride (requires a non-null, numeric result)			Yes	DM AUDIT TRIGLYCERIDE TAX
HDL (requires a non-null, numeric result)			Yes	DM AUDIT HDL TAX

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Fasting Glucose- <i>Denominator Definition</i> (requires a non-null, numeric result)			Yes	DM AUDIT FASTING GLUCOSE TESTS
Fasting Glucose- <i>Numerator Definition</i>		V POV: 790.21	Yes	DM AUDIT FASTING GLUCOSE TESTS
LDL	80061, 83700, 83701, 83704, 83715 (old code), 83716 (old code), 83721, 3048F, 3049F, 3050F		Yes	DM AUDIT LDL CHOLESTEROL TAX
Estimated GFR			Yes	BGP GPRA ESTIMATED GFR TAX
Quantitative Urinary Protein Assessment	82042, 82043, or 84156		Yes	BGP QUANT URINE PROTEIN
End Stage Renal Disease	36145, 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831- 36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951-90970 or old codes 90918- 90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, G0257, G0308-G0327 (old codes), G0392 (old code), G0393 (old code), or S9339	V POV: 585.5, 585.6, V42.0, V45.1 (old code), V45.11, V45.12, or V56.* V Procedure: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93-39.95, 54.98, or 55.6*		

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Tobacco Screening	D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455-G8457 (old codes), G8402 (old code) or G8453 (old code)	Any health factor for category Tobacco, TOBACCO (SMOKING), TOBACCO (SMOKELESS – CHEWING/DIP), or TOBACCO V POV or current Active Problem List: 305.1, 305.1* (old codes), 649.00-649.04, V15.82 Patient education codes: containing "TO-" or "-TO" or "-SHS", 305.1, 305.1* (old codes), 649.00-649.04, V15.82, D1320, 99406, 99407, G0375 (old code), G0376 (old code), 1034F, 1035F, 1036F, 1000F, G8455-G8457 (old codes), G8402 (old code) or G8453 (old code) Dental code: 1320		
Lifestyle Counseling – Medical Nutrition Counseling	97802-97804, G0270, G0271	Primary or secondary provider codes: 07, 29 Clinic codes: 67 (dietary) or 36 (WIC)		
Lifestyle Counseling – Nutrition Education		V POV: V65.3 dietary surveillance and counseling <i>or</i> Patient education codes: ending "-N" (nutrition) or "-MNT" (medical nutrition therapy) (or old code "-DT" (diet)) or containing V65.3, 97802-97804, G0270, or G0271.		

Test	CPT Codes	ICD and Other Codes	LOINC Codes	Taxonomy
Lifestyle Counseling – Exercise Education		V POV: V65.41 exercise counseling <i>or</i> Patient education codes: ending “-EX” (exercise) or containing V65.41.		
Lifestyle Counseling – Related Exercise and Nutrition Education		Patient education codes: ending “-LA” (lifestyle adaptation) or containing “OBS-” (obesity) or 278.00 or 278.01.		
Depression Screening		V Exam: Exam Code 36 V POV: V79.0 CPT: 1220F BHS Problem Code: 14.1 (Screening for Depression) V Measurement in PCC or BHS: PHQ2 or PHQ9 Refusals: Exam Code 36		
Mood Disorders		At least two visits in PCC or BHS during Report Period for: Major Depressive Disorder, Dysthymic Disorder, Depressive Disorder NOS, Bipolar I or II Disorder, Cyclothymic Disorder, Bipolar Disorder NOS, Mood Disorder Due to a General Medical Condition, Substance-induced Mood Disorder, or Mood Disorder NOS (see codes below). V POV: 296.*, 291.89, 292.84, 293.83, 300.4, 301.13, or 311 BHS POV: 14, 15		

Key Logic Changes from CRS Version 11.0

1. Added CPT code 36147 to ESRD definition.
2. Added CPT code 1220F to Depression Screening definition.
3. Added CPT codes 0001F and 2000F to BP Documented definition.
4. Removed refusals from BMI numerator and logic.
5. Added codes 62238-1 and 50110-4 to LOINC taxonomy for Estimated GFR.
6. Added codes 57937-5, 57936-7, 56133-2, 56132-4, 54471-8, 54470-0, 54469-2, 54468-4, and 54467-6 to LOINC taxonomy for HDL.
7. Added code 43396-1 to LOINC taxonomy for LDL.
8. Added codes 56553-1, 57369-1, 58448-2, 58992-9, and 59159-4 to LOINC taxonomy for Quantitative Urine Protein.
9. For Tobacco Screening, noted that G8453, G8455, G8456, G8457, and G8402 are old codes.

Patient List Description

List of patients age 18 and older with Prediabetes/Metabolic Syndrome with assessments received, if any. The denominator column displays the condition the patient met, either the diagnosis of 277.7 or the 3 conditions the patient met (e.g. BMI=35,TG=155,HDL=35).

Measure Source

“IHS Guidelines for Care of Adults with Prediabetes and/or the Metabolic Syndrome in Clinical Settings (April 2005)”

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 309
*** IHS 2011 Selected Measures with Community Specified Report ***							
DEMO INDIAN HOSPITAL							
Report Period: Jan 01, 2011 to Dec 31, 2011							
Previous Year Period: Jan 01, 2010 to Dec 31, 2010							
Baseline Period: Jan 01, 2000 to Dec 31, 2000							

Prediabetes/Metabolic Syndrome							
	REPORT	%	PREV YR	%	CHG from	BASE	%
	PERIOD		PERIOD		PREV YR %	PERIOD	BASE %
Active Clinical Pts =>18							
w/PreDiabetes/							

Met Syn	50		40			29		
# w/ All screenings	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# w/ BP documented	50	100.0	34	85.0	+15.0	27	93.1	+6.9
# w/LDL done	34	68.0	27	67.5	+0.5	18	62.1	+5.9
# w/ fasting glucose	1	2.0	1	2.5	-0.5	0	0.0	+2.0
# w/ est GFR & quant UP assmt or w/ ESRD	9	17.6	1	2.4	+15.2	0	0.0	+17.6
# w/Tobacco Screening w/in 1 yr	48	96.0	33	82.5	+13.5	21	72.4	+23.6
# w/BMI calculated -No Refusals	48	96.0	40	100.0	-4.0	29	100.0	-4.0
# w/lifestyle adaptation counseling	24	48.0	15	37.5	+10.5	8	27.6	+20.4
# w/Depression screening, DX, or refusal	9	18.0	1	2.5	+15.5	1	3.4	+14.6

Figure 2-124: Sample Report, Prediabetes/Metabolic Syndrome

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Prediabetes/Metabolic Syndrome: List of patients 18 and older with
 Prediabetes/Metabolic Syndrome with assessments received, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,HALEY DEBRA	000001	COMMUNITY #1	F	25	UP,AC BMI=33.64; TRIG=271; HDL=45.4; HTN DX: 01/13/11NUM: 2 BPs - CPT 3074F//;TOB SCRN: 01/13/11 305.1-DP;BMI: 33.64;LIFE: 12/12/11:97804-HM	
PATIENT2,CYNTHIA	000002	COMMUNITY #1	F	36	UP;AC BMI=37.08; TRIG=242; HDL=41.1NUM: 2 BPs;LDL: 07/16/11 111;TOB SCRN: 11/18/11 NEVER SMOKED;BMI: 37.08	
PATIENT3,ABIGAIL	000003	COMMUNITY #1	F	39	UP;AC TRIG=150; HDL=38.5; HTN DX: 08/06/11; BP=142/76NUM: 2 BPs;LDL: 08/06/11 108;GFR: 12/02/11 & QUANT UP: MICROALBUMIN-12/02/11;TOB SCRN: 03/25/11 NEVER SMOKED;BMI: 29.65;LIFE: 12/03/11:Prv: 29	
PATIENT4,ANNA LINDA	000004	COMMUNITY #1	F	44	UP BMI=34.87; TRIG=286; HDL=34.4; BP=138/68NUM: 2 BPs;ESRD: V42.0-05/25/11;TOB SCRN: 11/10/11 G8453-FU;BMI: 34.87	
PATIENT5,DARLENE T	000005	COMMUNITY #1	F	54	UP;AC TRIG=260; HDL=36.0; HTN DX: 11/15/11; BP=134/56NUM: 2 BPs;TOB SCRN: 12/18/11 NEVER SMOKED;BMI: 24.75;LIFE: 11/07/11:ELD-N SN	

Figure 2-125: Sample Patient List, Prediabetes/Metabolic Syndrome

2.10.12 Proportion of Days Covered by Medication Therapy

Denominators

Active Clinical patients ages 18 and older who had two or more prescriptions for *beta-blockers* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *ACEI/ARBs* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *calcium channel blockers (CCB)* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *biguanides* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *sulfonylureas* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *thiazolidinediones* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *statins* during the Report Period.

Active Clinical patients ages 18 and older who had two or more prescriptions for *antiretroviral agents* during the Report Period.

Numerators

Patients with proportion of days covered (PDC) $\geq 80\%$ during the Report Period.

Patients with a gap in medication therapy ≥ 30 days.

Patients with proportion of days covered (PDC) $\geq 90\%$ during the Report Period.

Logic Description

Age is calculated at the beginning of the report period.

To be included in the denominator, patients must have at least two prescriptions for that particular type of medication on two unique dates of service at any time during the Report Period. Medications must not have a comment of RETURNED TO STOCK.

The Index Prescription Start Date is the date when the medication was first dispensed within the Report Period. This date must be greater than 90 days from the end of the Report Period to be counted in the denominator.

The medications in the measures are defined with medication taxonomies: BGP PQA BETA BLOCKER MEDS, BGP PQA ACEI ARB MEDS, BGP PQA CCB MEDS, BGP PQA BIGUANIDE MEDS, BGP PQA SULFONYLUREA MEDS, BGP PQA THIAZOLIDINEDIONE MEDS, BGP PQA STATIN MEDS, BGP PQA ANTIRETROVIRAL MEDS.

For each PDC numerator:

Proportion of days covered = # of days the patient was covered by at least one drug in the class / # of days in the patient's measurement period.

Measurement Period definition:

The patient's measurement period is defined as the number of days between the Index Prescription Start Date and the end of the Report Period. When calculating the number of days the patient was covered by at least one drug in the class, if prescriptions for the same drug overlap, the prescription start date for the second prescription will be adjusted to be the day after the previous fill has ended.

Note: If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e. visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4.

Example of Proportion of Days Covered:

- Report Period: Jan 1 - Dec 31, 2011
- 1st Rx is Index Rx Start Date: 3/1/11, Days Supply=90
- Rx covers patient through 5/29/11
- 2nd Rx: 5/26/11, Days Supply=90
- Rx covers patient through 8/27/11
- 3rd Rx: 9/11/11, Days Supply=180
- Gap = (9/11/11 - 8/27/11) = 15 days
- Rx covers patient through 3/8/12
- Patient's measurement period: 3/1/11 through 12/31/11 = 306 Days
- Days patient was covered: 3/1/11 through 8/27/11 + 9/11/11 through 12/31/11 = 292 Days
- PDC = 292 / 306 = 95%

For each Gap numerator:

CRS will calculate whether a gap in medication therapy of 30 or more days has occurred between each consecutive medication dispensing event during the Report Period. A gap is calculated as the days not covered by the days supply between consecutive medication fills.

Example of Medication Gap >=30 Days:

- Report Period: Jan 1 - Dec 31, 2011
- 1st Rx: 4/1/11, Days Supply=30
- Rx covers patient through 4/30/11
- 2nd Rx: 7/1/11, Days Supply=90
- Gap #1 = (7/1/11 - 4/30/11) = 61 days
- Rx covers patient through 9/28/11
- 3rd Rx: 10/1/11, Days Supply=90
- Gap #2 = (10/1/11 - 9/28/11) = 2 days
- Rx covers patient through 12/29/11
- Gap #1 >=30 days, therefore patient will be included in the numerator for that medication.

Key Logic Changes from CRS Version 11.0

New topic.

Patient List Description

List of patients 18 and older prescribed medication therapy medication with proportion of days covered and gap days.

Measure Source

PQA (Pharmacy Quality Alliance)

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011										Page 314		
*** IHS 2011 Selected Measures with Community Specified Report ***													
DEMO INDIAN HOSPITAL													
Report Period: Jan 01, 2011 to Dec 31, 2011													
Previous Year Period: Jan 01, 2010 to Dec 31, 2010													
Baseline Period: Jan 01, 2000 to Dec 31, 2000													

Proportion of Days Covered by Medication Therapy													
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from					
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %					
Active Clinical Pts													
w/beta-blockers	53		44			37							
# w/ PDC >=80%	31	58.5	24	54.5	+3.9	22	59.5	-1.0					

# w/ gap >=30 days	27	50.9	23	52.3	-1.3	15	40.5	+10.4
Active Clinical Pts w/ACEI/ARBs	97		80			74		
# w/ PDC >=80%	60	61.9	55	68.8	-6.9	39	52.7	+9.2
# w/ gap >=30 days	45	46.4	32	40.0	+6.4	42	56.8	-10.4
Active Clinical Pts w/ CCBs	56		48			55		
# w/ PDC >=80%	34	60.7	30	62.5	-1.8	37	67.3	-6.6
# w/ gap >=30 days	30	53.6	20	41.7	+11.9	22	40.0	+13.6
Active Clinical Pts w/ biguanides	30		26			11		
# w/ PDC >=80%	12	40.0	18	69.2	-29.2	3	27.3	+12.7
# w/ gap >=30 days	19	63.3	10	38.5	+24.9	8	72.7	-9.4
Active Clinical Pts w/ sulfonylureas	7		6			7		
# w/ PDC >=80%	3	42.9	1	16.7	+26.2	4	57.1	-14.3
# w/ gap >=30 days	5	71.4	5	83.3	-11.9	3	42.9	+28.6
Active Clinical Pts w/ thiazolidinediones	20		15			4		
# w/ PDC >=80%	13	65.0	10	66.7	-1.7	2	50.0	+15.0
# w/ gap >=30 days	8	40.0	7	46.7	-6.7	1	25.0	+15.0
Active Clinical Pts w/ statins	60		49			37		
# w/ PDC >=80%	43	71.7	33	67.3	+4.3	25	67.6	+4.1
# w/ gap >=30 days	24	40.0	19	38.8	+1.2	16	43.2	-3.2
Active Clinical Pts w/ antiretroviral agents	2		0			1		
# w/ PDC >=90%	1	50.0	0	0.0	+50.0	0	0.0	+50.0

Figure 2-126: Sample Report, Proportion of Days Covered by Medication Therapy

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
 PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Proportion of Days Covered by Medication Therapy: List of patients 18 and older prescribed medication therapy medication with proportion of days covered and gap days.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
Patient75,PAULA KAY	000075	COMMUNITY #1	F	34	AC	CCB: IXRD: 03/24/11
[282] Days=228 >80 GAP=52						
Patient76,CRSCT	000076	COMMUNITY #1	F	36	AC	CCB: IXRD: 06/01/11
[213] Days=180 >80 GAP=31						

Patient77,CRSAC	000077	COMMUNITY #1	F	44	AC	ACEI/ARB: IXRD:
06/05/11 [209]	Days=60	<80	GAP=118			
Patient78,DEBORA ELLEN	000078	COMMUNITY #1	F	45	AC	BB: IXRD: 07/23/11
[161]	Days=126	<80;	ACEI/ARB: IXRD: 01/29/11	[336]	Days=267	<80
01/29/11 [336]	Days=272	>80				
Patient79,STELLA LYNN	000079	COMMUNITY #1	F	46	AC	BB: IXRD: 01/21/11
[344]	Days=299	>80	GAP=38;	CCB: IXRD: 01/21/11	[344]	Days=299
Patient80,TARA MARIE	000080	COMMUNITY #1	F	51	AC	BB: IXRD: 08/25/11
[128]	Days=56	<80	GAP=70;	ACEI/ARB: IXRD: 01/15/11	[350]	Days=314
01/15/11 [350]	Days=218	<80	GAP=103			
Patient81,CRSNK	000081	COMMUNITY #1	F	51	AC	SULF: IXRD:
09/01/11 [121]	Days=120	>80				

Figure 2-127: Sample Patient List, Proportion of Days Covered by Medication Therapy

2.10.13 Medications Education

Denominators

Active Clinical patients with medications dispensed at their facility during the Report Period.

All User Population patients with medications dispensed at their facility during the Report Period.

Numerator

Patients who were provided patient education about their medications in any location.

Patients who refused patient education about their medications in any location.

Logic Description

Patients receiving medications at their facility are identified by any entry in the VMed file for your facility. The purpose of this definition is to ensure that sites are not being held responsible for educating patients about medications received elsewhere that may be recorded in RPMS. CRS assumes that the appropriate facility is the one the user has logged onto to run the report.

Note: If a site's system identifier, i.e., ASUFAC code, has changed during the period between the Baseline start date and the Current Year end date, due to compacting/contracting or other reasons, your report may display zeros (0s) or very low counts for some time periods.

CRS uses the following patient education codes to define the numerator:

Medication Education	Any Patient Education code containing "M-" or "-M" (medication) or DMC-IN (Diabetes Medicine–Insulin) FP-DPO (Family Planning–Depot Medroxyprogesterone Injections) FP-OC (Family Planning–Oral Contraceptives) FP-TD (Family Planning–Transdermal (Patch)) ASM-NEB (Asthma–Nebulizer) ASM-MDI (Asthma–Metered Dose Inhalers) PL-NEB (Pulmonary Disease–Nebulizer) PL-MDI (Pulmonary Disease–Metered Dose Inhalers)
Refusal	Any refusal in the past year with Patient Education codes containing "M-" or "-M" or PFE codes DMC-IN, FP-DPO, FP-OC, ASM-NEB, ASM-MDI, PL-NEB, PL-MDI, or FP-TD In the past year, any Patient Education code containing "M-" or "-M" or PFE codes DMC-IN, FP-DPO, FP-OC, ASM-NEB, ASM-MDI, PL-NEB, PL-MDI, or FP-TD with a level of understanding of "refused".

Key Logic Changes from CRS Version 11.0

1. Added refusal measure and definition to logic.

Patient List Description

List of patients receiving medications with med education, if any.

Measure Source

None

Measure Past Performance and Long-Term Targets

Measure	Target
<i>IHS 2020 Goal</i>	75.0%

DU	May 25, 2011	Page 316
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Medications Education (con't)		
REPORT PERIOD	% PREV YR PERIOD	% CHG from BASE % PERIOD
Active Clinical Pts receiving		

medications	757		631			592		
# receiving								
medication educ	490	64.7	268	42.5	+22.3	81	13.7	+51.0
# refusals	3	0.4	0	0.0	+0.4	0	0.0	+0.4
User Pop Pts receiving								
medications	981		800			753		
# receiving								
medication educ	599	61.1	306	38.3	+22.8	87	11.6	+49.5
# refusals	3	0.3	0	0.0	+0.3	0	0.0	+0.3

Figure 2-128: Sample Report, Medications Education

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease								
Medications Education: List of patients receiving medications with med education or refusal, if any								
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR		

PATIENT1,ANDREA MARY	000001	COMMUNITY #1	F	0	UP;AC			
PATIENT2,VIRGINIA A	000002	COMMUNITY #1	F	0	UP	08/06/11	HTN-M	
PATIENT3,MICHAELA	000003	COMMUNITY #1	F	0	UP	03/10/11	M-I	
PATIENT4,MISTY	000004	COMMUNITY #1	F	5	UP;AC	05/16/11	M-DI	
PATIENT5,RITA ANN	000005	COMMUNITY #1	F	15	UP;AC	07/05/11	M-I	
PATIENT6,DIANE LOUISE	000006	COMMUNITY #1	F	15	UP	08/21/11	M-I	
PATIENT7,ALICIA	000007	COMMUNITY #1	F	15	UP;AC			
PATIENT8,ALYSHA	000008	COMMUNITY #1	F	16	UP;AC	02/15/11	Refused	
M-DI								
PATIENT9,SHELLY	000009	COMMUNITY #1	F	18	UP;AC	03/12/11	PP-M	

Figure 2-129: Sample Patient List, Medications Education

2.10.14 Public Health Nursing

Patient-Related Measures

Denominator

All User Population patients.

Numerators

For User Population only, the number of patients in the denominator served by Public Health Nurses (PHNs) in any setting, including Home.

For User Population only, the number of patients in the denominator served by a PHN driver/interpreter in any setting.

For User Population only, the number of patients in the denominator served by PHNs in a HOME setting.

For User Population only, the number of patients in the denominator served by a PHN driver/interpreter in a HOME Setting.

Visit-Related Measures

Denominators

Number of visits to User Population patients by PHNs in any setting, including Home

- Number of visits to patients ages 0–28 days (Neonate) in any setting.
- Number of visits to patients ages 29 days–12 months (infants) in any setting.
- Number of visits to patients ages 1–64 years in any setting
- Number of visits to patients ages 65 and older (Elders) in any setting
- Number of PHN driver/interpreter (Provider Code 91) visits

Number of visits to User Population patients by PHNs in Home setting

- Number of Home visits to patients age 0–28 days (Neonate)
- Number of Home visits to patients age 29 days to 12 months (Infants)
- Number of Home visits to patients ages 1–64 years
- Number of Home visits to patients aged 65 and over (Elders).
- Number of PHN driver/interpreter (Provider Code 91) visits in a HOME setting.

Numerator

No numerator: count of visits only

Logic Description

PHN visit is defined as any visit with primary or Secondary Provider Code 13 or 91. Home visit defined as: (1) Clinic 11 and a primary or Secondary Provider Code 13 or 91 or (2) Location Home (as defined in Site Parameters) and a primary or Secondary Provider Code 13 or 91.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients with PHN visits documented.

Numerator codes in patient list: All PHN = Number of PHN visits in any setting; Home = Number of PHN visits in home setting; Driver All = Number of PHN driver/interpreter visits in any setting; Driver Home = Number of PHN driver/interpreter visits in home setting.

Measure Source

None

Measure Past Performance and Long-Term Targets

	All PHN visits	PHN Home Visits
IHS FY 2005 Performance	438,376	Not Reported
IHS FY 2004 Performance	423,379	192,121
IHS FY 2003 Performance	359,089	160,650
IHS FY 2002 Performance	343,874	156,263

DU	May 25, 2011						Page 317	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Public Health Nursing (con't)								
	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR %	BASE PERIOD	%	CHG from BASE %
All User Population patients	2,896		2,456			2,346		
# served by PHNs in any Setting	13	0.4	13	0.5	-0.1	13	0.6	-0.1
# served by PHN drivers/interpreter - in any Setting	0	0.0	0	0.0	+0.0	0	0.0	+0.0
# served by PHNs in a Home Setting	1	0.0	1	0.0	+0.0	0	0.0	+0.0
# served by PHN drivers/interpreters in Home Setting	0	0.0	0	0.0	+0.0	0	0.0	+0.0
Total # PHN Visits - Any Setting	18		16		+2	19		-1
A. Ages 0-28 days	0		0		+0	0		+0
B. Ages 29 days - 12 months	1		3		-2	0		+1
C. Ages 1-64 years	16		13		+3	19		-3
D. Ages 65+	1		0		+1	0		+1
E. Driver/Interpreter visits - any setting	0		0		+0	0		+0

Total # PHN Visits - Home Setting	3	1	+2	0	+3
A. Ages 0-28 days	0	0	+0	0	+0
B. Ages 29 days- 12 months	0	1	-1	0	+0
C. Ages 1-64 years	3	0	+3	0	+3
D. Ages 65+	0	0	+0	0	+0
E. Driver/interpreter visits - Home Setting	0	0	+0	0	+0

Figure 2-130: Sample Report, Public Health Nursing

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Public Health Nursing: List of patients with PHN visits documented						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,HELENE MARIE	000001	COMMUNITY #1	F	29	UP	2 all PHN; 0 home;
0 driver all; 0 driver home						
PATIENT2,KATHLEEN	000002	COMMUNITY #1	F	38	UP	3 all PHN; 3 home;
0 driver all; 0 driver home						
PATIENT40,ERIKA SUE	000040	COMMUNITY #2	F	37	UP	1 all PHN; 0 home;
0 driver all; 0 driver home						
PATIENT41,DANIEL RAY	000041	COMMUNITY #2	M	0	UP	1 all PHN; 0 home;
0 driver all; 0 driver home						

Figure 2-131: Sample Patient List, Public Health Nursing

2.10.15 Breastfeeding Rates

Denominators

Active Clinical patients who are 30–394 days old

Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of *two months* (45–89 days) (PART Denominator)

Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of *six months* (165–209 days)

Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of *nine months* (255–299 days)

Active Clinical patients who are 30–394 days old who were screened for infant feeding choice at the age of *one year* (350–394 days)

Numerators

Patients who were screened for infant feeding choice *at least once*

Patients who were screened for infant feeding choice at the age of *two months* (45–89 days)

Patients who were screened for infant feeding choice at the age of *six months* (165–209 days)

Patients who were screened for infant feeding choice at the age of *nine months* (255–299 days)

Patients who were screened for infant feeding choice at the age of *one year* (350–394 days)

Patients who, at the age of *two months* (45–89 days), were either exclusively or mostly breastfed (PART Numerator)

Patients who, at the age of *six months* (165–209 days), were either exclusively or mostly breastfed

Patients who, at the age of *nine months* (255–299 days), were either exclusively or mostly breastfed

Patients who, at the age of *one year* (350–394 days), were either exclusively or mostly breastfed

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Infant feeding choice definition: The documented feeding choice from the file V Infant Feeding Choice that is closest to the exact age that is being assessed will be used. For example, if a patient was assessed at 45 days old as 1/2 breastfed and 1/2 formula and assessed again at 65 days old as mostly breastfed, the mostly breastfed value will be used since it is closer to the exact age of two months (i.e., 60 days). Another example is a patient who was assessed at 67 days as mostly breastfed and again at 80 days as mostly formula. In this case, the 67 days value of mostly breastfed will be used. The other exact ages are 180 days for six months, 270 days for nine months, and 365 days for one year.

In order to be included in the age-specific screening numerators, the patient must have been screened at the specific age range. For example, if a patient was screened at six months and was exclusively breastfeeding but was not screened at two months, then the patient will only be counted in the six months numerator.

Key Logic Changes from CRS Version 11.0

1. Changed denominators to ages 30-394 days.

Patient List Description

List of patients 30–394 days old, with infant feeding choice value, if any.

Note: “DO” represents “Days Old.”

Measure Source

HP 2020, MICH-21.4 Exclusive breastfeeding-through 3 months, MICH-21.5 Exclusive breastfeeding-through 6 months.

Measure Past Performance and Long-Term Targets

Performance	Percent
IHS FY 2010 Performance	33%
IHS FY 2008 Performance	28%
<i>HP 2020 goal for breastfeeding through 3 months of age</i>	<i>44.3%</i>
<i>HP 2020 goal for breastfeeding through 6 months of age</i>	<i>23.7%</i>

DU	May 25, 2011						Page 321	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Breastfeeding Rates								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
30-394 days	46		30			34		
# w/infant feeding choice screening	11	23.9	0	0.0	+23.9	1	2.9	+21.0
# w/screening @ 2 mos	4	8.7	0	0.0	+8.7	1	2.9	+5.8
# w/screening @ 6 mos	3	6.5	0	0.0	+6.5	0	0.0	+6.5
# w/screening @ 9 mos	4	8.7	0	0.0	+8.7	0	0.0	+8.7

# w/screening @ 1 yr	3	6.5	0	0.0	+6.5	0	0.0	+6.5
AC Pts 30-394 days screened @ 2 mos (PART)	4		0			1		
# @ 2 mos exclusive/ mostly breastfed (PART)	4	100.0	0	0.0	+100.0	1	100.0	+0.0
AC Pts 30-394 days screened @ 6 mos	3		0			0		
# @ 6 mos exclusive/mostly breastfed	2	66.7	0	0.0	+66.7	0	0.0	+66.7
AC Pts 30-394 days screened at 9 mos	4		0			0		
# @ 9 mos exclusive/mostly breastfed	3	75.0	0	0.0	+75.0	0	0.0	+75.0
AC Pts 30-394 days screened @ 1 yr	3		0			0		
# @ 1 year exclusive/mostly breastfed	2	66.7	0	0.0	+66.7	0	0.0	+66.7

Figure 2-132: Sample Report, Breastfeeding Rates

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Breastfeeding Rates: List of patients 30-394 days old, with infant feeding
choice value, if any.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR
PATIENT1,AMANDA DEBRA	000001	COMMUNITY #1	F	0	AC	:
PATIENT2,LEROY JAMES	000002	COMMUNITY #1	M	0	AC	Scrn: 9 MOS: 269
DO, 10/29/10 MOSTLY BREASTFEEDING						
PATIENT3,TERRY SCOTT	000003	COMMUNITY #1	M	0	AC	:
PATIENT4,ROBERT	000004	COMMUNITY #1	M	1	AC	Scrn: 2 MOS: 48 DO,
01/20/11 EXCLUSIVE BREASTFEEDING; 6 MOS: 178 DO, 05/30/11 EXCLUSIVE BREASTFEEDING; 9						
MOS: 276 DO, 09/05/11 EXCLUSIVE BREASTFEEDING; 1 YR: 382 DO, 12/20/11 MOSTLY						
BREASTFEEDING						
PATIENT11,STEVEN CODY	000011	COMMUNITY #2	M	0	AC	Scrn: 6 MOS: 187
DO, 08/11/11 EXCLUSIVE BREASTFEEDING						
PATIENT12,ANDREW THOMAS	000012	COMMUNITY #2	M	0	AC	Scrn: 9 MOS: 278
DO, 10/16/11 MOSTLY BREASTFEEDING						
PATIENT13,ROBERT	000013	COMMUNITY #2	M	0	AC	:
PATIENT14,RICHARD ABE	000014	COMMUNITY #2	M	0	AC	Scrn: 1 YR: 361 DO,
02/05/11 FORMULA ONLY						
PATIENT15,JEFFREY LYLE	000015	COMMUNITY #2	M	0	AC	:

PATIENT16, JASON EDWARD 000016 COMMUNITY #2 M 0 AC 11/05/11 EXCLUSIVE BREASTFEEDING	Scrn: 1 YR: 383 DO,
--	---------------------

Figure 2-133: Sample Patient List, Breastfeeding Rates

2.10.16 Use of High-Risk Medications in the Elderly

Denominators

Active Clinical patients ages 65 and older. Broken down by gender and age groups (65-74, 75-84, >85).

User Population patients ages 65 and older. Broken down by gender.

Numerators

Patients who received at least one high-risk medication for the elderly during the Report Period. (GPRA Developmental Numerator)

Patients who received at least two different high-risk medications for the elderly during the Report Period. (GPRA Developmental Numerator)

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Medication definitions: High-risk medications for the elderly (i.e., potentially harmful drugs) defined with medication taxonomies:

- BGP HEDIS ANTIANXIETY MEDS (Includes combination drugs) (Aspirin-Meprobamate, Meprobamate)
- BGP HEDIS ANTIEMETIC MEDS ((Scopolamine, Trimethobenzamide)
- BGP HEDIS ANALGESIC MEDS (Includes combination drugs) (Acetaminophen-diphenhydramine, diphenhydramine-magnesium salicylate, Ketorolac)

- BGP HEDIS ANTIHISTAMINE MEDS (Includes combination drugs)
(APAP/dextromethorphan/diphenhydramine,
APAP/diphenhydramine/phenylephrine,
APAP/diphenhydramine/pseudoephedrine, Acetaminophen-diphenhydramine,
Atropine/CPM/hyoscyamine/PE/PPA/scopolamine,
Carbetapentane/diphenhydramine/phenylephrine,
Codeine/phenylephrine/promethazine, Codeine-promethazine, Cyproheptadine,
Dexchlorpheniramine, Dexchlorpheniramine/dextromethorphan/PSE,
Dexchlorpheniramine/guaifenesin/PSE,
Dexchlorpheniramine/hydrocodone/phenylephrine,
Dexchlorpheniramine/methscopolamine/PSE, Dexchlorpheniramine-
pseudoephedrine, Dextromethorphan-promethazine, Diphenhydramine,
Diphenhydramine/hydrocodone/phenylephrine, Diphenhydramine-magnesium
salicylate, Diphenhydramine-phenylephrine, Diphenhydramine-pseudoephedrine,
Hydroxyzine hydrochloride, Hydroxyzine pamoate, Phenylephrine-promethazine,
Promethazine, Tripelemnamine)
- BGP HEDIS ANTIPSYCHOTIC MEDS (Thioridazine, Mesoridazine)
- BGP HEDIS AMPHETAMINE MEDS (Amphetamine-destroamphetamine,
Benzphetamine, Dexmethylphenidate, Dextroamphetamine, Diethylpropion,
Methamphetamine, Methylphenidate, Pemoline, Phendimetrazine, Phenteramine)
- BGP HEDIS BARBITURATE MEDS (Amobarbital, Butabarbital,
Mephobarbital, Pentobarbital, Phenobarbital, Secobarbital)
- BGP HEDIS BENZODIAZEPINE MEDS (Includes combination drugs)
(Amitriptyline-Chlordiazepoxide, Chlordiazepoxide, Chlordiazepoxide-clidinium,
Diazepam, Flurazepam)
- BGP HEDIS CALCIUM CHANNEL MEDS (Nifedipine - short acting only)
- BGP HEDIS GASTRO ANTISPASM MED (Dicyclomine, Propantheline)
- BGP HEDIS BELLADONNA ALKA MEDS (Includes combination drugs)
(Atropine, Atropine/CPM/hyoscyamine/PE/scopolamine,
Atropine/hyoscyamine/PB/scopolamine, Atropine-difenoxin, Atropine-
diphenoxylate, Atropine-edrophonium, Belladonna,
Belladonna/cafeine/ergotamine/pentobarbital,
Belladonna/ergotamine/phenobarbital,
Butabarbital/hyoscyamine/phenazopyridine, Digestive enzymes/hyoscyamine/
phenyltoloxamine, Hyoscyamine, Hyoscyamine/methenam/m-blue/phenyl salicyl,
Hyoscyamine-phenobarbital)
- BGP HEDIS SKL MUSCLE RELAX MED (Includes combination drugs)
(ASA/cafeine/orphenadrine, ASA/carisoprodol/codeine, Aspirin-carisoprodol,
Aspirin-meprobamate, Aspirin-methocarbamol, Carisoprodol, Chlorzoxazone,
Cyclobenzaprine, Metaxalone, Methocarbamol, Orphenadrine)

- BGP HEDIS ORAL ESTROGEN MEDS (Includes combination drugs) (Conjugated estrogen, Conjugated estrogen-medroxyprogesterone, Esterified estrogen, Esterified estrogen-methyltestosterone, Estropipate)
- BGP HEDIS ORAL HYPOGLYCEMIC RX (Chlorpropamide)
- BGP HEDIS NARCOTIC MEDS (Includes combination drugs) (ASA/caffeine/propoxyphene, Acetaminophen-pentazocine, Acetaminophen-propoxyphene, Belladonna-opium, Meperidine, Meperidine-promethazine, Naloxone-pentazocine, Pentazocine, Propoxyphene hydrochloride, Propoxyphene napsylate)
- BGP HEDIS VASODILATOR MEDS (Cyclandelate, Dipyridamole-short acting only, Ergot mesyloid, Isoxsuprine)
- BGP HEDIS OTHER MEDS AVOID ELD (Includes androgens and anabolic steroids, thyroid drugs, and urinary anti-infectives) (Methyltestosterone, Nitrofurantoin, Nitrofurantoin macrocrystals, Nitrofurantoin macrocrystals-monohydrate, Thyroid desiccated).

Note: For each medication, the days supply must be >0. If the medication was started and then discontinued, CRS will recalculate the # Days Prescribed by subtracting the prescription date (i.e., visit date) from the V Medication Discontinued Date. Example: Rx Date=11/15/2011, Discontinued Date=11/19/2011, Recalculated # Days Prescribed=4. Medications must not have a comment of RETURNED TO STOCK.

Key Logic Changes from CRS Version 11.0

1. Added denominators for age group breakdowns.

Patient List Description

List of patients 65 and older with at least one prescription for a potentially harmful drug.

Measure Source

HEDIS

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011	Page 324
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Use of High-Risk Medications in the Elderly									
	REPORT PERIOD	%	PREV YR PERIOD	%	CHG from PREV YR	BASE % PERIOD	%	CHG from BASE %	
Active Clinical Pts =>65	116		64			65			
# w/exposure to at least 1 high-risk med (GPRA Dev.)	23	19.8	12	18.8	+1.1	18	27.7	-7.9	
# w/exposure to multiple high-risk meds (GPRA Dev.)	11	9.5	1	1.6	+7.9	8	12.3	-2.8	
Male Active Clinical =>65	53		28			27			
# w/exposure to at least 1 high-risk med (GPRA Dev.)	10	18.9	4	14.3	+4.6	6	22.2	-3.4	
# w/exposure to multiple high-risk meds (GPRA Dev.)	4	7.5	0	0.0	+7.5	2	7.4	+0.1	
Female Active Clinical =>65	63		36			38			
# w/exposure to at least 1 high-risk med (GPRA Dev.)	13	20.6	8	22.2	-1.6	12	31.6	-10.9	
# w/exposure to multiple high-risk meds (GPRA Dev.)	7	11.1	1	2.8	+8.3	6	15.8	-4.7	
User Pop Pts =>65	222		149			143			
# w/exposure to at least 1 high-risk med	25	11.3	13	8.7	+2.5	18	12.6	-1.3	
# w/exposure to multiple high-risk meds	11	5.0	2	1.3	+3.6	8	5.6	-0.6	

Figure 2-134: Sample Report, Drugs to be Avoided in the Elderly

DU	May 25, 2011	Page 326
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		
Previous Year Period: Jan 01, 2010 to Dec 31, 2010		
Baseline Period: Jan 01, 2000 to Dec 31, 2000		

Use of High-Risk Medications in the Elderly (con't)		
ACTIVE CLINICAL PATIENTS 65+		

	Age Distribution		
	65-74	75-84	85+
CURRENT REPORT PERIOD			
AC Patients 65+	70	36	10
# w/exposure to at least 1 high-risk med	15	6	2
% w/exposure to at least 1 high-risk med	21.4	16.7	20.0
# w/exposure to multiple high-risk meds	8	2	1
% w/exposure to multiple high-risk meds	11.4	5.6	10.0
PREVIOUS YEAR PERIOD			
AC Patients 65+	39	19	6
# w/exposure to at least 1 high risk med	7	4	1
% w/exposure to at least 1 high-risk med	17.9	21.1	16.7
# w/exposure to multiple high-risk meds	1	0	0
% w/exposure to multiple high-risk meds	2.6	0.0	0.0
CHANGE FROM PREV YR %			
w/exposure to at least 1 high risk med	+3.5	-4.4	+3.3
w/exposure to multiple high-risk meds	+8.9	+5.6	+10.0

Figure 2-135: Sample Report, Drugs to be Avoided in the Elderly

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic
PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease

Use of High Risk Medications in the Elderly: List of patients 65 and older
with at least one high-risk medication.

PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,JONELLE	000001	COMMUNITY #1	F	69	UP;AC	2 drugs: 08/04/11
ESTERIFIED ESTROGENS 0.625MG TAB (ORAL ESTROGEN); 08/04/11 PROPOXYPHENE-N 100MG/APAP 650MG TAB (NARCOTIC)						
PATIENT2,PAULINE	000002	COMMUNITY #1	F	70	UP;AC	1 drug:11/02/11
PROPOXYPHENE-N 100MG/APAP 650MG TAB (NARCOTIC)						
PATIENT3,NADINE	000003	COMMUNITY #1	F	82	UP;AC	2 drugs: 09/25/03
DIAZEPAM 5MG TAB (BENZODIAZEPINE); 09/25/11 PROPOXYPHENE-N 100MG/APAP 650MG TAB (NARCOTIC)						
PATIENT4,JESSE NATHAN	000004	COMMUNITY #1	M	77	UP;AC	1 drug:08/27/11
CYCLOBENZAPRINE HCL 10MG TAB (SKL MUSCLE)						

Figure 2-136: Sample Patient List, Drugs to be Avoided in the Elderly

2.10.17 Functional Status Assessment in Elders

Denominator

Active Clinical patients ages 55 and older. Broken down by gender.

Numerator

Patients screened for functional status at any time during the Report Period.

Logic Description

Age is calculated at the beginning of the Report Period.

Functional status screening definition: Any non-null values in V Elder Care for (1) at least one of the following ADL fields: toileting, bathing, dressing, transfers, feeding, or continence *and* (2) at least one of the following IADL fields: finances, cooking, shopping, housework/chores, medications or transportation during the Report Period.

Key Logic Changes from CRS Version 11.0

None

Patient List Description

List of patients =>55 with functional status codes, if any. The following are the abbreviations used in the Numerator column:

- TLT–Toileting
- BATH–Bathing
- DRES–Dressing
- XFER–Transfers
- FEED–Feeding
- CONT–Continence
- FIN–Finances
- COOK–Cooking
- SHOP–Shopping
- HSWK–Housework/Chores
- MEDS–Medications
- TRNS–Transportation

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 328	
*** IHS 2011 Selected Measures with Community Specified Report ***								
DEMO INDIAN HOSPITAL								
Report Period: Jan 01, 2011 to Dec 31, 2011								
Previous Year Period: Jan 01, 2010 to Dec 31, 2010								
Baseline Period: Jan 01, 2000 to Dec 31, 2000								

Functional Status Assessment in Elders								
	REPORT	%	PREV YR	%	CHG from	BASE	%	CHG from
	PERIOD		PERIOD		PREV YR %	PERIOD		BASE %
Active Clinical Pts								
=>55	271		162			127		
# w/functional status screening	2	0.7	0	0.0	+0.7	0	0.0	+0.7
Male Active Clinical								
=>55	135		77			60		
# w/functional status screening	1	0.7	0	0.0	+0.7	0	0.0	+0.7
Female Active Clinical								
=>55	136		85			67		
# w/functional status screening	1	0.7	0	0.0	+0.7	0	0.0	+0.7

Figure 2-137: Sample Report, Functional Status Assessment in Elders

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Functional Status Assessment in Elders: List of patients => 55 with functional status codes, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,GLENDA JOYCE	000001	COMMUNITY #1	F	57	AC	
PATIENT2,NADINE	000002	COMMUNITY #1	F	61	AC	
PATIENT3,CHARLOTTE MAE	000003	COMMUNITY #1	F	64	AC	YES: 02/24/11 BATH,
CONT, COOK, DRES, FEED, FIN, HSWK, MEDS, SHOP, TLT, TRNS, XFER						
PATIENT4,KATHERINE ANN	000004	COMMUNITY #1	F	66	AC	YES: 07/11/11 BATH,
FIN						
PATIENT5,ANNA MARIE	000005	COMMUNITY #1	F	66	AC	
PATIENT6,DIANA	000006	COMMUNITY #1	F	67	AC	
PATIENT7,PEGGY ANN	000007	COMMUNITY #1	F	70	AC	NO: 05/20/11: FIN

Figure 2-138: Sample Patient List, Functional Status Assessment in Elders

2.10.18 Fall Risk Assessment in Elders

Denominators

Active Clinical patients ages 65 and older. Broken down by gender.

User Population patients ages 65 and older. Broken down by gender.

Numerators

Patients who have been screened for fall risk or with a fall-related diagnosis in the past year.

Note: This numerator does *not* include refusals.

- a. Patients who have been screened for fall risk in the past year
- b. Patients with a documented history of falling in the past year
- c. Patients with a fall-related injury diagnosis in the past year
- d. Patients with abnormality of gait/balance or mobility diagnosis in the past year

Patients with a documented refusal of fall risk screening exam in the past year

Logic Description

Age of the patient is calculated at the beginning of the Report Period.

Fall risk screening/fall related diagnosis is defined as any of the codes in the table below.

	ICD and Other Codes	Exam Code	E Codes (Injury)
Fall Risk Exam	CPT: 1100F, 1101F, 3288F	V Exam: 37 (Fall Risk)	
History of Falling	V POV: V15.88 (Personal History of Fall)		
Fall-related Injury			V POV (Cause Codes #1-3): E880.*, E881.*, E883.*, E884.*, E885.*, E886.*, E888.*
Abnormality of Gait/Balance or Mobility	V POV: 781.2, 781.3, 719.7, 719.70 (old code), 719.75–719.77 (old codes), 438.84, 333.99, 443.9		
Refusal		V Exam: 37 (Fall Risk)	

Key Logic Changes from CRS Version 11.0

1. Removed refusals from main numerator. Refusal numerator is a separate numerator, and no longer a subnumerator.
2. Added CPT codes 1100F, 1101F, and 3288F to Fall Risk Exam definition.

Patient List Description

List of patients 65 years or older with fall risk assessment, if any.

Measure Source

HP 2010 15–28 Reduce hip fractures among older adults.

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011						Page 331		
*** IHS 2011 Selected Measures with Community Specified Report ***									
DEMO INDIAN HOSPITAL									
Report Period: Jan 01, 2011 to Dec 31, 2011									
Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Fall Risk Assessment in Elders (con't)									
Active Clinical Pts									
65+	116			64			65		
# w/ fall risk screen/ Dx-No Refusals	14	12.1	8	12.5	-0.4	8	12.3	-0.2	
A. # w/ fall risk screen	5	4.3	0	0.0	+4.3	0	0.0	+4.3	
B. # w/ history of fall	1	0.9	0	0.0	+0.9	0	0.0	+0.9	
C. # w/ fall injury	2	1.7	1	1.6	+0.2	3	4.6	-2.9	
D. # w/ abnormal gait	6	5.2	7	10.9	-5.8	5	7.7	-2.5	
# w/ refusal	3	2.6	0	0.0	+2.6	0	0.0	+2.6	
Male Active Clinical									
65+	53			28			27		
# w/ fall risk screen/ Dx-No Refusals	6	11.3	3	10.7	+0.6	2	7.4	+3.9	
A. # w/ fall risk screen	2	3.8	0	0.0	+3.8	0	0.0	+3.8	
B. # w/ history of fall	1	1.9	0	0.0	+1.9	0	0.0	+1.9	
C. # w/ fall injury	0	0.0	0	0.0	+0.0	1	3.7	-3.7	
D. # w/ abnormal gait	3	5.7	3	10.7	-5.1	1	3.7	+2.0	
# w/ refusal	1	1.9	0	0.0	+1.9	0	0.0	+1.9	
Female Active Clinical									

65+	63			36			38		
# w/ fall risk screen/ Dx-No Refusals	8	12.7		5	13.9	-1.2	6	15.8	-3.1
A. # w/ fall risk screen	3	4.8		0	0.0	+4.8	0	0.0	+4.8
B. # w/ history of fall	0	0.0		0	0.0	+0.0	0	0.0	+0.0
C. # w/ fall injury	2	3.2		1	2.8	+0.4	2	5.3	-2.1
D. # w/ abnormal gait	3	4.8		4	11.1	-6.3	4	10.5	-5.8
# w/ refusal	2	3.2		0	0.0	+3.2	0	0.0	+3.2

Figure 2-139: Sample Report, Fall Risk Assessment in Elders

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease						
Fall Risk Assessment in Elders: List of patients 65 years or older with fall risk assessment, if any.						
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR

PATIENT1,SHERRY	000001	COMMUNITY #1	F	68	UP;AC	Refused: 03/03/11
exam 37						
PATIENT2,LORETTA LYNN	000002	COMMUNITY #1	F	78	UP	
PATIENT17,NICOLE	000017	COMMUNITY #2	F	71	UP;AC	Abnormal Gait:
11/25/11 443.9						
PATIENT18,VERONICA	000018	COMMUNITY #2	F	72	UP;AC	Screen: 05/01/11
Exam 37						
PATIENT19,STEPHANIE	000019	COMMUNITY #2	F	76	UP;AC	11/10/03
PATIENT87,MICHAEL JOHN	000021	COMMUNITY #3	M	81	UP;AC	Hx of Fall:
07/25/11 Hx of Fall: 07/25/11						
PATIENT88,KENNETH RAY	000028	COMMUNITY #3	M	85	UP;AC	Screen: 03/13/11
CPT 1100F						

Figure 2-140: Sample Patient List, Fall Risk Assessment in Elders

2.10.19 Palliative Care

Denominators

Active Clinical patients ages 18 and older with two or more types of cancer documented during the Report Period. Broken down by gender and age groups (18-54, >55).

Numerators

No denominator; count only. The total number of *Active Clinical patients* with at least one palliative care visit during the Report Period. Broken down by age groups (<18, 18-54, >55).

No denominator; count only. The total number of palliative care visits for *Active Clinical patients* during the Report Period. Broken down by age groups (<18, 18-54, >55).

Patients with at least two palliative care visits during the Report Period.

Logic Description

Age is calculated at the beginning of the Report Period.

Palliative care visit definition: POV V66.7.

Cancer Types:

	ICD Codes
Melanoma	V POV: 172*
Breast	V POV: 174*, 175*, 239.3
Colon	V POV: 153*, 154*, 235.2
Gyn	V POV: 180*, 182*, 183*, 184*, 236.1, 236.2
Prostate	V POV: 185* 236.5
Testes/Male GU	V POV: 186*, 187.3, 187.4, 187.9, 236.4, 236.6
Head and Neck	V POV: 140–149.9, 160*, 161*, 162*, 195.0
Urinary Tract	V POV: 188*, 189*, 236.7, 236.91, 239.4, 239.5
Non-melanomatous skin cancer	V POV: 173*, 238.2
Non-colon GI	V POV: 150–152.9, 155–159.9, 235*, 239.0
Lung	V POV: 162*, 235.9, 239.1
Brain	V POV: 190–192.9, 237.5, 237.6, 239.6
Bones/soft tissue	V POV: 170*, 171*, 238.1, 238.2
Endocrine	V POV: 193, 194*, 237.0, 237.4, 239.7
Pleura/mediastinum	V POV: 163*, 164*
Non-specific site	V POV: 195*, 199*, 238.8, 238.9, 239.8, 239.9
Lymph node spread	V POV: 196*
Secondary cancer	V POV: 196*, 197*

Patient List Description

List of patients with a palliative care visit.

Key Logic Changes from CRS Version 11.0

1. Added new measures and corresponding logic.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011					Page 334				
*** IHS 2011 Selected Measures with Community Specified Report ***										
DEMO INDIAN HOSPITAL										
Report Period: Jan 01, 2011 to Dec 31, 2011										
Previous Year Period: Jan 01, 2010 to Dec 31, 2010										
Baseline Period: Jan 01, 2000 to Dec 31, 2000										

Palliative Care										
	REPORT PERIOD		PREV YR PERIOD		CHG from PREV YR		BASE PERIOD		CHG from BASE	
Total # of Patients										
w/At Least 1 Palliative										
Care Visit										
	40		0		+40		0		+40	
A. Total # of Patients										
<18 w/At Least 1 Palliative										
Care Visit										
	1		0		+1		0		+1	
B. Total # of Patients 18-54										
w/At Least 1 Palliative										
Care Visit										
	25		0		+25		0		+25	
C. Total # of Patients 55+										
w/At Least 1 Palliative										
Care Visit										
	14		0		+14		0		+14	
Total # of Palliative										
Care Visits										
	62		0		+62		0		+62	
A. Total # of Palliative										
Care Visits-Pts										
<18										
	2		0		+2		0		+2	
B. Total # of Palliative										
Care Visits-Pts										
18-54										
	38		0		+38		0		+38	
C. Total # of Palliative										
Care Visits-Pts										
55+										
	22		0		+22		0		+22	
Active Clinical Pts 18+										
w/ 2+ types of										
cancer										
	28		3				3			
# w/ 2+ Palliative										
Care Visits										
	14 50.0		0 0.0		+50.0		0 0.0		+50.0	

Figure 2-141: Sample Report, Palliative Care

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Palliative Care: List of patients with a palliative care visit, if any.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
PATIENT1,JOHN	000012	Community #1	M	57	AC	1 visit: 05/01/11	
PATIENT2,ROBERT	000013	Community #1	M	59	AC,CAN	2 visits: 01/25/11, 05/10/11	
PATIENT3,JAMES	000014	Community #2	M	67	AC,CAN	0 visits:	
PATIENT4,TONYA	000015	Community #3	F	78	AC	1 visit: 06/01/11	
PATIENT5,RITA ANN	000016	Community #3	F	96	AC	2 visits: 06/01/11; 06/07/11	
PATIENT6,Clifford	000017	Community #3	M	24	AC,CAN	3 visits: 01/25/11, 05/10/11, 08/01/11	

Figure 2-142: Sample Patient List, Palliative Care

2.10.20 Annual Wellness Visit

Denominators

Active Clinical patients ages 65 and older. Broken down by gender.

Numerators

Patients with at least one Annual Wellness Exam in the past 15 months.

Logic Description

Age is calculated at the beginning of the Report Period.

Annual Wellness Exam: CPT G0438, G0439, G0402.

Patient List Description

List of patients with an annual wellness visit in the past 15 months.

Key Logic Changes from CRS Version 11.0

New topic.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011	Page 337
*** IHS 2011 Selected Measures with Community Specified Report ***		
DEMO INDIAN HOSPITAL		
Report Period: Jan 01, 2011 to Dec 31, 2011		

Previous Year Period: Jan 01, 2010 to Dec 31, 2010									
Baseline Period: Jan 01, 2000 to Dec 31, 2000									

Annual Wellness Visit									
	REPORT PERIOD		PREV YR PERIOD		CHG from PREV YR	BASE PERIOD	CHG from BASE		
Active Clinical Pts 65+	116		64			65			
# w/ Annual Wellness Exam	2	1.7	0	0.0	+1.7	0	0.0	+1.7	
Male Active Clinical Pts 65+	53		28			27			
# w/ Annual Wellness Exam	0	0.0	0	0.0	+0.0	0	0.0	+0.0	
Female Active Clinical Pts 65+	63		36			38			
# w/ Annual Wellness Exam	2	3.2	0	0.0	+3.2	0	0.0	+3.2	

Figure 2-143: Sample Report, Annual Wellness Visit

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease									
Annual Wellness Visit: List of patients with an annual wellness visit in the past 15 months.									
PATIENT NAME		HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR		

Patient1,DENISE		000001	Community #1	F	65	AC >65	12/31/10	G0402	
Patient2,MELISSA GAYLE		000002	Community #1	F	66	AC >65			
Patient3,JESSICA DAWN		000003	Community #1	F	67	AC >65	02/22/11	G0438	
Patient4,RUTH ALICE		000004	Community #1	F	69	AC >65			
Patient5,BRYSON DEWAY		000005	Community #1	F	72	AC >65			
Patient6,BRITTNEY ANN		000006	Community #1	F	73	AC >65	05/31/11	G0439	
Patient7,MARK		000007	Community #1	M	67	AC >65			
Patient8,HOWIE		000008	Community #1	M	72	AC >65	10/01/10	G0402	

Figure 2-144: Sample Patient List, Annual Wellness Visit

2.10.21 Goal Setting

Denominators

User Population patients who received patient education during the report period.

Numerators

Number of patients who set at least one goal during the Report Period.

Number of patients who met at least one goal during the Report Period.

Logic Description

Patient education codes must be the standard national patient education codes, which are included in the Patient and Family Education Protocols and Codes (PEPC) manual published each year. If codes are found that are not in the table, they will not be reported on (i.e. locally-developed codes).

Numerator Logic:

For Goal Set, the patient education code must have a "GS" value documented during the Report Period.

For Goal Met, the patient education code must have a "GM" value documented during the Report Period but the patient is not required to have set a goal during the Report Period.

Patient List Description

List of User Population patients who received patient education during the Report Period with goal setting information.

Key Logic Changes from CRS Version 11.0

New topic.

Measure Source

None

Measure Past Performance and Long-Term Targets

None

DU	May 25, 2011				Page 338
*** IHS 2011 Selected Measures with Community Specified Report ***					
DEMO INDIAN HOSPITAL					
Report Period: Jan 01, 2011 to Dec 31, 2011					
Previous Year Period: Jan 01, 2010 to Dec 31, 2010					
Baseline Period: Jan 01, 2000 to Dec 31, 2000					

Goal Setting					
	REPORT PERIOD	PREV YR PERIOD	CHG from PREV YR	BASE PERIOD	CHG from BASE
# User Pop w/ Pat Ed	1,024	858		644	

# w/ goal set	8	0.8	0	0.0	+0.8	0	0.0	+0.8
# w/ goal met	6	0.6	0	0.0	+0.6	0	0.0	+0.6

Figure 2-145: Sample Report, Goal Setting

UP=User Pop; AC=Active Clinical; AD=Active Diabetic; AAD=Active Adult Diabetic PREG=Pregnant Female; IMM=Active IMM Pkg Pt; IHD=Active Ischemic Heart Disease							
Goal Setting: List of User Population patients who received patient education during the Report Period with goal setting information.							
PATIENT NAME	HRN	COMMUNITY	SEX	AGE	DENOMINATOR	NUMERATOR	
Patient1,Paula	000001	Community #1	F	34	UP w/Educ	GS: 11/17/11	
Patient2,PENNY	000002	Community #1	F	43	UP w/Educ		
Patient3,RITA	000003	Community #1	F	64	UP w/Educ	GS: 03/13/11	
Patient4,HARRY	000004	Community #1	M	50	UP w/Educ	GM: 10/30/11	
Patient5,ROSS	000005	Community #1	M	55	UP w/Educ		
Patient6,FELIPE	000006	Community #1	M	57	UP w/Educ	GS: 09/10/11	
Patient7,MARK	000007	Community #1	M	67	UP w/Educ		
Patient8,CATHERINE	000008	Community #2	F	72	UP w/Educ	GM: 07/30/11	

Figure 2-146: Sample Patient List, Goal Setting

Contact Information

If you have any questions or comments regarding this distribution, please contact the OIT Help Desk (IHS).

Phone: (505) 248-4371 or (888) 830-7280 (toll free)

Fax: (505) 248-4363

Web: <http://www.ihs.gov/GeneralWeb/HelpCenter/Helpdesk/index.cfm>

Email: support@ihs.gov