


COVID-19 UPDATE

Richard Arakaki, M.D.

Division of Diabetes Treatment and Prevention


Indian Health Service

U.S. COVID-19 Cases by Race N=479,452; May 11, 2020




https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html

COVID-19 Cases: All IHS Reporting



https://www.ihs.gov/coronavirus/

COVID-19 Cases: All IHS Reporting (cont.)



https://www.ihs.gov/coronavirus/

CDC COVIDView Surveillance System Week Ending May 2, 2020

Adaptation of CDC Influenza-like Illness (ILI) to track COVID-19-like Illness (CLI) and provides national percentage of respiratory specimens testing positive for SARS-CoV-2, outpatient or emergency department (ED) visits for potential CLI, cumulative COVID-19-associated hospitalization rates, deaths attributed to COVID-19 and deaths attributed to pneumonia (other than COVID-19 and flu), and current data through week ending April 18th (lag in reporting and analysis).

- Key Points: compared to the previous three weeks
 - Nationally, the percentages of laboratory specimens testing positive for SARS-CoV-2 has decreased.
 - Nationally, visits to outpatient providers and emergency departments (EDs) for illnesses with symptoms consistent with COVID-19 continued to decline.
 - The decrease in the percentage of people presenting for care with ILI and CLI may be due to a decline in COVID-19 illness, which could be in part a result of widespread adoption of social distancing in addition to changes in healthcare seeking behavior.
 - Based on death certificate data, the percentage of deaths attributed to pneumonia, influenza, or COVID-19 (PIC) decreased from 23.6% during week 16 to 21.0% during week 17 to 10.6% during week 18.





Thank you!

