





**Indian Health Service**  
 Division of Epidemiology and Disease Prevention  
 Influenza Surveillance Report  
**IHS Influenza Awareness System (IIAS)**  
**#12 (Week 7: 02/13/2011 – 02/19/2011)**

**Percentage of Visits for Influenza-like Illness (ILI) per week**  
**Overall ILI % (ILI visits per week / Total visits per week)**  
**(ILI) in IHS, Weeks 32 (08/08/2010)**  
**through 07 (02/19/2011)**



All data are preliminary and may change as more reports are received.




### **% Visits for ILI Compared to Previous Week**

<b>IHS Area</b>	<b>% in week 6</b>	<b>% in week 7</b>
NATIONAL	4.2	3.4
Aberdeen	3.4	3.1
Alaska	3.0	1.0
Albuquerque	4.9	4.9
Bemidji	3.3	2.8
Billings	3.0	3.3
California	3.9	3.9
Nashville	3.4	2.2
Navajo	6.4	5.7
Oklahoma	2.5	1.8
Phoenix	4.6	3.5
Portland	2.9	2.7
Tucson	6.9	5.2

## IHS Influenza Vaccination Surveillance

- As of 02/19/2011, sites reporting to the IIAS have administered 389,556 doses (1st and 2nd) of seasonal influenza vaccine to Active Clinical\* patients.

### Influenza Vaccine Coverage, IHS Influenza Awareness System Weeks 32 (08/08/2010) through 07 (02/19/2011), 2010 - 2011



(Active Clinical Population covered by the IIAS)

### Influenza Vaccination Coverage by Age Group

Age Group	Active Clinical Population	1 dose Seasonal Flu Vaccine	% Seasonal flu (1 dose)	2 dose Seasonal Flu Vaccine	% Seasonal flu (2 dose)
0	3,649	93	2.1	34	0.7
6-23 months	22,133	18,754	65.1	6,484	22.5
2-4 years	59,109	26,680	34.9	3,345	4.1
5-17 years	192,016	80,365	31.4	.	.
18-49 years	418,732	134,223	24.7	.	.
50-64 years	148,478	73,592	37.8	.	.
65+ years	80,555	46,079	42.9	.	.
Children (6 months – 17 years)	273,258	125,799	34.9	.	.
Adults (18+)	647,765	253,894	30.0	.	.
All(6 months +)	921,023	379,693	31.4	.	.

\* Active Clinical population is defined as patients who have had at least 2 visits in the last 3 years

## Influenza Vaccination Coverage by Risk Group

<b>Risk Group</b>	<b>Active Clinical Population</b>	<b>1 dose Seasonal Flu Vaccine</b>	<b>% Seasonal flu (1 dose)</b>
Asthma	105,411	21,443	16.0
Diabetes	121,469	45,552	29.2
Pregnant	6,777	2,880	32.3

\* Active Clinical population is defined as patients who have had at least 2 visits in the last 3 years

### Summary

- This report includes data from the following # RPMS databases:

<b>IHS Area</b>	<b># of RPMS Databases Reporting</b>
Aberdeen	18
Alaska	5
Albuquerque	6
Bemidji	2
Billings	8
California	21
Nashville	11
Navajo	6
Oklahoma	19
Phoenix	15
Portland	22
Tucson	2
All Areas	135

- Nationwide, ILI visits are decreased; a trend seen in most IHS Areas.
- Modest increases in vaccination coverage continue.