




Indian Health Service
 Division of Epidemiology and Disease Prevention
Influenza Surveillance Report
IHS Influenza Awareness System (IIAS)
(Week 08: 02/17/2019 – 02/23/2019)


- Influenza-like-illness (ILI) activity in IHS is high at 5.3%, above the IHS baseline of 1.58%
 - All twelve IHS Areas reported having elevated ILI activity
- National ILI activity reported by CDC is high
 - CDC's ILINet ILI activity for week 7 was 5.1 % and is above the ILINet baseline (2.2%) and the IHS baseline (1.58%).
- To date, IHS has administered 301,000 doses of influenza vaccine.
- Overall, 35.5% of the IHS Active Clinical population received an influenza vaccine this season.

All IHS Areas
Influenza-Like Illness (ILI) for Current Surveillance Year and Previous Six Surveillance Years
Percentage of Visits for ILI per week
09/30/2012 to 02/23/2019




Percent ILI visits is number of ILI visits divided by total number of daily outpatient visits.
 The horizontal line is the 2018-2019 IIAS baseline. The ILINet baseline value for 2018-2019 is 2.2.
 Baselines are computed as the mean percentage of ILI patient visits during non-influenza weeks for the previous three seasons, plus two standard deviations.
 Sources: ILINet - CDC Sentinel Surveillance System for Influenza-Like Illness, IIAS - IHS Influenza Awareness System

Percentage of Outpatient Visits for Influenza-Like Illness (ILI)
Percent ILI for Flu Year 2018 - 2019
IIAS vs. ILINet
09/30/2018 to 02/23/2019




Percent ILI visits is number of ILI visits divided by total number of daily outpatient visits.
 The horizontal line is the 2018-2019 IIAS baseline. The ILINet baseline value for 2018-2019 is 2.2.
 Baselines are computed as the mean percentage of ILI patient visits during non-influenza weeks for the previous three seasons, plus two standard deviations.
 Sources: ILINet - CDC Sentinel Surveillance System for Influenza-Like Illness, IIAS - IHS Influenza Awareness System

Influenza-Like Illness (ILI) for Current Surveillance Year and Previous Six Years – Continuous
Percentage of Visits for ILI per week
IIAS vs. ILINet
09/30/2012 to 02/23/2019



Percent ILI visits is number of ILI visits divided by total number of daily outpatient visits.
 The horizontal line is the 2018-2019 IIAS baseline. The ILINet baseline value for 2018-2019 is 2.2.
 Baselines are computed as the mean percentage of ILI patient visits during non-influenza weeks for the previous three seasons, plus two standard deviations.
 Sources: ILINet - CDC Sentinel Surveillance System for Influenza-Like Illness, IIAS - IHS Influenza Awareness System

**Percentage of Influenza-Like Illness Visits by Region
For Week: 08**



Percent Visits for ILI Compared to Previous Week


| IHS Areas | Previous Week (7) | Current Week (8) |
|------------------|--------------------------|-------------------------|
| NATIONAL | 5.1 | 5.3 |
| ALASKA | 6.4 | 4.9 |
| ALBUQUERQUE | 3.8 | 3.7 |
| BEMIDJI | 3.7 | 6.2 |
| BILLINGS | 6.9 | 7.7 |
| CALIFORNIA | 1.3 | 2.0 |
| GREAT PLAINS | 5.3 | 6.6 |
| NASHVILLE | 8.3 | 7.5 |
| NAVAJO | 6.7 | 6.3 |
| OKLAHOMA | 4.5 | 5.0 |
| PHOENIX | 3.7 | 3.5 |
| PORTLAND | 4.1 | 3.1 |
| TUCSON | 1.8 | 2.5 |

Percentage of ILI visits is number of ILI visits divided by total number of daily visits.

IHS Influenza Vaccination Surveillance

As of 02/23/2019, sites reporting to the IIAS have administered 301,171 doses (1st and 2nd) of seasonal influenza vaccine to Active Clinical patients.

All IHS Areas
Cumulative One-Dose Seasonal Influenza Vaccine Coverage
Active Clinical Population
07/01/2018 to 02/23/2019



Influenza Vaccination Coverage by Age Group*

| Age Group | Active Clinical Population ** | Seasonal Flu Vaccine (At Least 1 Dose) *** | % Seasonal Flu (At Least 1 Dose) | Seasonal Flu Vaccine (At Least 2 Doses) | % Seasonal Flu (At Least 2 Doses) | Total # Doses Administered **** |
|------------------------------|----------------------------------|--|-------------------------------------|--|--------------------------------------|------------------------------------|
| 0 - 5 months | 3,590 | 29 | 0.8 | 4 | 0.1 | 33 |
| 6 - 23 months | 16,798 | 10,205 | 60.8 | 2,938 | 17.5 | 13,183 |
| 2 - 4 years | 43,033 | 14,583 | 33.9 | 503 | 1.2 | 15,125 |
| 5 - 17 years | 171,410 | 63,453 | 37.0 | 376 | 0.2 | 64,038 |
| 18 - 49 years | 353,092 | 96,566 | 27.3 | 477 | 0.1 | 97,407 |
| 50 - 64 years | 151,556 | 63,519 | 41.9 | 370 | 0.2 | 64,170 |
| 65 + years | 94,399 | 46,567 | 49.3 | 353 | 0.4 | 47,248 |
| Children (6 months-17 years) | 231,241 | 88,241 | 38.2 | 3,817 | 1.7 | 92,346 |
| Adults (18+ years) | 599,047 | 206,652 | 34.5 | 1,200 | 0.2 | 208,825 |
| All (6 months+) | 830,288 | 294,893 | 35.5 | 5,017 | 0.6 | 301,171 |

* Limited to sites that exported both doses administered along with accompanying Active Clinical population data.

** Active Clinical is defined as patients who have had at least 2 visits in the last 3 years.

*** Number of unique patients with at least 1 dose.

**** Total # Doses Administered = All doses administered to patients including doses not included in coverage calculation due to lack of Active Clinical Population data.

Influenza Vaccine Doses Administered by Influenza Vaccine Type*

| Area | INACTIVE CVX CODES* | CVX 88 FLU-NOS** | CVX 135 FLU-HIGH | CVX 140 FLU-IIV3pf | CVX 141 FLU-IIV3 | CVX 149 FLU-LAIV4 | CVX 150 FLU-IIV4pf | CVX 158 FLU-IIV4 | CVX 161 IIV4pf ped | CVX 168 FLU-IIV3ad | CVX 171 FLU-ccl4pf | CVX 185 FLU-RIV4 | CVX 186 FLU-ccl4 | Proc 99.52 Vac Inf | ICD-9 V04.81 Vac Inf | CPT Billing Codes+ | Cumulative # of Doses Seasonal Flu |
|------|---------------------|------------------|------------------|--------------------|------------------|-------------------|--------------------|------------------|--------------------|--------------------|--------------------|------------------|------------------|--------------------|----------------------|--------------------|------------------------------------|
| AKA | 0 | 48 | 358 | 1 | 30 | 0 | 2,638 | 422 | 199 | 1 | 55 | 10 | 0 | 0 | 0 | 14 | 3,776 |
| ALB | 49 | 290 | 142 | 43 | 1,615 | 1 | 11,005 | 7,489 | 686 | 2,114 | 180 | 37 | 17 | 0 | 0 | 112 | 23,780 |
| BEM | 6 | 53 | 127 | 11 | 213 | 39 | 4,448 | 3,129 | 224 | 336 | 2 | 68 | 0 | 0 | 0 | 40 | 8,696 |
| BIL | 6 | 164 | 1,288 | 1 | 248 | 780 | 12,009 | 3,333 | 292 | 1 | 5 | 7 | 4 | 0 | 0 | 46 | 18,184 |
| CAO | 51 | 1,259 | 1,480 | 3 | 610 | 0 | 3,991 | 1,748 | 370 | 103 | 499 | 1 | 104 | 0 | 0 | 195 | 10,414 |
| GPA | 20 | 347 | 1,066 | 6 | 777 | 0 | 15,781 | 12,169 | 554 | 593 | 404 | 91 | 15 | 0 | 0 | 140 | 31,963 |
| NSA | 52 | 201 | 248 | 5 | 342 | 13 | 5,513 | 5,464 | 504 | 505 | 24 | 32 | 73 | 0 | 0 | 270 | 13,246 |
| NAV | 37 | 418 | 2,627 | 24 | 711 | 2 | 49,219 | 20,512 | 2,165 | 3 | 323 | 46 | 16 | 0 | 0 | 135 | 76,238 |
| OKC | 120 | 3,159 | 1,127 | 10 | 2,269 | 37 | 31,110 | 10,277 | 1,364 | 2,339 | 68 | 41 | 5 | 0 | 0 | 204 | 52,130 |
| PHX | 18 | 1,701 | 828 | 41 | 467 | 0 | 21,302 | 11,182 | 752 | 1,914 | 12 | 191 | 3 | 0 | 0 | 93 | 38,504 |
| POR | 40 | 253 | 611 | 68 | 246 | 160 | 9,355 | 4,792 | 469 | 494 | 184 | 416 | 276 | 0 | 0 | 67 | 17,431 |
| TUC | 3 | 56 | 34 | 4 | 1 | 0 | 7,634 | 108 | 12 | 0 | 0 | 4 | 0 | 0 | 0 | 44 | 7,900 |
| | 402 | 7,949 | 9,936 | 217 | 7,529 | 1,032 | 174,005 | 80,625 | 7,591 | 8,403 | 1,756 | 944 | 513 | 0 | 0 | 1,360 | 302,262 |


* Inactive CVX Codes 15 (FLU-TIVhx), 16 (FLU-WHOLE), 111 (FLU-LAIV3), 123 (FLU-H5N1), 144 (FLU-DERMAL), 151 (FLU-NasNOS), 153 (FLU-cclIIV3), 155 (FLU-RIV3), 160 (H5N1, adj) and 166 (FLU-DERM4) should not be used to document flu vaccines given this flu season.

** CVX Code 88 (FLU-NOS) should only be used to document flu vaccines given elsewhere when flu vaccine type is unknown.

+ CPT Flu codes include: 90654-90662, 90672-90673, 90685-90686, 90724, G0008, G8108.

^ ICD9 codes V04.8 and V06.6 are not displayed on this report and are not used in any of the influenza vaccine coverage calculations.

All IHS Areas
Influenza Vaccine Coverage by Age Groups (Children, Adults, Seniors and All)
% Seasonal Flu (At least 1 dose) for Active Clinical Population
07/01/2018 to 02/23/2019



All IHS Areas
Influenza Vaccine Coverage by Age Groups (Children, Adults, Seniors, and All)
% Seasonal Flu (At least 1 dose) for Active Clinical Population
07/01/2018 to 02/23/2019

| Area | Children (6 mos-17 yrs) | Adults (18+ yrs) | Seniors (65+ yrs) | All (6 mos+) |
|--------------|----------------------------|---------------------|----------------------|-----------------|
| NATIONAL | 38.2 | 34.5 | 49.3 | 35.5 |
| ALASKA | 21.4 | 25.2 | 32.6 | 24.1 |
| ALBUQUERQUE | 38.6 | 38.1 | 52.9 | 38.3 |
| BEMIDJI | 18.7 | 18.1 | 29.2 | 18.2 |
| BILLINGS | 42.9 | 35.5 | 45.2 | 37.8 |
| CALIFORNIA | 17.6 | 19.6 | 29.1 | 19.1 |
| GREAT PLAINS | 35.2 | 31.6 | 48.5 | 32.8 |
| NASHVILLE | 35.3 | 36.7 | 55.3 | 36.3 |
| NAVAJO | 54.4 | 45.6 | 59.1 | 48.1 |
| OKLAHOMA | 32.1 | 30.5 | 49.8 | 30.9 |
| PHOENIX | 38.6 | 36.6 | 51.0 | 37.3 |
| PORTLAND | 43.5 | 38.6 | 56.6 | 39.7 |
| TUCSON | 56.8 | 49.7 | 53.6 | 51.9 |

Number of RPMS Databases Reporting for Week 08, 2019

| IHS Area | # of RPMS Databases Reporting* | # of Facilities Reporting** |
|--------------|--------------------------------|-----------------------------|
| NATIONAL | 132 | 225 |
| ALASKA | 4 | 22 |
| ALBUQUERQUE | 10 | 16 |
| BEMIDJI | 4 | 6 |
| BILLINGS | 12 | 19 |
| CALIFORNIA | 14 | 23 |
| GREAT PLAINS | 16 | 32 |
| NASHVILLE | 20 | 24 |
| NAVAJO | 7 | 19 |
| OKLAHOMA | 17 | 24 |
| PHOENIX | 12 | 20 |
| PORTLAND | 15 | 16 |
| TUCSON | 1 | 4 |

* Exporting databases may include data from multiple facilities.

** Includes only facility categories: Hospital, Health Center, Health Station, Health Location, and Alaska Village Clinic.