



To: Director of IHS Maternal & Child Health Program
IHS Area Chief Medical Officers
IHS Area Immunization Coordinators
Tribal Epidemiology Centers

From: Amy Groom, IHS Immunization Program Manager

Subject: Fourth Quarter FY 2011 Immunization Report (July 1, 2011 – September 30, 2011)

Date: November 22, 2011

The Indian Health Service, Tribal and Urban Indian immunization programs report on the immunization status of American Indian and Alaska Native (AI/AN) children 3-27 months of age, 19 – 35 months of age, and AI/AN adolescents 13 – 17 years of age. They also report on influenza vaccine coverage for all age groups. These reports are submitted to the IHS Division of Epidemiology and Disease Prevention on a quarterly basis.

3-27 Month Old Report

The 3 – 27 month report is designed to help programs ensure timely vaccination and identify children who may be falling behind so they can be brought up to date. The criteria listed below are used to monitor coverage in the following age groups:

Age Group	Vaccines	Age Group	Vaccines
3-4 months	<ul style="list-style-type: none"> • DTAP1 • IPV 1 • Hib 1 • Hepatitis B 1 • PCV1 • (ROTA1)[†] 	16-18 months	<ul style="list-style-type: none"> • DTAP 3 • IPV 2 • MMR 1 • Hib 3 • Hepatitis B 2 • Varicella • PCV4 • (ROTA3)[†]
5-6 Months	<ul style="list-style-type: none"> • DTAP 2 • IPV 2 • Hib 2 • Hepatitis B 2 • PCV2 • (ROTA2)[†] 	19-23 months	<ul style="list-style-type: none"> • DTAP 4 • IPV 3 • MMR 1 • Hib 3 • Hepatitis B 3 • Varicella • PCV4 • (ROTA3)[†]
7-15 months	<ul style="list-style-type: none"> • DTAP 3 • IPV 2 • Hib 2 • Hepatitis B 2 • PCV3 • (ROTA3)[†] 	24-27 months	<ul style="list-style-type: none"> • DTAP 4 • IPV 3 • MMR 1 • Hib 3 • Hepatitis B 3 • Varicella • PCV4 • (ROTA3)[†] • (Hepatitis A1)[†]


† Not included in Age appropriate immunization coverage calculations

Rotavirus vaccine (ROTA) and hepatitis A, though recommended, are not included in overall age appropriate coverage calculations. Data on ROTA and hepatitis A vaccine coverage are reported where available.

Summary

For FY 2011 Q4, 12 IHS Areas submitted immunization reports from IHS, Tribal, and Urban Indian health centers (I/T/U). This composite report provides information on the immunization status of 30,151 children 3-27 months old. Of these, 21,405 or 71% received all age-appropriate vaccinations compared to 71% (22,647/31,797) of children 3-27 months in the 3rd Quarter of FY 2011 [Figure 1]. Coverage by Area is included in the spreadsheet.

Figure 1



Two Year Old Report

The Two year old reports (children 19-35 months old) are used to monitor progress towards the GPRA childhood immunization indicator.


As of 2010, the 4:3:1:3:3:1 (4 DTaP, 3 IPV, 1 MMR, 3 Hib, 3 Hep B, 1 VAR) series is used to monitor coverage in this age group; in 2011, the measure will be expanded to include 4 doses of PCV (4:3:1:3:3:1:4). The HP 2020 goal is 80% with the 4:3:1:3:3:1:4 series, and 90% coverage with each individual vaccine in the series.

4:3:1:3:3:1 and 4:3:1:3:3:1:4 Series

In FY 2011 Q4, 12 IHS Areas completed reports. Overall 80% (17,948/22,579) of two year olds had completed the 4:3:1:3:3:1 vaccine series compared to 80% (19,593/24,388) in FY 2011 Q3 [Figure 2]. For the 4:3:1:3:3:1:4 series, coverage was 74% (16,618/22,579) among two year olds compared to 76% (18,444/24,388) in FY 2011 Q3. Coverage for both series by Area is included in the spreadsheet.

Figure 2

**4:3:1:3:3:1 and 4:3:1:3:3:1:4 Coverage
2 Year Olds
FY 2005 Q3 - FY 2011 Q4**



ADOLESCENT REPORT

IMPORTANT – CHANGES TO THE ADOLESCENT REPORT INCLUSION CRITERIA

The Adolescent Immunization Report was collected for the first time in FY 2008 Q1 and included adolescents manually deemed “active” by the facility. Starting with the FY 2010 Quarter 4 report, the adolescent report inclusion criteria changed. The report now includes all adolescents meeting the electronically-determined “Active Clinical” user definition – i.e. 2 primary care visits in the last 3 years. Because of this change, in this and all future adolescent reports we will include data as of FY 2010 Q4 rather than FY 2008 Q1. In general, the change in the report inclusion criteria resulted in an overall increase in the number of adolescents captured in the reports, and an overall decrease in coverage for most vaccines.


This report collects data on adolescents 13 – 17 years and is designed to monitor uptake of “recently” recommended adolescent vaccines (e.g. tetanus toxoid, reduced diphtheria toxoid and acellular pertussis [Tdap], meningococcal conjugate [MCV4], and human papillomavirus [HPV] vaccines) and monitor coverage in the adolescent population with the following recommended childhood vaccines: 1 dose of Td or Tdap, 3 doses of hepatitis B, 2 doses of MMR, and 2 doses of Varicella/Hx of chickenpox. A summary of immunization coverage for 13 year olds and 13 –17 year olds with these vaccines is included below.

Immunization Coverage with Tdap and MCV4 for 13 year olds, Males and Females

For FY 2011 Q4, the 12 IHS Areas combined reported on 15,675 thirteen year olds compared to 16,416 thirteen year olds in the FY 2011 Q3 report. For FY 2011 Q4, 84% (13,215/15,675) received 1 dose of Tdap vaccine, 87% (13,573/15,675) received 1 dose of Tdap/Td vaccine and 82% (12,777/15,675) had received 1 dose of MCV4 vaccine [Figure 3].

In FY 2011 Q4, coverage by Area ranged from 76% - 91% for Tdap, 79% - 92% for Tdap/Td, and 69% - 93% for MCV4. Coverage by Area is included in the spreadsheet.

Figure 3




Immunization Coverage for 13 – 17 year olds, Males and Females

For FY 2011 Q4, the 12 IHS Areas combined reported on 77,793 13 – 17 year olds compared to the FY 2011 Q3 report which included 82,277 13 – 17 year olds.

For FY 2011 Q4, 90% (70,365/77,793) received 3 doses of hepatitis B vaccine, 91% (70,416/77,793) received 2 doses of MMR vaccine, 81% (63,392/77,793) received 2 doses of Varicella vaccine or had a documented history of chickenpox, 83% (64,899/77,793) received 1 dose of Tdap vaccine, 88% (68,763/77,793) received 1 dose of Tdap or Td and 82% (64,007/77,793) received 1 dose of MCV4 vaccine. Coverage by Area is included in the spreadsheet [Figure 4].

Figure 4



HPV Coverage for 13 – 17 year olds, Females Only

For FY 2011 Q4, the 12 IHS Areas combined reported on 41,063 females 13 – 17 years of age compared to FY 2011 Q3 which included 44,391 females 13-17 years. For FY 2011 Q4, 75% (30,878/41,063) had received 1 dose of HPV, 63% (25,754/41,063) had received 2 doses of HPV and 49% (20,004/41,063) had received 3 doses of HPV [Figure 5].

For FY 2011 Q4, coverage by Area ranged from 61% - 88% for HPV1, 47% - 83% for HPV2, and 34% – 73% for HPV 3. Coverage by Area is included in the spreadsheet.

Figure 5

HPV Coverage for Females 13 - 17 years FY10 Q4 - FY11 Q4

