



February 23, 2022

**\*\*Emergency Use Instructions\*\***  
**CDC Issues Updated Guidance on COVID-19 mRNA Vaccine Dosing Interval**

**CDC Emergency Use Instructions<sup>1-3</sup>:**

On February 22, 2022 the CDC issued [Emergency Use Instructions](#) (EUI) which provide information about emergency use of the licensed COVID-19 vaccines by [Pfizer-BioNTech \(Comirnaty<sup>®</sup>\)](#) and [Moderna \(Spikevax<sup>®</sup>\)](#). **The CDC-issued EUI provide instructions and information for the use of these vaccines that are beyond the FDA-approved labeling.** Authority for the issuance of EUI is allowed under the [Pandemic and All Hazards Preparedness Reauthorization Act](#).

**Summary of Revisions to CDC Interim Clinical Considerations for Use of COVID-19 Vaccines<sup>1,4</sup>:**

Update of COVID-19 vaccine guidance to include addition of considerations for an 8-week interval between the first and second doses of a primary mRNA vaccine schedule **for certain persons ages 12-64 years of age:** (Note that this applies only to use of Spikevax<sup>®</sup> (Moderna) for people ages 18 years and older and Comirnaty<sup>®</sup> (Pfizer-BioNTech) for people ages 12 years and older).

Vaccine	First Dose	Second Dose	Booster Dose
<b>Pfizer-BioNTech</b> (ages 5-11 years)	Month 0	3 weeks after 1st dose	NA
<b>Pfizer-BioNTech</b> (ages 12 years and older)	Month 0	3-8 weeks after 1st dose	At least 5 months after 2nd dose
<b>Moderna</b> (ages 18 years and older)	Month 0	4-8 weeks after 1st dose	At least 5 months after 2nd dose
<b>Janssen</b> (ages 18 years and older)	Month 0	NA	At least 2 months after 1st dose

For a link to the original CDC table graphic, please click here: [https://www.cdc.gov/vaccines/covid-19/images/covid19-vac-schedule-chart-01.png?\\_=42169?noicon](https://www.cdc.gov/vaccines/covid-19/images/covid19-vac-schedule-chart-01.png?_=42169?noicon)

The originally recommended interval (3 weeks for Pfizer-BioNTech; 4 weeks for Moderna) between the first and second primary doses remains the recommended interval for:

- people who are moderately to severely immunocompromised;
- adults ages 65 years and older; and
- others who need early protection due to increased concern about community transmission or risk of severe disease.

**Rationale<sup>5</sup>:**

International data on risk of myocarditis, especially among the highest risk group (males ages 12-39 years of age post dose 2) along with international data on an extended primary series interval related to the risk of myocarditis and vaccine effectiveness suggest that an extended 8-week dosing interval between the 1<sup>st</sup> and 2<sup>nd</sup> dose of an mRNA COVID-19 primary vaccine series may;

- increase duration of vaccine protection
- reduce the small risk of myocarditis & pericarditis

## **Updated CDC mRNA COVID-19 Vaccine EUI Fact Sheets<sup>2-3</sup>:**

For Healthcare Providers:

- [Pfizer-BioNTech COVID-19 Vaccine EUI Fact Sheet](#)
- [Moderna COVID-19 Vaccine EUI Fact Sheet](#)

For Recipients and Caregivers:

- [Pfizer-BioNTech COVID-19 Vaccine EUI Fact Sheet](#)
- [Moderna COVID-19 Vaccine EUI Fact Sheet](#)

### **References:**

1. U.S. Centers for Disease Control and Prevention, Vaccine [Interim Clinical Considerations for Use of COVID-19 Vaccines](#). Accessed 2/23/22.
2. U.S. Centers for Disease Control and Prevention, [Emergency Use Instructions for Healthcare Providers](#): Pfizer-BioNTech COVID-19 vaccine for Primary, Additional, and/or Booster Doses. Accessed 2/23/22.
3. U.S. Centers for Disease Control and Prevention, [Emergency Use Instructions for Healthcare Providers](#): Moderna COVID-19 vaccine for Primary, Additional, and/or Booster Doses. Accessed 2/23/22.
4. U.S. Centers for Disease Control and Prevention, [Questions and Answers about Emergency Use Instructions](#). Accessed 2/22/22.
5. Advisory Committee on Immunization Practices, [Myocarditis and COVID-19 Vaccines Intervals](#), International Data & Policies, Danielle Moulia, Feb 4, 2022.