



**Indian Health Service  
National Pharmacy and Therapeutics Committee  
Formulary Brief: ACEi and ARB in Heart Failure  
-May 2017-**



---

---

**Background:**

In May 2017, the IHS National Pharmacy and Therapeutics Committee (NPTC) convened to discuss pharmacotherapy in heart failure (HF). Angiotensin converting enzyme inhibitors (ACEi) and angiotensin receptor blockers (ARB) have important roles in the management of heart failure and are considered a cornerstone in the treatment of heart failure with reduced ejection fraction (HFrEF)<sup>1,2</sup>. The NPTC last reviewed the ACEi and ARB classes (for hypertension) in August 2014 and, at that time, made no changes to the National Core Formulary (NCF). At that time, lisinopril and losartan were listed on the NCF as sole representatives in their respective medication classes. Based on the findings of this new review, **no changes were made to the NCF.**

**Discussion:**

*Angiotensin Converting Enzyme Inhibitors:*

The review focused heavily on HFrEF as ACEi and ARBs are most strongly recommended and show the greatest benefit in these patients. It is well established that ACEi reduce morbidity and mortality in HFrEF based on primary literature of trial data<sup>3,4</sup> but most trials have used enalapril. All ACEi are similar with regard to safety, dosing adjustments, monitoring and precautions<sup>5</sup>. A landmark meta-analysis published in 1995 concluded that there was an ACEi class effect in HF<sup>6</sup>. The study showed a significant reduction in mortality (OR=0.77; 95% CI: 0.67 to 0.88,  $p<0.001$ ), a reduction in the combined endpoint of mortality or hospitalization in HF (OR=0.65; 95% CI: 0.57 to 0.74,  $p<0.001$ ) and an equivalent effect across the ACEi class (no statistical heterogeneity was detected among the various agents).

Subsequent studies have suggested that differences among the ACEi class may exist. A meta-analysis published in 2016 reported enalapril to have the greatest hemodynamic effect but also the most adverse events. The meta-analysis noted that ramipril had the lowest incidence all-cause mortality and lisinopril had the highest all-cause mortality, comparatively<sup>7</sup>. Despite these findings, authors concluded that there was no statistical difference in any ACEi and that no conclusive recommendations could be made<sup>7</sup>. Another study published in 2016 also suggested heterogeneity among the ACEi class<sup>8</sup>. This study reported that ramipril had the highest probability of reducing death among ACEi. However, due to short-term durations and limited number of ramipril trials included in the analysis, the study did not demonstrate statistically significant results (i.e., benefit) for any ACEi.

*Angiotensin Receptor Blockers:*

Only candesartan and valsartan are FDA indicated in the treatment of HF. Losartan is the current NCF agent for the ARB class. Losartan does not have an FDA indication for HF but is routinely used off label for this indication. All ARBs are similar regarding safety, dosing adjustments, monitoring and precautions<sup>9</sup>. The outcomes data to support candesartan and valsartan are derived from well-known studies (CHARM and Val-HeFT, respectively).

In an effort to determine whether losartan had benefit over an ACEi in HF, the ELITE study was undertaken in 1997<sup>10</sup>. Losartan was compared against captopril and, as a secondary measure, demonstrated a non-statistically significant 32% reduction in death and/or hospitalization for HF. This trend was largely driven by a 46% decrease in all-cause mortality ( $p=0.035$ ). Because these results were secondary endpoints and thus not hypothesized a priori, the investigators attempted to replicate these results in the ELITE II study<sup>11</sup>. The ELITE II study concluded with no mortality benefit seen with losartan versus captopril. Interestingly, both ELITE studies used the 50mg doses of losartan; researchers suggested the dosing may have been suboptimal in this trial. Subsequently, the HEAAL trial was conducted to evaluate the dose-dependent effect of losartan (50mg vs. 150mg) in HF<sup>12</sup>. Losartan 150mg was found to be superior in the composite endpoint of all-cause mortality and HF admission (HR=0.90; 95% CI, 0.82 to 0.99;  $p=0.027$ ). Hypotension, hyperkalemia and renal impairment were all more common in the 150mg arm but did not lead to differences in discontinuation rates between study groups.

A subsequent study published in 2009 compared 4 ARBs (candesartan, irbesartan, losartan, valsartan) using VA data from 1996 to 2002 and concluded there are no statistical differences in mortality between the ARBs studied<sup>13</sup>.

Neither American nor European consensus guidelines recommend a specific ACEi or ARB over other agents within their respective drug classes<sup>1,2</sup>. Additionally, these guidelines recommend using ARBs only when ACEi are not tolerated.

### Findings:

The benefits of ACEi can generally be considered a class effect in HF<sub>rEF</sub> and no specific ACEi is favored among the ACEi indicated for HF (i.e., captopril, enalapril, fosinopril, lisinopril, perindopril, quinapril, ramipril, trandolapril). The strength and volume of evidence supporting the role of ARBs in HF<sub>rEF</sub>, when clinically indicated, is less clear (vs. ACEi) and only three ARBs are generally accepted for use in HF management (candesartan, losartan, valsartan). While candesartan and valsartan have FDA indications for HF treatment, losartan remains commonly prescribed based on clinical trial data supporting its use in HF<sub>rEF</sub><sup>12</sup>. When losartan is used, clinicians should target a 150mg daily dose in HF<sub>rEF</sub> patients<sup>12</sup>.

---

*If you have any questions regarding this document, please contact the NPTC at [IHSNPTC1@ihs.gov](mailto:IHSNPTC1@ihs.gov). For more information about the NPTC, please visit the [NPTC website](#).*

### References:

1. Yancy CW, Jessup M, Bozkurt B, et al. [2013 ACCF/AHA Guideline for the Management of Heart Failure: A report of the ACCF/AHA task force on practice guidelines](#). Circulation. 2013;128(16).
2. Ponikowski P, Voors AA, Anker SD, et al. [2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure](#). Eur Heart J. 2016; 37(27):2129–2200.
3. The CONSENSUS Trial Study Group. [Effects of enalapril on mortality in severe congestive heart failure: results of the Cooperative North Scandinavian Enalapril Survival Study \(CONSENSUS\)](#). NEJM 1987; 316:1429–35.
4. The SOLVD Investigators. [Effect of enalapril on survival in patients with reduced left ventricular ejection fraction and congestive heart failure](#). NEJM. 1991; 325(5):293–302.
5. Official reprint from UpToDate @ www.uptodate.com ©2017 UpToDate @ [ACE inhibitors in heart failure with reduced ejection fraction: Therapeutic use](#). (2017).
6. Garg R, Yusuf S. [Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure](#). JAMA. 1995; 273:1450–6.
7. Sun W, Zhang H, Guo J, et al. [Comparison of the Efficacy and Safety of Different ACE Inhibitors in Patients with Chronic Heart Failure: A PRISMA-Compliant Network Meta-Analysis](#). Medicine (Baltimore) [Internet]. 2016; 95(6):e2554.
8. Guo W-Q, Li L. [Angiotensin converting enzyme inhibitors for heart failure with reduced ejection fraction or left ventricular dysfunction: A complementary network meta-analyses](#). Int J Cardiol [Internet]. 2016; 214:10–12.
9. Jessup M. Official reprint from UpToDate @ www.uptodate.com ©2017 UpToDate @ [Use of angiotensin II receptor blocker and neprilysin inhibitor in heart failure with reduced ejection fraction](#). 2017;1.
10. Pitt B, Segal R, Martinez FA, et al. [Randomized trial of losartan versus captopril in patients over 65 with heart failure: Evaluation of Losartan in Elderly Study \(ELITE\)](#). Lancet 1997; 349:747-52.
11. Pitt B, Poole-Wilson PA, Segal R, et al. [Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomized trial: the Losartan Heart Failure Survival Study ELITE II](#). Lancet 2000; 355:1582-7.
12. Konstam MA, Neaton JD, Dickstein K, et al. [Effects of high-dose versus low-dose losartan on clinical outcomes in patients with heart failure \(HEAAL study\): a randomized, double-blind trial](#). Lancet 2009; 374:1840-8.
13. Desai RJ, Ashton CM, Deswal A, et al. [Comparative effectiveness of individual angiotensin receptor blockers on risk of mortality in patients with chronic heart failure](#). Pharmacoepidemiol Drug Saf. 2012; 21(3):233–40.