




Influenza Planning 2015-2016

CAPT Thomas Weiser, MD, MPH
Medical Epidemiologist
Portland Area Indian Health Service/
Northwest Tribal Epidemiology Center

Key Points for Area Immunizations Coordinators/Planners

- Use area-level data sources to determine historic flu vaccine coverage and to monitor progress
 - IIAS, CRS, NIRS
- Advocate for early vaccination
 - Address concerns about supply availability, waning immunity.
- Anticipate adverse media coverage and develop messages to mitigate the effects
 - Decreased coverage because of vaccine mis-match last year led to increased influenza B later in the flu season.

IPC Care Model



Rationale

Table 1. Pooled Average Vaccine Effectiveness (VE)


Age range (yrs.)	Average VE	Range
0.5–4	52%	39%-67%
5–19	50.25%	46%-59%
20–64	50%	46%-52%
≥65	37.5%	32%-43%

(Adapted from Foppa, *et al.* Vaccine, 2015)¹

“An annual seasonal flu vaccine ... is the best way to reduce the chances that you will get seasonal flu and spread it to others. When more people get vaccinated against the flu, less flu can spread through that community.”

CDC. <http://www.cdc.gov/flu/protect/keyfacts.htm>

Critical vaccination coverage as a function of vaccine effectiveness for given level of R_0




Average Seasonal flu: $R_0=1.3$

1918 Pandemic flu: $R_0=2.0$

(Adapted from Plans-Rubio, *et al*, 2012)³


Critical vaccination coverage as a function of vaccine effectiveness for given level of R_0




Critical vaccine coverage 0.5-64 years - - - - -
Critical vaccine coverage ≥ 65 years - - - - -

(Adapted from Plans-Rubio, *et al*, 2012)³

Cumulative Percent of Active User Population Receiving Influenza Immunization and ILI Activity Portland Area IHS 2014-2015 Season




Weekly count of influenza vaccine doses given in Portland Area IHS for the 2014-15 influenza season



Strategies to increase the uptake of influenza vaccine in the Portland Area IHS


- 1. Starting sooner:** vaccinate as many people as possible as soon as the vaccine arrives; even starting 1 or 2 weeks sooner would increase the early uptake of vaccine.
- 2. Sustain maximum vaccination rate longer:** try to keep vaccinating at the same high rate in October for four more weeks, until the end of November. The advantages of this strategy is there is no expectation of starting earlier, which is dependent on vaccine supplies, and the clinics (systems) already have a proven capacity to vaccinate at this rate.
- 3. Increase weekly vaccination uptake by a defined percentage (e.g, 25%):** requires that the clinics/systems adapt to provide more vaccinations/week than last year. This could include adding additional evening or weekend clinics or adding community-based vaccine sites/efforts. And it requires the increased effort to be maintained throughout the flu season, though once the “goal” is achieved the effort could be tapered off.
- 4. Combination Strategies:** would use two or more of these strategies in combination.

Projected cumulative influenza immunization rates using three single strategies compared to current practice.



Minimum herd immunity threshold to be reached by 11/30/2015 is shown in red. All three strategies are projected to show increased coverage but no single strategy will reach the goal of 50% before ILI activity begins nor would they reach HP2020 goal of 70%

Projected cumulative influenza immunization rates using three combination strategies compared to current practice



Minimum herd immunity threshold to be reached by 11/30/2015 is shown in red. All three strategies could meet/exceed the goal of 50% before ILI activity begins.

Driver Diagram for Improving Influenza Vaccine Coverage

Strategy (Change Concept)	Primary Drivers	Secondary Drivers	Constraints
<p>Start vaccinating sooner</p>	<p>Clinic Readiness</p>	<ul style="list-style-type: none"> • Pre-scheduled walk-in flu vaccine clinics • Pharmacists, Mas and nurses trained and ready to vaccinate • All necessary supplies in place prior to arrival of vaccines (gloves, syringes, needles, alcohol wipes, etc) 	<p>Highly dependent on timely vaccine supply delivery to clinic</p>
	<p>Community Readiness</p>	<ul style="list-style-type: none"> • Pre-placed articles/ads in local newspapers about when flu vaccines will be given, benefits of flu vaccines, etc • Messaging throughout the community- posters, brochures, PSAs, video-messages, Social Media, radio, etc • Community-based vaccine days/sites pre-planned 	


Strategy (Change Concept)	Primary Drivers	Secondary Drivers	Constraints
<p>Increase weekly number of vaccines given per week by some percent (e.g., by 25%)</p>	<p>Clinical systems change to increase capacity</p> <p>Community Demand or Acceptance</p>	<ul style="list-style-type: none"> • Remove barriers to getting flu vaccine (standing orders, walk-in clinics, offering to all patients, etc) • Provide multiple types of vaccine (e.g., live attenuated, preservative free, high-dose) • Providers educated and committed to providing flu vaccine to all patients • Vaccinate providers/staff • Create new vaccination venues – evening/weekend, community-based clinics • Develop/repeat messaging strategies • Anticipate and provide information specific to issues that may develop (vaccine mis-match, adverse events, reported “severity” of the circulating flu strain, special populations). 	<ul style="list-style-type: none"> • System must increase its daily capacity to give vaccines (staff must work harder than previous years) • Staff reluctance to promote vaccine or reluctance to receive their own flu vaccine • Insufficient staff to provide evening/weekend vaccination clinics • Mistrust of IHS/CDC • Negative media messages

Recommendations

IHS Areas should consider the following:

- Review local influenza policies and practices
- Review data on influenza immunization levels in prior years
- Set goals to achieve immunization levels that approach the IHS goal of 70% coverage for all aged 6 months and older.
- Consider adopting more than one single strategy
- Identify the primary and secondary drivers of flu vaccine uptake and adopt new policies and practices aligned with those drivers.
- At the clinic level:
 - Engage ALL staff in efforts to receive and provide influenza immunizations.
 - Engage patients through media/outreach materials (posters, postcards, PSAs and articles) and open communication.

Resources



- [NPAIHB Breaking News 2015-2016 Flu Season](#)
- www.cdc.gov/flu
- <https://www.ihs.gov/Flu/>
- www.facebook.com/IHSHDPDP
- www.flu.gov
- [Wes Study Flu Video](#)
- [More CDC Resources](#)

A digital widget for finding flu clinics. It has a blue background with white and orange text. The title is "Flu Vaccine Finder" with a sub-header "Find flu clinics near you". Below that, it says "Everyone six months of age and older needs a flu vaccine." At the bottom, it says "powered by HealthMap" and includes buttons for "VISIT FLU.GOV" and "SHARE THIS WIDGET".

Flu Vaccine Finder Find flu clinics near you

Everyone six months of age and older needs a flu vaccine.

powered by HealthMap

VISIT FLU.GOV SHARE THIS WIDGET

A vertical poster with a yellow top section and a white middle section. The top section says "Test Your Knowledge" with the CDC logo. The middle section features a circular graphic of various shoes arranged around a central circle divided into four quadrants (black, white, red, yellow). The text "Protect the Circle of Life" is written in a curve above the shoes. The bottom section says "Please Get the Flu Vaccine".

Test Your Knowledge CDC

Protect the Circle of Life

Please Get the Flu Vaccine

A vertical poster with a white background and a yellow bottom section. The title is "2015-2016 Influenza Vaccine Information" with a sub-header "Which Flu Shot should I get?". Below that is a graphic with the text "Life is a delicate balance" and a red feather. At the bottom, it says "Save Lives, Immunize!".

2015-2016 Influenza Vaccine Information

➤ Which Flu Shot should I get?

Life is a delicate balance

Your flu vaccine protects me. My flu vaccine protects you. When you get your flu vaccine, you can keep yourself healthy and keep from spreading the flu. Protect the circle of life. For more information, visit <http://www.flu.gov> or <http://www.cdc.gov/flu>

Save Lives, Immunize!

References

1. Deaths averted by influenza vaccination in the U.S. during the seasons 2005/06 through 2013/14. I Foppa, P Cheng, S Reynolds, D Shay, C Carias, J Bresee, I Kim, M Gambhir, A Fry. Article in Press, Vaccine (2015), <http://dx.doi.org/10.1016/j.vaccine.2015.02.042>
2. Estimates of the reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic review of the literature. M Biggerstaff, S Cauchemez, Carrie Reed, M Gambhir, Lyn Finelli. BMC Infectious Diseases (2014) 14:480 <http://www.biomedcentral.com/1471-2334/14/480>
3. The vaccination coverage required to establish herd immunity against influenza viruses. P Plans-Rubió. Preventive Medicine (2012) 55:72–77



Contact Information:
CAPT Thomas Weiser, MD, MPH
tweiser@npaihb.org
thomas.weiser@ihs.gov
(503) 416-3298 (Office)
(503) 927-4467 (Cell)



Northwest Portland Area
Indian Health Board
Indian Leadership for Indian Health