# The Impact of Caregiver Alcoholism on Youth and Families

Shawn S. Sidhu, M.D., F.A.P.A.

#### Disclosure

Dr. Sidhu receives royalties from the American Psychiatric Association for writing continuing medical education questions for the journal *FOCUS* 

## Objectives

At the end of this presentation, participants will be able to:

- Identify key statistics regarding the prevalence of alcoholism in adults and youth.
- Screen youth in homes with caregiver alcohol use for a myriad of concerns, including cognitive and physiological development.
- 3. Perform a 20-Item Questionnaire from Al-Anon which screens those who may have been impacted by growing up in a home with an alcoholic.

#### Resources

National Institutes of Alcohol and Alcohol Abuse (<a href="http://www.niaaa.nih.gov/">http://www.niaaa.nih.gov/</a>)

National Council on Alcoholism and Drug Dependence, Inc. (<a href="https://ncadd.org/about-addiction/alcohol-drugs-and-crime">https://ncadd.org/about-addiction/alcohol-drugs-and-crime</a>)

#### OUTLINE

I) Epidemiology

- II) Impact of Caregiver Alcoholism on Children and Families
  - I) Physical
  - II) Emotional
  - III) Cognitive
  - IV) Social
- III) Solutions/Treatments

Much of the research presented here is in the form of association, and <u>not</u> necessarily causation

#### Alcoholic Parents

Alcoholic parents may well have experienced traumatic childhoods themselves, as child victimization is a significant risk factor for the development of alcohol and drug problems in adulthood, and this risk is especially high for females but also present for males

For people with traumatic childhoods, alcohol/drugs are most commonly used:

- -As a coping mechanism
- -To reduce feelings of isolation/loneliness
- -Self-medication of mental health issues
- -To gain control, improve self-esteem and self-empowerment
- -As a form of self-destructive behavior

Epidemiology of Alcohol Use in Adults and Youth

# Adult Epidemiology

25% of people 18 and older admit to binge drinking recently

16.6 million adults in the U.S. have an Alcohol Use Disorder or AUD (9.4% of men and 4.7% of women)

3<sup>rd</sup> leading preventable cause of death (88,000 annually)



20% of adult Americans have lived with an alcoholic relative while growing up

#### **Youth Statistics**

700,000 teens (3.2 % of females and 2.5% of males) have an AUD

35% of 15 year olds have abused alcohol, and 23% of 12-20 year olds report drinking within the past month (23% males, 22.5% females)

Children of alcoholics are four times more likely than other children to become alcoholics themselves

Impact of Caregiver Alcoholism on Physical Health

#### QUESTION 1

Which of the following types of cancer has been linked to alcohol consumption that exceeds moderation (> 2 drinks per day)?

- A) Brain
- B) Bone
- C) Blood (Leukemia/Lymphoma)
- D) Breast
- E) Lung

#### Impact of Alcohol on Physical Health

Nearly 50% of all liver disease deaths (70,000) annually are alcohol related

Alcohol consumption that exceeds moderation (>2 drinks per day) increases risk of oral, esophageal, pharyngeal, colorectal, liver, pancreatic, and breast cancer, and very heavy consumption (4 drinks or greater daily) can also increase risk for endometrial, gallbladder, ovarian, prostate, and small intestine cancer.

While < 1% (0.7%) of children are born with Fetal Alcohol Syndrome, 5% are born with Fetal Alcohol Spectrum Disorders (FASD). According to the CDC, 7 to 8% of pregnant mothers continue to drink alcohol despite massive education campaigns aimed at this problem. FASD is associated with hyperactivity, impulsivity, growth retardation, facial anomalies, and cognitive problems

http://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/alcohol-facts-and-statistics

### Impact of Alcohol on Physical Health of Children

Additionally, mothers who are abusing alcohol may be more likely to be malnourished and to not receive appropriate prenatal care during their pregnancies, further placing their children at risk for birth defects

Given the high rate of neglect in children of alcoholics, it is highly likely that things like yearly physicals, basic dentition, and eye exams for corrected vision will go forgotten in some alcoholic households

#### QUESTION 2

Alcohol increases which of the following phases of sleep in the second half of the evening?

- A) Stage I
- B) Stage II
- C) Stage III
- D) Stage IV
- E) Beta-Wave

#### Alcohol and Sleep

30% of people with difficulty sleeping use alcohol as a remedy


Dose-dependent (1 drink regularly +/-, approaching 3-4 drinks per night negative impact)

Improves initiation and 1<sup>st</sup> half of sleep, greatly disrupts 2<sup>nd</sup> half

REM and Light Sleep (Stage I) follow the same pattern:

- -Decrease in 1<sup>st</sup> half of the night
- -Rebound increase well beyond baseline in 2<sup>nd</sup> half of night

Result: 2<sup>nd</sup> half of the night is filled with increased awakenings and non-restorative sleep



# Impact of Parental Alcohol Use on the Health of Children

#### DWI/DUI:

> 1 million arrests annually

3<sup>rd</sup> most common crime in the United States

#1 cause of death, injury, and disability of those under 21

40% of traffic fatalities alcohol-related

Daily: 36 MVA deaths and 700 injuries due to alcohol

\*Not only places children at risk for great harm, but also places children at risk for having their caretakers injured/killed, arrested, and without drivers licenses for transportation

#### Question 3

Which of the following developmental problems is seen most frequently in children living with an alcoholic parent?

- (A) Cognitive
- (B) Behavioral
- (C) Psychological
- (D) Social
- (E) Physical (health)

Although all of these developmental issues may be present in children of alcoholics, cognitive problems are observed most frequently



Children of alcoholics have been found to have:

-Lower full-scale IQ scores

-Lower score son performance tests (abstract and conceptual reasoning)

-Lower verbal scores

Even in educated alcoholic families with parents living in the home, children were found to have:

- -Lower IQ
- -Lower arithmetic scores
- -Lower reading scores
- -Lower verbal scores

\*Lower does not mean below normed values but rather in the low-normal range

Even children of alcoholics who are considered to perform differently from their peers academically, still:

- -They often underestimate their own competence (self-esteem)
- -Their mothers often underrate the abilities of their child (parental pessimism)
- \*These perceptions affect motivation, self-esteem, and future performance

School performance may be a more practical measure of cognitive health. Children of alcoholics demonstrate:

#### -Academic problems in school

- -Repeating/failing grades
- -Failing to graduate form high school
- -Requiring referrals to school mental health professionals
- -Truancy
- -School behavioral problems

\*While cognitive issues may partially explain this, these cognitive issues are compounded by <u>decreased motivation</u> and <u>stress in the home environment</u>

# Poor School Performance Directly Linked to Decreased Earning Potential

Impact on Social Issues Involving Children

#### Impact on Social Issues Involving Children

Households with adult alcoholics have a higher rate of :

- -Divorce
- -Parental anxiety
- -Parental affective disorders
- -More frequent/undesirable changes in the family/life situation

#### Impact on Social Issues Involving Children

#### Families of alcoholics have:

- -Lower levels of cohesion
- -Lower levels of expressiveness
- -Lower levels of independence
- -Increased family conflict

Children of alcoholics are also at greater risk for teen pregnancy (1)

## Alcoholism and Poverty

Longer durations of poverty and involuntary unemployment significantly predict heavy drinking, independent of gender, age, race/ethnicity, marital status, prior heavy drinking, and present socioeconomic status (1)

Neighborhood poverty is also a significant predictor of alcohol consumption (2)

#### Question 4

What percentage of violent crimes (in a sample of nearly 2 million convicted offenders in jail) involve alcohol?

- A) 1%
- B) 10%
- C) 25%
- D) 40%
- E) 70%

#### Parental Alcohol Use and Violence/Abuse

Alcohol is a factor in 40% of all violent crimes (Department of Justice), including closer to 50% of homicides and assaults

Associated more than any other drug with murder, rape, assault, child, and spousal abuse

Domestic Violence: Up to 69% of incidents involve alcohol. This includes domestic violence versus spouses/significant others and other family members. Children who witness domestic violence against their maternal caregiver are 50% more likely to abuse alcohol and/or drugs.

#### Parental Alcohol Use and Violence/Abuse

40% of child victimizers/abusers report drinking at the time of the abuse, and 50% of these abusers had been consuming alcohol for 6 or more hours preceding the abuse

Children of alcohol/drug-abusing parents are three times more likely to be abused and four times more likely to be neglected than peers

The overall level of stress in alcoholic homes can make these parents especially at risk for lashing out at their family members, and the presence of alcohol decreases the inhibition to engage in violence

#### Juvenile Crime and Alcohol/Drugs

80% of children arrested in state juvenile justice systems are:

- -Under the influence of alcohol/drugs while committing crimes
- -Test positive for drugs
- -Are arrested for committing an alcohol or drug offense
- -Admit to having a substance abuse or addiction problem

\*While 1,900,000 out of 2,400,000 juveniles had substance abuse involvement, less than 70,000 had received treatment

# Impact on Emotional and Behavioral Development in Children

#### Mental Health

#### Children of Alcoholics have:

- -Higher rates of depression (more depressive affect, more frequent and extreme depressive episodes than peers)
- -Higher rates of anxiety
- -More symptoms of generalized stress, including low self-esteem
- -A pervasive sense of a lack of control

\*Some of these symptoms can reverse when parents stop drinking

### Behavioral Issues

Children from homes with alcoholic parents demonstrate increased:

- -Conduct-disordered behavior:
  - -Lying
  - -Stealing
  - -Fighting
  - -Truancy/School Behavioral Problems

- -Hyperactivity
- -Impulsivity

### **Emotional Issues**

Many children find themselves in a bind between:

- -Not having their own needs met
- -Forced parentification at an early age, especially older children
- -Leading to the loss of a typical childhood
- -The inability to form close friendships given inconsistency
- -The inability to perform well in school given inconsistency

### **Emotional Issues**

Needs not met: lack of empathy, low self-esteem

<u>Forced parentification:</u> lifelong caretaker, avoidance of responsibility, desire to not have their own children/families

Loss of a typical childhood: grief/depression

The inability to form close friendships given inconsistency: social isolation and the inability to develop social skills

The inability to perform well in school given inconsistency: decreased earning potential, increased unemployment, decreased self esteem

### **Emotional Issues**

Common emotions in children of alcoholics:

*Guilt*: unable to care for parent/family, view themselves as the cause

Anxiety: worry constantly about abuse, health issues, financial/housing

**Embarrassment**: of family behavior in public or of clothing/housing

*Fear of Intimacy*: inability to trust people via repeated disappointments

**Confusion**: via inconsistent parental behavior and environment

<u>Anger/Frustration</u>: can also include anger and enablers in the family

<u>Depression</u>: sense of pervasive despair and hopeless/helpless

### 20 Item Questionnaire from Al-Anon

#### Did you grow up with a problem drinker? If so:

- 1) Do you constantly seek approval and affirmation?
- 2) Do you fail to recognize your accomplishments?
- 3) Do you fear criticism?
- 4) Do you overextend yourself?
- 5) Have you had problems with your own compulsive behavior?
- 6) Do you have a need for perfection?
- 7) Are you uneasy when your life is going smoothly, continually anticipating problems?
- 8) Do you feel more alive in the midst of a crisis?
- 9) Do you still feel responsible for others, as you did for the problem drinker?
- 10) Do you care for others easily, yet find it difficult to care for yourself?

### 20 Item Questionnaire from Al-Anon

- 11) Do you isolate yourself from other people?
- 12) Do you respond with fear to authority figures and angry people?
- 13) Do you feel that individuals and society in general are taking advantage of you?
- 14) Do you have trouble with intimate relationships?
- 15) Do you confuse pity with love, as you did with the problem drinker?
- 16) Do you attract and/or seek people who are compulsive or abusive?
- 17) Do you cling to unhealthy relationships because you are afraid of being alone?
- 18) Do you often mistrust your own feelings and the expressed feelings of others?
- 19) Do you find it difficult to identify and express your own emotions?
- 20) Do you think someone's drinking may have affected you?

Solutions/Treatments

## Resiliency

What contributes to resiliency in children of alcoholics?

Those who go on to function well as adults have:

- -Other adults and people who were able to give them positive attention
- -Average to strong communication skills
- -Average intelligence or beyond
- -A <u>caring</u> attitude
- -A desire to achieve
- -A belief in self-help

# Resiliency

Children of alcoholic parents are less likely to become alcoholics themselves when they have a parent who:

- -ls consistent
- -Sets limits
- -Follows through on plans
- -Maintains daily rituals such as mealtimes
- -Maintains holiday rituals

# Solutions/Treatment

Be sure to refer <u>parents</u> to treatment

Typically, substance abuse treatment is divided into:

- -Acute inpatient detoxification
- -Long-term residential treatment
- -Ongoing outpatient therapy (Motivational Interviewing, Harm Reduction, 12 Step Groups, Faith Groups, etc.)
- -Ongoing outpatient med-assisted treatment (Naltrexone, Acamprosate, Naloxone, Methadone, Suboxone)

# Solutions/Treatment

Evaluate children for mental health, behavioral, cognitive, and/or emotional issues

Consider <u>family therapy</u> as substance abuse tends to have a profound impact on the entire family

<u>Identify additional supports for the child</u>, including extended family, friends, or other community supports (youth group leaders, spiritual leaders, big brothers/sisters, mentors, teachers)

# Solutions/Treatment

In the case of severe ongoing abuse and neglect, a <u>child protective services and/or</u> <u>tribal social services report</u> will need to be made

<u>Alternative living arrangements</u> can be made via other caring relatives and/or therapeutic foster care

Highly important to <u>continue screening</u> children for the development of substance use disorders of their own and school failure as well

Questions?

ssidhu@salud.unm.edu